

8. REFERENCES

- *Abedin Z, Cook RC, Jr, Milberg RM. 1980. Cardiac toxicity of perchloroethylene (a dry cleaning agent). *South Med J* 73:1081-1083.
- *Abrahamsson K, Ekdahl A, Cohen J, et al. 1995. Marine algae-a source of trichloroethylene and perchloroethylene. *Limnology and Oceanography* 40(7): 1321-1326.
- *ACGIH. 1991. Documentation of the Threshold Limit Values and Biological Exposure Indices. Sixth Edition. American Conference of Governmental Industrial Hygienists. Cincinnati, OH.
- *ACGIH. 1995. Threshold limit values and biological exposure indices for 1995 - 1996. American Conference of Governmental Industrial Hygienists. Cincinnati, OH.
- *Aggazzotti G, Fantuzzi G, Predieri G, et al. 1994a. Indoor exposure to perchloroethylene (PCE) in individuals living with dry cleaning workers. *Sci Total Environ* 156:133-137.
- *Aggazzotti G, Fantuzzi G, Righi E, et al. 1994b. Occupational and environmental exposure to perchloroethylene (PCE) in dry cleaners and their family members. *Arch Environ Health* 49(6):487-493.
- *Ahlborg G Jr. 1990. Pregnancy outcome among women working in laundries and dry cleaning shops using tetrachloroethylene. *Am J Ind Med* 17:567-575.
- *Albee RR, Mattsson JL, Bradley GJ, et al. 1991. Acute neurophysiologic effects of 1,1,2,2-tetrachloroethylene in rats. Sponsored by Halogenated Solvents Industry Alliance. OTS0540058. 88-920003407.
- *Alden CL. 1986. A review of unique male rat hydrocarbon nephropathy. Symposium. *Toxicol Pathol* 14(1):109-111.
- *Altmann L, Bottger A, Wiegand H. 1990. Neurophysiological and psychophysical measurements reveal effects of acute low-level organic solvent exposure in humans. *Int Arch Occup Environ Health* 62:493-499.
- *Altmann L, Florian Neuhann H, Kramer U, et al. 1995. Neurobehavioral and neurophysiological outcome of chronic low-level tetrachloroethene exposure measured in neighborhoods of dry cleaning shops. *Environmental Research* 69:83-89.
- *Altmann L, Wiegand H, Bottger A, et al. 1992. Neurobehavioral and neurophysiological outcomes of acute repeated perchloroethylene exposure. *Applied Psychology: An International Review* 41(3):269-279.
- *Anders MW, Lash L, Dekant W, et al. 1988. Biosynthesis and biotransformation of glutathione S-conjugates to toxic metabolites. *CRC Crit Rev Toxicol* 18:311-341.

8. REFERENCES

- Anderson DC, Jones SG. 1984. Fate of organic liquids spilled on soil. In: Hazard Wastes Environ Emerg; Manage Prev Cleanup Control Natl Conf Exhib, 384-388.
- Andersen ME, Krishnan K. 1994. Relating *in vitro* to *in vivo* exposures with physiologically based tissue dosimetry and tissue response models. In: Salem H, ed. Animal test alternatives: Refinement, reduction, replacement. New York, NY: Marcel Dekker, Inc., 9-25.
- *Andrasik I, Cloutet D. 1990. Monitoring solvent vapors in dry cleaning plants. International Fabricare Institute Focus on Drycleaning 14(3):1-8.
- *Anttila A, Pukkala E, Sallmen M, et al. 1995. Cancer incidence among Finnish workers exposed to halogenated hydrocarbons. J Occup Environ Med 37(7):797-806.
- *Antoine SR, DeLeon IR, O'Dell-Smith RM. 1986. Environmentally significant volatile organic pollutants in human blood. Bull Environ Contam Toxicol 36:364-371.
- Antti-Poika M. 1982. Overall prognosis of patients with diagnosed chronic organic solvent intoxication. Int Arch Occup Environ Health 5 1: 127- 138.
- Aoki A, Suzaki H, Kawabata Y, et al. 1994. Effect of perchloroethylene inhalation on nasal mucosa in mice. Eur Arch Otorhinolaryngol 25 1:361-365.
- *APHA. 1992. Standard methods for the examination of water and wastewater, 18th ed. American Public Health Association, American Water Works Association and Pollution Control Federation, Washington, DC, 6-42-6-47.
- *Aranyi C, O'Shea WJ, Graham JA, et al. 1986. The effects of inhalation of organic chemical air contaminants on murine lung host defenses. Fundam Appl Toxicol 6:713-720.
- *Aschengrau A, Ozonoff D, Paulu C, et al. 1993. Cancer risk and tetrachloroethylene-contaminated drinking water in Massachusetts. Archives of Environmental Health 48(5):284-292.
- Askergen A, et al. 1981. Studies on kidney function in subjects exposed to organic solvents. I. Excretion of albumin and β_2 -microglobulin in the urine. Acta Med Stand 209:479-483.
- ASTM. 1978. Compilation of odor and taste threshold values data. Philadelphia, PA: American Society for Testing and Materials, 155.
- *Atkinson R. 1985. Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds under atmospheric conditions. Chem Rev 85:130-132,193.
- *Atkinson R, Carter WPL. 1984. Kinetics and mechanisms of the gas-phase reactions of ozone with organic compounds under atmospheric conditions. Chem Rev 84:437-470.
- *ATSDR. 1989. Agency for Toxic Substances and Disease Registry. Part V. Federal Register 54:37619-37633.
- *ATSDR. 1993. Toxicological profile for trichloroethylene. U.S. Department of Health & Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry, TP-92/18.

8. REFERENCES

- Axelsson O. 1986. Epidemiological studies of workers with exposure to tri- and tetrachloroethylene. *Dev Toxicol Environ Sci* 12:233-230.
- *Bagnell PC, Ellenberger HA. 1977. Obstructive jaundice due to a chlorinated hydrocarbon in breast milk. *Can Med Assoc J* 117:1047-1048.
- *Barbash J, Roberts PV. 1986. Volatile organic chemical contamination of groundwater resources in the U.S. *J Water Pollut Control Fed* 58:343-348.
- *Barber LB II, Thurman EM, Schroeder MP. 1988. Long-term fate of organic micropollutants in sewage-contaminated groundwater. *Environ Sci Technol* 22:205-211.
- *Barkley J, Bunch J, Bursley JT, et al. 1980. Gas chromatography/mass spectrometry computer analysis of volatile halogenated hydrocarbons in man and his environment. A multimedia environmental study. *Biomed Mass Spectrom* 7:139-147.
- Barnes D, Dourson M. 1988. Reference dose (RfD): Description and use in health risk assessments. *Regul Toxicol Pharmacol* 8:471-486.
- *Barrows ME, Petrocelli SR, Macek KJ, et al. 1980. Bioconcentration and elimination of selected water pollutants by bluegill sunfish (*Lepomis macrochirus*). In: Haque R, ed., Dynamics, exposure and hazard assessment of toxic chemicals. Ann Arbor, MI: Ann Arbor Science, 379-392.
- *Bartels MJ. 1994. Quantitation of the tetrachloroethylene metabolite N-acetyl-S-(trichlorovinyl)cysteine in rat urine via negative ion chemical ionization gas chromatography/tandem mass spectrometry. *Biol Mass Spectrom* 23:689-694.
- *Bartsch H, Malaveille C, Barbin A, et al. 1979. Mutagenic and alkylating metabolites of haloethylenes, chlorobutadienes, and dichlorobutenes produced by rodent or human liver tissues. *Arch Toxicol* 41:249-277.
- *Bayer CW, Black MS. 1987. Capillary chromatographic analysis of volatile organic compounds in the indoor environment. *J Chromatogr Sci* 25(2):60-64.
- *Baykut G, Voigt A. 1992. Spray extraction of volatile organic compounds from aqueous systems into the gas phase for gas chromatography/mass spectrometry. *Anal Chem* 64:677-681.
- Bazylewicz-Walczak B, Marszal-Wisniewska M. 1986. Evaluation of mental effects of chronic tetrachloroethylene exposure. *Med Pr* 37:315-320.
- *Bellar TA, Lichtenberg JJ, Kroner RC. 1974. The occurrence of organohalides in chlorinated drinking waters. *Journal of American Water Works Association* 66:703-706.
- *Bentley P, Calder I, Elcombe C, et al. 1993. Hepatic peroxisome proliferation in rodents and its significance for humans. *Food Chemical Toxicology* 31(11):857-907.
- *Bergamaschi E, Mutti A, Bocchi MC, et al. 1992. Rat model of perchloroethylene-induced renal dysfunctions. *Environmental Research* 59:427-439.

8. REFERENCES

- *Berman E, Schlicht M, Moser VC, et al. 1995. A multidisciplinary approach to toxicological screening: I. Systemic toxicity. *J Tox Environ Health* 45:127-143.
- Bernard A, Lauwerys R. 1989. Epidemiological application of early markers of nephrotoxicity. *Toxicol Lett* 46:293-306.
- *Birner G, Richling C, Henschler D, et al. 1994. Metabolism of tetrachloroethylene in rats: identification of N-(dichloroacetyl)-L-lysine and N-(trichloroacetyl)-L-lysine as protein adducts. *Chem Res Toxicol* 7:724-732.
- *Birner G, Rutkowska A, Dekant W. 1996. *N*-Acetyl-*S*-(1,2,2-trichlorovinyl)-L-cysteine and 2,2,2-trichloroethanol: Two novel metabolites of tetrachloroethene in humans after occupational exposure. *Drug Metabolism and Disposition* 24(1):41-48.
- *Blair A, Decoufle P, Grauman D. 1979. Causes of death among laundry and dry cleaning workers. *Am J Public Health* 69:508-511.
- *Blair A, Stewart PA, Tolbert PE, et al. 1990. Cancer and other causes of death among a cohort of dry cleaners. *Br J Ind Med* 47:162-168.
- *Blattner WA, Strober W, Muchmore AV, et al. 1976. Familial chronic lymphocytic leukemia: immunologic and cellular characterization. *Ann Intern Med* 84:554-557.
- *Boekhold AJ, Van Der Schee HA, Kaandorp BH. 1989. [Rapid gas-chromatographic determination of trichloroethylene and/or tetrachloroethylene in lettuce by direct head-space analysis]. *Z Lebensm Unters Forsch* 189:550-553. (German)
- *Bogen KT, Colston BW Jr., Machicao LK. 1992. Dermal absorption of diluted aqueous chloroform, trichloroethylene, and tetrachloroethylene in hairless guinea pigs. *Fund Appl Toxicol* 18:30-39.
- *Bois FY, Zeise L, Tozer TN. 1990. Precision and sensitivity of pharmacokinetic models for cancer risk assessments: Tetrachloroethylene in mice, rats, and humans. *Toxicol Appl Pharmacol* 102:300-315.
- Borm PJA, Barbanson B. 1988. Bias in biologic monitoring caused by concomitant medication. *J Occup Med* 30:214-223.
- *Bosco MG, Figa-Talamanca I, Salerno S. 1986. Health and reproductive status of female workers in dry cleaning shops. *Int Arch Occup Environ Health* 59:295-301.
- *Bouwer EJ, McCarty PL. 1982. Removal of trace chlorinated organic compounds by activated carbon and fixed film bacteria. *Environ Sci Technol* 16:836-843.
- *Bouwer EJ, McCarty PL. 1984. Modeling of trace organics biotransformation in the subsurface. *Ground Water* 22:433-440.
- *Bouwer EJ, Rittmann BE, McCarty PL. 1981. Anaerobic degradation of halogenated 1- and 2-carbon organic compounds. *Environ Sci Technol* 15:596-599.

8. REFERENCES

- *Bove FJ, Fulcomer MC, Klotz JB. 1995. Public drinking water contamination and birth outcomes. *Am J Epidemiol* 141:850-862.
- *Briving C, Jacobson I, Hamberger A, et al. 1986. Chronic effects of perchloroethylene and trichloroethylene on the gerbil brain amino acids and glutathione. *Neurotoxicology* 7:101-108.
- *Brodkin CA, Daniell W, Checkoway H, et al. 1995. Hepatic ultrasonic changes in workers exposed to perchloroethylene. *Occup Environ Med* 52:679-685.
- *Brodzinsky R, Singh HB. 1982. Volatile organic chemicals in the atmosphere: An assessment of available data. Menlo Park, CA: Atmospheric Science Center, SRI International. Contract 68-02-3452, 109-110.
- *Bronstein AC, Curran PL. 1988. Emergency care for hazardous materials exposure. Washington, DC: The C.V. Mosby Company, 93-94.
- *Bronzetti G, Bauer C, Corsi R, et al. 1983. Genetic and biochemical studies on perchloroethylene *in vitro* and *in vivo*. *Mutat Res* 116:323-331.
- Brown DP, Kaplan SD. 1985. Retrospective cohort mortality study of dry cleaner workers using perchloroethylene. National Institute for Occupational Safety and Health, Department of Health and Human Services, Cincinnati, OH.
- *Brown DP, Kaplan SD. 1987. Retrospective cohort mortality study of dry cleaning workers using perchloroethylene. *J Occup Med* 29:535-541.
- *Bruckmann P, Kersten W, Funcke W, et al. 1988. The occurrence of chlorinated and other organic trace compounds in urban air. *Chemosphere* 17:2363-2380.
- *Brugnone F, Perbellini L, Giuliani C, et al. 1994. Blood and urine concentrations of chemical pollutants in the general population. *Med Lav* 85(5):370-389.
- *Buben JA, O'Flaherty EJ. 1985. Delineation of the role of metabolism in the hepatotoxicity of trichloroethylene and perchloroethylene: A dose-effect study. *Toxicol Appl Pharmacol* 78:105-122.
- *Buckley JD, Robison LL, Swotinsky R, et al. 1989. Occupational exposures of parents of children with acute nonlymphocytic leukemia: A report from the children's cancer study group. *Cancer Res* 49:4030-4037.
- *Budavari S, ed. 1989. The Merck index. 11th ed. Rahway, NJ: Merck and Co., Inc., 1449.
- *Bundschuh I, Herbort C, Fels LM, et al. 1993. Renal fibronectin excretion as a marker for renal environmental toxins. *Contrib Nephrol* 101: 177- 184.
- *Burmester DE. 1982. The new pollution-groundwater contamination. *Environment* 24:6-13, 33-36.
- *Byczkowski JZ, Fisher JW. 1994. Lactational transfer of tetrachloroethylene in rats. *Risk Analysis* 14(3): 339-349.

8. REFERENCES

- *Byczkowski JZ, Fisher JW. 1995. A computer program linking physiologically based pharmacokinetic model with cancer risk assessment for breast-fed infants. *Comput Methods Programs Biomed* 46:155-163.
- *Byczkowski JZ, Kinkead ER, Leahy HF, et al. 1994. Computer simulation of the lactational transfer of tetrachloroethylene in rats using a physiologically based model. *Toxicol Appl Pharmacol* 125:228-236.
- *Byers VS, Levin AS, Ozonoff DM, et al. 1988. Association between clinical symptoms and lymphocyte abnormalities in a population with chronic domestic exposure to industrial solvent-contaminated domestic water supply and a high incidence of leukemia. *Cancer Immunol Immunother* 27:77-81.
- *Cai S-X, Huang M-Y, Chen Z, et al. 1991. Subjective symptom increase among dry cleaning workers exposed to tetrachloroethylene vapor. *Industrial Health* 29: 111-121.
- *Callen DF, Wolf CR, Philpot RM. 1980. Cytochrome P-450 mediated genetic activity and cytotoxicity of seven halogenated aliphatic hydrocarbons in *Saccharomyces cerevisiae*. *Mutat Res* 77:55-63.
- Cantelli-Forti G, Bronzetti G. 1988. Mutagenesis and carcinogenesis of halogenated ethylenes. *Ann NY Acad Sci* 534:679-693.
- *Carpenter CP. 1937. The chronic toxicity of tetrachloroethylene. *J Ind Hyg Toxicol* 19:323-336.
- Cattley RC, Popp JA. 1989. Differences between the promoting activities of the peroxisome proliferator WY-14,643 and phenobarbital in rat liver. *Cancer Res* 49:3246-3251.
- *Cavalleri A, Gobba F, Paltrinieri M, et al. 1994. Perchloroethylene exposure can induce colour vision loss. *Neurosci Lett* 179:162-166.
- *CELDS. 1994. Computer-environmental legislative data system database. University of Illinois, Urbana, IL.
- *CEPA. 1993. Canadian Environmental Protection Act. Priority substances list assessment report: Tetrachloroethylene. Government of Canada, Environment Canada, Health Canada, 10-14, 29-30.
- *Chapman JAW, Connolly JG, Rosenbaum L. 1981. Occupational bladder cancer: A case-control study. In: Connolly JE, ed. *Carcinoma of the bladder*. New York, NY: Raven Press, 45-54.
- *Chaudhuri RN, Mukerji AK. 1947. Death following administration of tetrachloroethylene. *Indian Medical Gazette* 82:115-116.
- *C&EN. 1994. Facts and Figures for the Chemical Industry. *Chemical and Engineering News*, July 4, 1994.

8. REFERENCES

- *Chemical Products Synopsis. 1985. Perchloroethylene. Mannsville Chemical Products Corp., Cortland, N.Y.
- Chemical Profile. 1995. Perchloroethylene. Chemical Marketing Reporter, March 13, 1995.
- *Chen CW, Blancato JN. 1987. Role of pharmacokinetic modeling in risk assessment: Perchloroethylene as an example. In: Pharmacokinetics in risk assessment, drinking water and health. Vol. 8. Washington, DC: National Academy of Sciences, National Research Council, 369-390.
- Chen CW, Hiremath CB, Parker JC. 1986. Updated carcinogenicity assessment for tetrachloroethylene (perchloroethylene, pert, PCE): Addendum to the health assessment document for tetrachloroethylene (perchloroethylene). External review draft. Issue 12.
- *Chen KM, Dallas CE, Muralidhara S, et al. 1993. Analyses of volatile C2 haloethanes and haloethenes in tissues: Sample preparation and extraction. *J Chromatogr* 612:199-208.
- *Chilton PJ, Lawrence AR, Barker A. 1990. Chlorinated solvents in chalk aquifers: some preliminary observations on behavior and transport. In: Telford T, International Chalk Symposium. London, England, 605-610.
- Chiou CT, Peters LJ, Freed VH. 1979. A physical concept of soil-water equilibria for nonionic organic compounds. *Science* 206:831-832.
- Chiou CT, Freed VH, Peters LJ, et al. 1980. Evaporation of solutes from water. *Environment International* 3:231-236.
- *Chodola GR, Biswas N, Bewtra JK, et al. 1989. Fate of selected volatile organic substances in aqueous environment. *Water Pollution Research Journal of Canada* 24:119-142.
- *Christensen JM, Rasmussen K, Koppen B. 1988. Automatic headspace gas chromatographic method for the simultaneous determination of trichloroethylene and metabolites in blood and urine. *J Chromatogr* 442:317-323.
- *Class TH, Ballschmitter K. 1986. Chemistry of organic traces in air. VI: Distribution of chlorinated Cl-C4 hydrocarbons in air over the northern and southern Atlantic Ocean. *Chemosphere* 15:413-427.
- Clewell HJ III, Andersen ME. 1985. Risk assessment extrapolations and physiological modeling. *Toxicol Ind Health* 1(4):111-131.
- Cochran JW, Yates MV, Henson JM. 1988. A modified purge-and-trap-gas chromatography method for analysis of volatile halocarbons in microbiological degradation studies. *J Microbial Methods* 8:347-354.
- Coffin RR, Witherell LE, Novick LF, et al. 1987. Establishment of an exposure level to tetrachloroethylene in ambient air in Vermont. *Public Health Rep* 102:681-686.
- Cohen AJ, Grasso P. 1981. Review of the hepatic response to hypolipidaemic drugs in rodents and assessment of its toxicological significance to man. *J Cosmet Toxicol* 19:585-605.

8. REFERENCES

- Cohen MA, Ryan PB, Yanagisawa Y, et al. 1989. Indoor/outdoor measurements of volatile organic compounds in Kanawha Valley of West Virginia. *J Air Pollut Control Assoc* 39:1086-1093.
- *Cohn P, J Klotz, F Bove, et al. 1994. Drinking water contamination and the incidence of leukemia and non-Hodgkin's lymphoma. *Environmental Health Perspectives* 102(6-7): 556-561.
- *Coler HR, Rossmiller HR. 1953. Tetrachloroethylene exposure in a small industry. *AMA Arch Ind Hyg Occup Med* 8:227-233.
- Conway JG, Tomaszewski KE, Olson MJ, et al. 1989. Relationship of oxidative damage to the hepatocarcinogenicity of the peroxisome proliferators di(2-ethylhexyl)phthalate and WY-14,643. *Carcinogenesis* 10:513-519.
- Corapcioglu MY, Hossain MA. 1990. Theoretical modeling of biodegradation and biotransformation of hydrocarbons in subsurface environments. *J Theor Biol* 142:503-516.
- *Cornish HH, Adefuin J. 1966. Ethanol potentiation of halogenated aliphatic solvent toxicity. *Am Ind Hyg Assoc J* 27:57-61.
- *Cornish HH, Ling BP, Barth ML. 1973. Phenobarbital and organic solvent toxicity. *Am Ind Hyg Assoc J* 34:487-492.
- *Costa AK, Ivanetich KM. 1980. Tetrachloroethylene metabolism by the hepatic microsomal cytochrome P-450 system. *Biochem Pharmacol* 29:2863-2869.
- *Cramer PH, Boggess KE, Hosenfeld JM, et al. 1988. Determination of organic chemicals in human whole blood: Preliminary method development for volatile organics. *Bull Environ Contam Toxicol* 40(4):612-618.
- *CRIS. 1994. Current Research Information System. U.S. Department of Agriculture, Beltsville, MD.
- Crist HL, Mitchell WJ. 1986. Field adult results with organic gas standards on volatile organic ambient air samplers equipped with Tenax GC. *Environ Sci Technol* 20:1260-1262.
- *Cupitt LT. 1987. Atmospheric persistence of eight air toxics. Environmental Protection Agency, Research Triangle Park, NC, Atmospheric Sciences Research Lab. EPA/600/3-87/004.
- *Daft JL. 1988. Rapid determination of fumigant and industrial chemical residues in food. *J Assoc Off Anal Chem* 71:748-760.
- *Daft JL. 1989. Determination of fumigants and related chemicals in fatty and nonfatty foods. *J Agric Food Chem* 37:560-564.
- *Dallas CE, Chen XM, Muralidhara S, et al. 1994a. Use of tissue disposition data from rats and dogs to determine species differences in input parameters for physiological model for perchloroethylene. *Environ Res* 67:54-67.

8. REFERENCES

- *Dallas CE, Chen XM, O'Barr K, et al. 1994b. Development of a physiologically based pharmacokinetic model for perchloroethylene using tissue concentration - time data. *Tox Appl Pharm* 128:50-59.
- *Dallas CE, Muralidhara S, Chen XM, et al. 1994c. Use of a physiologically based model to predict systemic uptake and respiratory elimination of perchloroethylene. *Tox Appl Pharm* 128:60-68.
- *Dallas CE, Chen XM, Muralidhara S, et al. 1995. Physiologically based pharmacokinetic model useful in prediction of the influence of species, dose, and exposure route on perchloroethylene pharmacokinetics. *J Toxicol Environ Health* 44:301-317.
- *Daniel JW. 1963. The metabolism of ³⁶Cl-labeled trichloroethylene and tetrachloroethylene in the rat. *Biochem Pharmacol* 12:795-802.
- *Deipser A, Stegmann, R. 1994. The origin and fate of volatile trace components in municipal solid waste landfills. *Waste Management and Research* 12:129-1393.
- *Dekant W. 1986. Metabolic conversion of tri- and tetrachloroethylene: Formation and deactivation of genotoxic intermediates. In: Chambers PL, Gehring P, Sakai F, eds. *New Concepts and Developments in Toxicology*. 211-221.
- *Dekant W, Martens G, Vamavakas S. et al. 1987. Bioactivation of tetrachloroethylene. Role of glutathione s-transferase-catalyzed conjugation versus cytochrome P-450 dependent phospholipid alkylation. *Drug Metab Dispos* 15:702-709.
- *Dekant W, Metzler M, Henschler D. 1986. Identification of S-1,2,2-trichlorovinyl-n-acetylcysteine as a urinary metabolite of tetrachloroethylene: Bioactivation through glutathione conjugation as a possible explanation of its nephrocarcinogenicity. *J Biochem Toxicol* 1:57-72.
- De La Iglesia FA, Farber E. 1982. Hypolipidemics carcinogenicity and extrapolation of experimental results for human safety assessments. *Toxicol Pathol* 10:152-170.
- *Dewalle FB, Chian ESK. 1981. Detection of trace organics in well water near a solid waste landfill. *J Am Water Works Assoc* 73:206-211.
- *DHHS. 1994. Seventh annual report on carcinogens. Summary 1994. U.S. Department of Health and Human Services, Public Health Service. Report to the National Institute of Environmental Health Sciences, Research Triangle Park, NC, by Technical Resources, Inc., Rockville, MD, 372-376.
- *Dickson AG, Riley JP. 1976. The distribution of short-chain halogenated aliphatic hydrocarbons in some marine organisms. *Mar Pollut Bull* 7:167-169.
- *Dietz EA, Singley KF. 1979. Determination of chlorinated hydrocarbons in water by headspace gas chromatography. *Anal Chem* 51(11):1809-1814.
- *Dilling WL. 1977. Interphase transfer processes. II. Evaporation rates of chloromethanes, ethanes, ethylenes, propanes, and propylenes from dilute aqueous solutions. Comparisons with theoretical predictions. *Environ Sci Technol* 11:405-409.

8. REFERENCES

- *Dilling WL, Tefertiller NB, Kallos GJ. 1975. Evaporation rates and reactivities of methylene chloride, chloroform, 1,1, 1-trichloroethane, trichloroethylene, tetrachloroethylene, and other chlorinated compounds in dilute aqueous solutions. *Environ Sci Technol* 9:833-838.
- *Doust HG, Huang J. 1992. The fate and transport of hazardous chemicals in the subsurface environment. *Water Science and Technology* 25:169-176.
- Dresen MD, Nichols EM. 1987. Environmental protection investigations and corrections series: Distribution of VOCs (Volatile organic compounds) in ground water in the southeast area of LLNL and vicinity. Washington, DC: U.S. Department of Energy, Issue 13.
- Droz PO, Guillemin MP. 1986. Occupational exposure monitoring using breath analysis. *J Occup Med* 28:593-603.
- *Dub RW, Asal NR. 1984. Mortality among laundry and dry cleaning workers in Oklahoma. *Am J Public Health* 74:1278-1280.
- *Dyksen JE, Hess AF III. 1982. Alternatives for controlling organics in groundwater supplies. *J Am Water Works Assoc* 74:394-402.
- *Earnest GS. 1996. Evaluation and control of perchloroethylene exposure during dry cleaning. *Applied Occupational and Environmental Hygiene* 11 (2) 125- 132.
- *Earnest GS. 1995. In-depth survey report: Chemical exposures and fire safety in commercial dry cleaners at A-One Cleaners, Cincinnati, Ohio. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Report No. ECTB 201-18a. 66 pages.
- *Earnest GS, Spencer AB. 1995. In-depth survey report: Perchloroethylene exposures in commercial dry cleaners at Brown's Cleaners, Santa Monica, California. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Report No. ECTB 201-16a. 47 pages.
- *Earnest GS, Moran AJ, Spencer AB et al. 1995a. In-depth survey report: Control of perchloroethylene exposures in commercial dry cleaners at Appearance Plus Cleaners, Cincinnati, Ohio. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Report No. ECTB: 201-15a. 45 pages.
- *Earnest GS, Moran AJ, Spencer AB. 1995b. In-depth survey report: Perchloroethylene exposures in commercial dry cleaners at Tuchman Cleaners (shop #27), Fishers, Indiana. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Report No. ECTB: 201-17a. 45 pages.
- *Earnest GS, Spencer AB, Smith SS, et al. 1995c. In-depth survey report: Perchloroethylene exposures and ergonomic risk factors in commercial dry cleaners at Tuchman Cleaners (shop #24), Carmel, Indiana. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Report No. ECTB: 201-19a. 61 pages.

8. REFERENCES

- *Ebrahim AS, Gopalakrishnan R, Murugesan A, et al. 1995. *In vivo* effect of vitamin E on serum and tissue glycoprotein levels in perchloroethylene induced cytotoxicity. *Mol Cell Biol* 144: 13-18.
- *ECETOC. 1990. Technical report No. 37. Tetrachloroethylene: Assessment of human carcinogenic hazard. Brussels, Belgium: European Chemical Industry Ecology and Toxicology Centre.
- *Echeverria D, White RF, Sampaio C. 1995. A behavioral evaluation of PCE exposure in patients and dry cleaners: a possible relationship between clinical and preclinical effects. *J Occup Environ Med* 37(6):667-680.
- Eichler DL, Mackey JH. 1986. The levels of certain volatile organic compounds in the ambient air of the United States. *Proc APCA Annu Meet* 79th. 6:3-17.
- *Ellenhorn MJ, Barceloux DG. 1988. *Medical toxicology: Diagnosis and treatment of human poisoning*. Elsevier, New York 986-988.
- *Entz RC, Diachenko GW. 1988. Residues of volatile halocarbons in margarines. *Food Addit Contam* 5:267-276.
- *Entz RC, Hollifield HC. 1982. Headspace gas chromatographic analysis of foods for volatile halocarbons. *J Agric Food Chem* 30:84-88.
- Entz RC, Thomas KW, Diachenko GW. 1982. Residues of volatile halocarbons in foods using headspace gas chromatography. *J Agric Food Chem* 30:846-849.
- *EPA. 1978. Pesticide chemicals. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 455.
- EPA. 1980. Hazardous waste management system: Identification and listing of hazardous waste: Interim rule. U.S. Environmental Protection Agency. Federal Register 45:78530-78550.
- *EPA. 1982a. Exposure and risk assessment for tetrachloroethylene. Office of water. U.S. Environmental Protection Agency, available through the National Technical Information Service (NTIS), Springfield, VA, PB85-221497.
- *EPA. 1982b. Methods for organic chemical analysis of municipal and industrial wastewater. Test method: Purgeable halocarbons - Method 601 and Purgeables - Method 624. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Monitoring and Support Laboratory, EPA 600/4-82-057. 601-1 to 601-10 and 624-1 to 624-12.
- *EPA. 1982c. Test methods for evaluating solid waste. Physical/chemical methods. Method 8010 - Halogenated volatile organics. Washington, DC: U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response, 8010-1 to 8010-12.
- *EPA. 1985a. Assessment of perchloroethylene as a potentially toxic air pollutant. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 61.

8. REFERENCES

- *EPA. 1985b. Designation, reportable quantities, and notification. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 302.
- *EPA. 1985c. Drinking water criteria document for tetrachloroethylene. (Final). Washington, DC: U.S. Environmental Protection Agency, Office of Drinking Water. NTIS PB86-118114.
- *EPA. 1985d. Health assessment document for tetrachloroethylene (perchloroethylene) - Final report. Washington, DC: U.S. Environmental Protection Agency, EPA/600/8-82/005F.
- EPA. 1986a. Addendum to the health assessment document for tetrachloroethylene (perchloroethylene): Updated carcinogenicity assessment. EPA/600/8-82/005FA. External review draft.
- *EPA. 1986b. Environmental Protection Agency: Guidelines for carcinogen risk assessment. Federal Register, 51(185):33992-34003.
- EPA. 1987a. Ground-water monitoring list. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 264, Appendix IX.
- *EPA. 1987b. Health advisories for 25 organics. Office of drinking water. U.S. Environmental Protection Agency, available through the National Technical Information Service (NTIS), Springfield, VA, PB87-235578IAS.
- EPA. 1987c. Reportable quantity adjustment. U.S. Environmental Protection Agency. Federal Register 52:8140-8186.
- *EPA. 1987d. Toxic chemical release reporting: Community right-to-know. U.S. Environmental Protection Agency. Federal Register 52:21152-21208.
- *EPA. 1988. Toxic chemical release reporting: Community right-to-know. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 372.
- EPA. 1989a. Interim methods for development of inhalation reference concentrations. U.S. Environmental Protection Agency. EPA/600/8-88/066F.
- EPA. 1989b. National primary and secondary drinking water regulations; Proposed rule. U.S. Environmental Protection Agency. Federal Register 54:22062-22160.
- EPA. 1989c. Reportable Quantity Adjustments. U.S. Environmental Protection Agency. Federal Register 54:33418-33426.
- *EPA. 1991a. Hazardous waste and superfund programs; amendment to preambles published at 54 FR 33418 (August 14, 1989) and 54 FR 50968 (December 11, 1989). U.S. Environmental Protection Agency. Federal Register 56:643-645.
- *EPA. 1991b. Science Advisory Board's review of the Office of Research and Development draft document: Response to issues and data submissions on the carcinogenicity of tetrachloroethylene (perchloroethylene) (February 1991). U.S. Environmental Protection Agency. EPA-SAB-EHC-91-013.

8. REFERENCES

- *EPA. 1995. EPA Office of Compliance sector notebook project. Profile of the dry cleaning industry. Office of Compliance, Office of Enforcement and Compliance Assurance, U.S. EPA, Washington, DC. EPA/310-R-95-001.
- *EPA. 1996a. Air Quality; Revision to definition of volatile organic compounds - exclusion of perchloroethylene. Federal Register 61(26):4588-4591.
- *EPA. 1996b. Discarded commercial chemical products, off-specification species, container residues and spill residues thereof. U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 261.33. FR 60:19165.
- *EPA. 1996c. Drinking water regulations and health advisories. Washington, D.C. U.S. Environmental Protection Agency. Office of Water. EPA 822-R-96-001.
- *EPA. 1996d. Ground-water monitoring list. U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 264, Appendix IX. FR 60:56952.
- *EPA. 1996e Hazardous wastes from specific sources. U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 261.32. FR 60:7848.
- *EPA. 1996f. List of hazardous substances and reportable quantities. U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 302. FR 60:35991.
- *EPA. 1996g. Lists of pollutants and applicability of part 61. Code of Federal Regulations 40 CFR 61.01. FR 60:12427.
- *EPA. 1996h. Specific toxic chemical listings. U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 372.65. FR 60:49803.
- *EPA. 1996i. Toxics criteria for those states not complying with the Clean Water Act Section 303(c)(2)(B). U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 131.36. FR 60:22235.
- *EPA. 1996j. US Environmental Protection Agency. Code of Federal Regulations. 40 CFR 455. FR 58: 50689.
- *EPA. 1996k. US Environmental Protection Agency. Code of Federal Regulations. 40 CFR 401. FR 58: 45035.
- *EPA. 1996l. US Environmental Protection Agency. Code of Federal Regulations. 40 CFR 261.31. FR 60: 33913.
- *Eskenazi B, Fenster L, Hudes M, et al. 1991a. A study of the effect of perchloroethylene exposure on the reproductive outcomes of wives of dry cleaning workers. American Journal of Industrial Medicine 20:593-600.
- *Eskenazi B, Wyrobek AJ, Fenster L, et al. 1991b. A study of the effect of perchloroethylene exposure on semen quality in dry cleaning workers. American Journal of Industrial Medicine 20:575-591.

8. REFERENCES

- Fagliano J, Berry M, Bove F, et al. 1990. Drinking water contamination and the incidence of leukemia: An ecologic study. *Am J Public Health* 80:1209-1212.
- *FDA. 1977. Indirect food additives: Adhesive coatings and components. *Federal Register* 42(50):14534-14555.
- FDA. 1984. Indirect food additives: Adjuvants production aids and sanitizers. 21 CFR Part 178:308.
- Feldman RG, Chirico-Post J, Proctor SP. 1988. Blink reflex latency after exposure to tetrachloroethylene in well water. Pittsburgh, PA: Meeting on Epidemiology in Environmental Health held at the First International Symposium on Environmental Epidemiology, June 3-5, 1987. *Arch Environ Health* 42:143-148.
- *Ferrario JB, Lawler GC, Deleon IR, et al. 1985. Volatile organic pollutants in biota and sediments of Lake Pontchartrain. *Bull Environ Contam Toxicol* 34:246-255.
- *Ferroni C, Selis L, Mutti A, et al. 1992. Neurobehavioral and neuroendocrine effects of occupational exposure to perchloroethylene. *NeuroToxicology* 13:243-247.
- Foresman TW. 1986. Mapping monitoring and modelling of hazardous waste sites. Venice, Italy: International Conference on Man's Role in Changing the Global Environment, Part B, October 21-26, 1985. *Sci Total Environ* 56:255-263.
- *Franchini I, Cavotorta A, Falzoi M, et al. 1983. Early indicators of renal damage in workers exposed to organic solvents. *Int Arch Occup Environ Health* 52:1-9.
- *Frantz SW, Watanabe PG. 1983. Tetrachloroethylene: Balance and tissue distribution in male Sprague-Dawley rats by drinking water administration. *Toxicol Appl Pharmacol* 69:66-72.
- *Fredriksson A, Danielsson BRG, Eriksson P. 1993. Altered behavior in adult mice orally exposed to tri- and tetrachloroethylene as neonates. *Toxicology Letters* 66: 13- 19.
- *Freedman DL, Gossett JM. 1989. Biological reductive dechlorination of tetrachloroethylene and trichloroethylene to ethylene under methanogenic conditions. *Appl Environ Microbiol* 55:2144-2151.
- *Friberg L, Kylin B, Nystrom A. 1953. Toxicities of trichloroethylene and tetrachloroethylene and Fujiwara's pyridine-alkali reaction. *Acta Pharmacol Toxicol* 9:303-312.
- *Friesel P, Milde G, Steiner B. 1984. Interactions of halogenated hydrocarbons with soils. *Fresenius Z Anal Chem* 319:160-164.
- *FSTRAC. 1990. Summary of state and federal drinking water standards and guidelines. Federal-State Toxicology and Regulatory Alliance Committee, Chemical Communication Subcommittee.
- *Gamier R, Bedouin J, Pepin G, et al. 1996. Coin-operated dry cleaning machines may be responsible for acute tetrachloroethylene poisoning: Report of 26 cases including one death. *Clinical Toxicol* 34(2):191-197.

8. REFERENCES

- *Gay BW Jr, Hanst PL, Bulfalini JJ, et al. 1976. Atmospheric oxidation of chlorinated ethylenes. *Environmental Science and Technology* 10:58-67.
- *Gearhart JM, Mahle DA, Greene RJ. 1993. Variability of physiologically based pharmacokinetic (PBPK) model parameters and their effects on PBPK model predictions in a risk assessment for perchloroethylene. 68131-144.
- *Gennari P, Naldi M, Motta R, et al. 1992. Gamma-glutamyltransferase isoenzyme pattern in workers exposed to tetrachloroethylene. *Am J Ind Med* 21:661-671.
- *Germolec DR, Yang RSH, Ackermann MF, et al. 1989. Toxicology studies of a chemical mixture of 25 groundwater contaminants: II. Immunosuppression in B6C3F₁ mice. *Fundam Appl Toxicol* 13:377-387.
- Geyer H, Scheunert I, Korte F. 1986. Bioconcentration potential of organic environmental chemicals in humans. *Regul Toxicol Pharmacol* 6:313-347.
- *Ghantous H, Danielsson BRG, Dencker L, et al. 1986. Trichloroacetic acid accumulates in murine amniotic fluid after tri- and tetrachloroethylene inhalation. *Acta Pharmacol Toxicol* 58: 105-114.
- *Ghittori S, Imbraini M, Pezzagno G. 1987. The urinary concentration of solvents as a biological indicator of exposure: Proposal for the biological equivalent exposure limit for nine solvents. *Am Ind Hyg Assoc J* 48:786-790.
- *Gimmi T, Fluehler H, Studer B, et al. 1993. Transport of volatile chlorinated hydrocarbons in unsaturated aggregated media. *Water Air Soil Pollut* 68:291-305.
- *Giovannini L, Guglielmi G, Casini T, et al. 1992. Effect of ethanol chronic use on hepatotoxicity in rats exposed to tetrachloroethylene. *Int J Tiss React* XIV(6):281-285.
- *Goldberg ME, Johnson HE, Pozzani UC, et al. 1964. Effect of repeated inhalation of vapors of industrial solvents on animal behavior. I. Evaluation of nine solvent vapors on pole-climb performance in rats. *Am Ind Hyg Assoc J* 25:369-375.
- *Goldsworthy TL, Popp JA. 1987. Chlorinated hydrocarbon-induced peroxisomal enzyme activity in relation to species and organ carcinogenicity. *Toxicol Appl Pharmacol* 88:225-233.
- *Goldsworthy TL, Lyght O, Burnett VL, et al. 1988. Potential rate of α -2 μ -globulin, protein droplet accumulation, and cell replication in the renal carcinogenicity of rats exposed to trichloroethylene, perchloroethylene, and pentachloroethane. *Toxicol Appl Pharmacol* 96:367-379.
- *Gossett JM. 1987. Measurement of Henry's law constants for Cl and C2 chlorinated hydrocarbons. *Environmental Science and Technology* 21:202-208.
- *Green T. 1990. Species differences in carcinogenicity: The role of metabolism in human risk evaluation. *Teratogenesis, Carcinog, and Mutagen* 10: 103-113.

8. REFERENCES

- *Green T, Odum J, Nash JA, et al. 1990. Perchloroethylene-induced rat kidney tumors: An investigation of the mechanisms involved and their relevance to humans. *Toxicol Appl Pharmacol* 103:77-89.
- *Gregersen P. 1988. Neurotoxic effects of organic solvents in exposed workers: Two controlled follow-up studies after 5.5 and 10.6 years. *Am J Ind Med* 14:681-701.
- *Gregersen P, Angelso B, Nielsen TE, et al. 1984. Neurotoxic effects of organic solvents in exposed workers: An occupational neuropsychological and neurological investigation. *Am J Ind Med* 5:201-225.
- *Greim H, Bonse G, Radwan Z, et al. 1975. Mutagenicity *in vitro* and potential carcinogenicity of chlorinated ethylenes as a function of metabolic oxirane formation. *Biochem Pharmacol* 24:2013-2017.
- *Grob K, Frauenfelder C, Artho A. 1990. Uptake by foods of tetrachloroethylene, trichloroethylene, toluene, and benzene from air. *Z Lebensm Unters Forsch* 191:435-441.
- *Guberan E, Fernandez J. 1974. Control of industrial exposure to tetrachloroethylene by measuring alveolar concentrations: theoretical approach using a mathematical model. *Br J Ind Med* 31:159-167.
- *Gulyas H, Hemmerling L. 1990. Tetrachloroethene air pollution originating from coin-operated dry cleaning establishments. *Environ Res* 53:90-99.
- *Guo Z, Tichenor BA, Mason MA, et al. 1990. The temperature dependence of the emission of perchloroethylene from dry cleaned fabrics. *Environ Res* 52: 107- 115.
- *Haddad LM, Winchester JF. 1990. Clinical management of poisoning and drug overdose. Second edition. Philadelphia, PA: W.B. Saunders Company, Harcourt Brace Jovanovich, Inc., 13-20.
- *Haerer AF, Udelman HD. 1964. Acute brain syndrome secondary to tetrachloroethylene ingestion. *Am J Psychiatry* 12:78-79.
- Hajimiragha H, Ewers U, Jansen-Rosseck R. 1986. Human exposure to volatile halogenated hydrocarbons from the general environment. *Int Arch Occup Environ Health* 58:141-150.
- *Hake CL, Stewart RD. 1977. Human exposure to tetrachloroethylene: Inhalation and skin contact. *Environ Health Perspect* 21:231-238.
- *Hanioka N, Jinno H, Toyo'oka T, et al. 1995. Induction of rat liver drug-metabolizing enzymes by tetrachloroethylene. *Arch Environ Contam Toxicol* 28:273-280.
- Hara M, Kino M, Yamamoto et al. 1985. Cardiovascular effects of organic solvents (perchloroethylene) (PERC). *Jpn Circ J* 49:887-888.
- *Hardin BD, Bond GP, Sikov MR, et al. 1981. Testing of selected workplace chemicals for teratogenic potential. *Stand J Work Environ Health [Suppl 4]* 7:66-75.
- *Harkov R, Gianti SJ, Jr., Bozzelli JW, et al. 1985. Monitoring volatile organic compounds at hazardous and sanitary landfills in New Jersey. *J Environ Sci Health* 20:491-501.

8. REFERENCES

- *Harkov R, Kezbekus B, Bozzelli JW, et al. 1984. Comparison of selected volatile organic compounds during the summer and winter at urban sites in New Jersey. *Sci Total Environ* 38:259-274.
- Harkov R, Kezbekus B, Bozzelli JW. 1987. Volatile organic compounds at urban sites in New Jersey. In: Lioy & Daisey, eds. *Toxic Air Pollution*: Chelsea, MI: Lewis Pub. Inc, 69-90.
- *Hartwell TD, Crowder JH, Sheldon LS, et al. 1985. Levels of volatile organics in indoor air. In: *Proceedings of the Air Pollution Control Association 78th Annual Meeting, Volume 3* 85-30.B., 2-12.
- *Hartwell TD, Pellizzari ED, Perritt RL, et al. 1987. Comparison of volatile organic levels between sites and seasons for the total exposure assessment methodology (TEAM) study. *Atmos Environ* 21:2413-2424.
- *Hasanen E, Soinen V, Pyysalo M, et al. 1979. The occurrence of aliphatic chlorine and bromine compound in automobile exhaust. *Atmos Environ* 13:1217-1219.
- *Hattis D, White P, Koch P. 1993. Uncertainties in pharmacokinetic modeling for perchloroethylene: II. Comparison of model predictions with data for a variety of different parameters. *Risk Analysis* 13(6):599-610.
- *Hattis D, White P, Marmorstein L, et al. 1990. Uncertainties in pharmacokinetic modeling for perchloroethylene. I. Comparison of model structure, parameters, and predictions for low-dose metabolism rates for models derived by different authors. *Risk Analysis* 10(3):449-458.
- *Hauser TR, Bromberg SM. 1982. EPA's monitoring program at Love Canal 1980. *Environmental Monitoring and Assessment* 2:249-271.
- *Haworth S, Lawlor T, Mortelmans K, et al. 1983. *Salmonella* mutagenicity test results for 250 chemicals. *Environ Mutagen Supplement* 1:3-142.
- *Hayes JR, Condie LW Jr, Borzelleca JF. 1986. The subchronic toxicity of tetrachloroethylene (perchloroethylene) administered in the drinking water of rats. *Fundam Appl Toxicol* 7:119-125.
- *HAZDAT. 1996. Agency for Toxic Substances and Disease Registry (ATSDR), Atlanta, GA.
- *Heikes DL, Hopper ML. 1986. Purge and trap method for determination of fumigants in whole grains, milled grain products, and intermediate grain-based foods. *J Assoc Off Anal Chem* 69:990-998.
- *Heineman EF, Cocco P, Gomez MR, et al. 1994. Occupational exposure to chlorinated aliphatic hydrocarbons and risk of astrocytic brain cancer. *Am J Ind Med* 26: 155-169.
- Hemminiki K, Franssila E, Vainio H, et al. 1980. Spontaneous abortion among female chemical workers in Finland. *Int Arch Occup Environ Health* 45:123-126.
- Hemminiki K, Saloniemi I, Luoma K, et al. 1980. Transplacental carcinogens and mutagens: Childhood cancer, malformation and abortions as risk indicators. *J Toxicol Environ Health* 6:1115-1126.

8. REFERENCES

- *Henschler D. 1977. Metabolism and mutagenicity of halogenated olefins: A comparison of structure and activity. *Environ Health Perspect* 21:61-64.
- *Herren-Freund SL, Pereira MA, Khoury MD, et al. 1987. The carcinogenicity of trichloroethylene and its metabolites, trichloroacetic acid and dichloroacetic acid, in mouse liver. *Toxicol Appl Pharmacol* 90: 183- 189.
- Herzfeld D, Van Der Gun K-D, Louw R. 1989. Quantitative determination of volatile organochlorine compounds in water by GC-headspace analysis with dibromoethane as an internal standard. *Chemosphere* 18:1425-1430.
- *Hook EB. 1982. Perspectives in mutation epidemiology: 2. Epidemiologic and design aspects of studies of somatic chromosome breakage and sister-chromatic exchange. *Mutat Res* 99:373-382.
- *Hoover R, Mason TJ, McKay FW, et al. 1975. Geographic patterns of cancer mortality in the United States. In: Fraumeni JF, ed. *Persons at High Risk of Cancer: An Approach to Cancer Etiology and Control*. New York, NY: Academic Press. 343-360.
- *Hov O, Penkett SA, Isaksen ISA, et al. 1984. Organic gases in the Norwegian arctic. *Geophys Res Lett* 11:425-428.
- *Hoyt SD, Smith VL. 1991. Measurement of toxic organic compounds in ambient air using EPA method TO-14. In: *Capillary chromatography - The applications*. Jennings W, Nikelly JG eds. Huthig Buch Verlag Heidelberg, 83-94.
- *HSDB. 1996. Hazardous Substance Data Bank. National Library of Medicine, National Toxicology Information Program, Bethesda, MD.
- *IARC. 1979. IARC monographs on the evaluation of the carcinogenic risk of chemicals in humans. Tetrachloroethylene. Lyon, France: World Health Organization, 20:491-514.
- IARC. 1982. IARC monographs on the evaluation of the carcinogenic risk of chemicals to humans. Suppl. 4. Chemicals, industrial processes, and industries associated with cancer in humans. Lyon, France: World Health Organization. 243-245.
- *IARC. 1987. IARC monographs on the evaluation of the carcinogenic risks to humans. Suppl 7: Overall evaluations of carcinogenicity: An updating of IARC Monographs Volumes 1 to 42. Lyon, France: World Health Organization.
- *IARC. 1995. Tetrachloroethylene. IARC monographs of the evaluation of carcinogenic risks to humans, Vol 63, Drycleaning, some chlorinated solvents and other industrial chemicals. 159-221.
- *Ikeda M, Imamura T. 1973. Biological half-life of trichloroethylene and tetrachloroethylene in human subjects. *Int Arch Arbeitsmed* 31:209-224.

8. REFERENCES

Ikeda M, Koizumi A, Kashara M. 1987. Validation of passive dosimetry through biological monitoring and its application in solvent workplaces. In: Berlin A, Brown RH, Saunders KJ, eds. Diffusive Sampling, An Alternative Approach to Workplace Air Monitoring, Proceedings of an International Symposium, Luxembourg, 22-26 September 1986, London, Royal Society of Chemistry.

*Ikeda M, Koizuma A, Watanabe T, et al. 1980. Cytogenetic and cytokinetic investigations on lymphocytes from workers occupationally exposed to tetrachloroethylene. *Toxicol Lett* 5:251-256.

*Ikeda M, Ohtsuji H, Imamura T, et al. 1972. Urinary excretion of total trichloro-compounds, trichloroethanol, and trichloroacetic acid as a measure of exposure to trichloroethylene and tetrachloroethylene. *Br J Ind Med* 29:328-333.

*Imbriani M, Ghittori S, Pezzagno G, et al. 1988. Urinary excretion of tetrachloroethylene (perchloroethylene) in experimental and occupational exposure. *Arch Environ Health* 43:292-298.
Ince N, Inel Y. 1989. Volatilization of organic chemicals from water. *Water Air Soil Pollut* 47:71-79.

*IRIS. 1996. Integrated Risk Information System. U.S. Environmental Protection Agency. Office of Health and Environmental Assessment, Environmental Criteria and Assessment Office, Cincinnati, OH.

*Itoh N, Kutsuna S, Ibusuki T. 1994. A product study of the OH radical initiated oxidation of perchloroethylene and trichloroethylene. *Chemosphere* 28(11):2029-2040.

Jakobson I, Holmberg B, Ekner A. 1986. Venous blood levels of inhaled trichloroethylene in female rats and changes induced by interacting agents. *Acta Pharmacol Toxicol* 59:135-143.

*Jakobson I, Wahlberg JE, Holmberg B, et al. 1982. Uptake via the blood and elimination of 10 organic solvents following epicutaneous exposure of anesthetized guinea pigs. *Toxicol Appl Pharmacol* 63:181-187.

*Jang J-Y, Kang S-K, Chung HK. 1993. Biological exposure indices of organic solvents for Korean workers. *Int Arch Occup Environ Health* 65:S219-S222.

*Jeffers PM, Ward LM, Woytowitch LM, et al. 1989. Homogenous hydrolysis rate constants for selected chlorinated methanes, ethanes, ethenes and propanes. *Environ Sci Technol* 23:965-969.

*Jung WT, Fujita M, Sohn DH. 1992. Levels of volatile halogenated hydrocarbons in Tokyo rain and their seasonal, time-series changes. *Japanese Journal of Toxicology and Environmental Health* 38:490-497.

*Kaiser KLE, Comba ME. 1986. Tracking river plumes with volatile halocarbon contaminants: The St. Clair River-Lake St. Clair Example. *Environ Toxicol Chem* 5:965-976.

*Kanazawa S, Filip Z. 1987. [Effects of trichloroethylene, tetrachloroethylene, and dichloromethane on soil biomass and microbial counts]. *Zentralbl Bakteriell Hyg [B]* 184:24-33. (German)

Kaplan SD. 1980. Dry-cleaner workers exposed to perchloroethylene. A Retrospective cohort mortality study. Department of Health, Education, and Welfare Contract No. 210-77-0094. Cincinnati, OH: National Institute for Occupational Safety and Health.

8. REFERENCES

- *Karlsson J-E, Rosengren LE, Kjellstrand P, et al. 1987. Effects of low-dose inhalation of three chlorinated aliphatic organic solvents on deoxyribonucleic acid in gerbil brain. *Stand J Work Environ Health* 13:453-458.
- *Katz RM, Jowett D. 1981. Female laundry and dry cleaning workers in Wisconsin. A mortality analysis. *Am J Public Health* 71:305-307.
- *Kawakami T, Takano T, Araki R. 1988. Synergistic interaction of tri- and tetra-chloroethylene, hypoxia, and ethanol on the atrioventricular conduction of the perfused rat heart. *Ind Health* 26:25-33.
- *Kawamura K, Kaplan IR. 1983. Organic compounds in the rainwater of Los Angeles. *Environ Sci Technol* 17:497-501.
- *Kawasaki M. 1980. Experiences with the test scheme under the chemical control law of Japan: An approach to structure-activity correlations. *Ecotoxicol Environ Safety* 4:444-454.
- *Kawata K, Fujieda Y. 1993. Volatile chlorinated hydrocarbons in ambient air at Niigata area. *Japanese Journal of Toxicology and Environmental Health* 39:474-479.
- Kawauchi T, Nishiyama K. 1989. Residual tetrachloroethylene in dry-cleaned clothes. *Environ Res* 48:296-301.
- *Keil SL. 1985. Tetrachloroethylene. In: Grayson M, and Eckroth P, eds., *Kirk-Othmer Concise Encyclopedia of Chemical Technology*. New York, NY: John Wiley and Sons, 754-762.
- *Keith LH, Garrison AW, Allen FR, et al. 1976. Identification of organic compounds in drinking water from thirteen United States cities. In: Keith LH, ed. *Identification and Analysis of Organic Pollutants in Water*. Ann Arbor, MI: Ann Arbor Press, 329-373.
- *Kenaga EE. 1980. Predicted bioconcentration factors and soil sorption coefficients of pesticides and other chemicals. *Ecotoxicol Environ Safety* 4:26-38.
- *Kendrick JF. 1929. The treatment of hookworm disease with tetrachloroethylene. *Am J Trop Med* 483-488.
- *Khalil MAK, Rasmussen RA. 1983. Gaseous tracer of arctic haze. *Environ Sci Technol*. 17:157-164.
- *Kido K, Shiratori T, Watanabe T, et al. 1989. Correlation of tetrachloroethylene in blood and in drinking water: A case of well water pollution. *Bull Environ Contam Toxicol* 43:444-453.
- *Kinkead ER, Leahy HF. 1987. Evaluation of the acute toxicity of selected groundwater contaminants. *Govt Reports Announcements & Index (GRA&I), NTIS/AD-A180 198/4, Issue 17*.
- *Kirchner K, Helf D, Ott P, et al. 1990. The reaction of OH radicals with 1,1-di, tri- and tetrachloroethylene. *Ber Bunsenges Phys Chem* 94:77-83.
- Kitamori S, Koga K, Matuka S, et al. 1986. Groundwater pollution by tetrachloroethylene. *Zenkoku Kogaiken Kaishi* 11:93-96.

8. REFERENCES

- *Kjellstrand P, Bjerkemo M, Adler-Maihofer M, et al. 1985. Effects of solvent exposure on testosterone levels and butyrylcholinesterase activity in mice. *Acta Pharmacol Toxicol* 57:242-249.
- *Kjellstrand P, Holmquist B, Kanje M, et al. 1984. Perchloroethylene: Effects on body and organ weights and plasma butyrylcholinesterase activity in mice. *Acta Pharmacol Toxicol* 54:414-424.
- *Klaassen CD, Plaa GL. 1966. Relative effects of various chlorinated hydrocarbons on liver and kidney function in mice. *Toxicol Appl Pharmacol* 9:139-151.
- *Klaassen CD, Plaa GL. 1967. Relative effects of various chlorinated hydrocarbons on liver and kidney function in dogs. *Toxicol Appl Pharmacol* 10:119-131.
- *Kline SA, McCoy EC, Rosenkranz HS, et al. 1982. Mutagenicity of chloroalkene epoxides in bacterial systems. *Mutat Res* 101:115-125.
- *Kluwe WM, Abdo KM, Huff J. 1984. Chronic kidney disease and organic chemical exposure: Evaluations of causal relationships in humans and experimental animals. *Fundam Appl Toxicol* 4:899-901.
- Kobayashi H, Hobara T, Sakai T. 1989. Effects of inhalation of several organic solvents on left ventricular DP-DT. *Jpn J Ind Health* 31:136-141.
- *Kobayashi S, Hutcheon DE, Regan J. 1982. Cardiopulmonary toxicity of tetrachloroethylene. *J Toxicol Environ Health* 10:23-30.
- *Koch R, Schlegelmilch R, Wolf HU. 1988. Genetic effects of chlorinated ethylenes in the yeast *Saccharomyces cerevisiae*. *Mutat Res* 206:209-216.
- *Koenig HP, Lahl U, Kock H. 1987. Determination of organic volatiles in ambient air in the area of a landfill. *J Aerosol Sci* 18:837-840.
- Koizumi A, Kumai M, Ikeda M. 1983. Dose-dependent induction and suppression of liver mixedfunction oxidase system in chlorinated hydrocarbon metabolism. *J Appl Toxicol* 3:208-217.
- *Koppel C, Arndt I, Arendt U, et al. 1985. Acute tetrachloroethylene poisoning: Blood elimination kinetics during hyperventilation therapy. *Clin Toxicol* 23: 103-115.
- Korpela M. 1988. Organic solvents affect rat synaptosome membrane acetylcholinesterase and adenosine triphosphatase *in vitro*. *Arch Toxicol Supplement* 12:384-386.
- *Korpela M, Tahti H. 1986. The effect of selected organic solvents on intact human red cell membrane acetylcholinesterase *in vitro*. *Toxicol Appl Pharmacol* 85:257-262.
- *Kosson DS, Dienemann EA, Ahlert RC. 1985. Characterization and treatability studies of an industrial landfill leachate (Kin-But I). *Proceedings of the Industrial Waste Conference* 39:329-341.
- *Kringstad KP, Ljungquist PO, deSousa F, et al. 1981. Identification and mutagenic properties of some chlorinated aliphatic compounds in the spent liquor from Kraft pulp chlorination. *Environ Sci Technol* 15:562-566.

8. REFERENCES

- Krishnan K, Andersen ME. 1994. Physiologically based pharmacokinetic modeling in toxicology. In: Hayes W, ed. Principles and methods of toxicology. 3rd ed. New York, NY: Raven Press, Ltd., 149-188.
- Krishnan K, Andersen ME, Clewell HJ III, et al. 1994. Physiologically based pharmacokinetic modeling of chemical mixtures. In: Yang, RSA, ed. Toxicology of chemical mixtures. New York, NY: Academic Press, 399-437.
- *Kroneld R. 1989a. Volatile pollutants in suburban and industrial air. *Bull Environ Contam Toxicol* 42:868-872.
- Kroneld R. 1989b. Volatile pollutants in the environment and human tissues. *Bull Environ Contam Toxicol* 42:873-877.
- Kroneld R, Reuanen M. 1990. Determination of volatile pollutants in human and animal milk by GC-MS. *Bull Environ Contam Toxicol* 44:917-923.
- *Krost KJ, Pellizzari ED, Walburn SG, et al. 1982. Collection and analysis of hazardous organic emissions. *Anal Chem* 54:810-817.
- *Krotoszynski BK, Bruneau GM, O'Neill HJ. 1979. Measurement of chemical inhalation exposure in urban population in the presence of endogenous effluents. *J Anal Toxicol* 3:225-234.
- *Krska R, Taga K, Kellner R. 1993. New IR fiber-optic chemical sensor for in situ measurements of chlorinated hydrocarbons in water. *Applied Spectroscopy* 47(9):1484-1487.
- *Kukongviriyapan V, Kukongviriyapan U, Stacey NH. 1990. Interference with hepatocellular substrate uptake by 1,1,1-trichloroethane and tetrachloroethylene. *Toxicol Appl Pharmacol* 102(1):80-90.
- *Kylin B, Reichard H, Sumegi I, et al. 1963. Hepatotoxicity of inhaled trichloroethylene, tetrachloroethylene, and chloroform--single exposure. *Acta Pharmacol Toxicol* 20: 16-26.
- *Kylin B, Sumegi I, Yllner S. 1965. Hepatotoxicity of inhaled trichloroethylene and tetrachloroethylene--long-term exposure. *Acta Pharmacol Toxicol* 22:379-385.
- *Kyrklund T, Kjellstrand P, Haglid KG. 1990. Long-term exposure of rats to perchloroethylene, with and without a post-exposure solvent-free recovery period: Effects on brain lipids. *Toxicology Letters* 52:279-285.
- *Kyrklund T, Alling C, Kjellstrand P, et al. 1984. Chronic effects of perchloroethylene on the composition of lipid and acyl groups in cerebral cortex and hippocampus of the gerbil. *Toxicol Lett* 22:343-349.
- *Kyrklund T, Haglid K. 1991. Brain lipid composition in guinea pigs after intrauterine exposure to perchloroethylene. *Pharmacology & Toxicology* 68:146-148.

8. REFERENCES

- *Kyrklund T, Kjellstrand P, Haglid KG. 1988. Effects of exposure to freon 11, 1,1,1-trichloroethane or perchloroethylene on the lipid and fatty-acid composition of rat cerebral cortex. *Stand J Work Environ Health* 14:91-94.
- *Kyyronen P, Taskinen H, Lindbohm ML, et al. 1989. Spontaneous abortions and congenital malformations among women exposed to tetrachloroethylene in dry cleaning. *J Epidemiol Community Health* 43:346-351.
- *Lafuente A, Mall01 J. 1986. Thioethers in urine during occupational exposure to tetrachloroethylene. *Br J Ind Med* 43:68-69.
- *Lagakos SW, Wessen BJ, Zelen M, et al. 1986. An analysis of contaminated well water and health effects in Wobum, Massachusetts. *Journal of the American Statistical Association* 81:583-614.
- *Lash LH, Qian W, Xu Y, et al. 1996. Renal and hepatic metabolism and cytotoxicity of perchloroethylene (PER) by GSH conjugation. *The Toxicologist*, 30(1):A1613.
- *Lauwerys R, Herbrand J, Buchet JP, et al. 1983. Health surveillance of workers exposed to tetrachloroethylene in dry cleaning shops. *Int Arch Occup Environ Health* 52:69-77.
- *Lawrence AR, Chilton PJ, Barron RJ, et al. 1990. A method for determining volatile organic solvents in chalk pore waters (southern and eastern England) and its relevance to the evaluation of groundwater contamination. *Journal of Contaminant Hydrology* 6:377-386.
- *Leibman KC, Ortiz E. 1977. Metabolism of halogenated ethylenes. *Environ Health Perspect* 21:91-97.
- *Leung H-W. 1993. Physiologically-based pharmacokinetic modeling. In: Ballantyne B, Marrs T, Turner P, eds. *General and applied toxicology*. New York, NY: Stockton Press, I: 153-164.
- *Levine B, Fierro MF, Goza SW, et al. 1981. A tetrachloroethylene fatality. *J Forensic Sci* 26:206-209.
- *Lide DR, ed. 1990. *CRC Handbook of Chemistry and Physics*. 71st ed. Ann Arbor, MI: CRC Press, Inc., 3-242.
- *Ligoeki MP, Leuenberger C, Pankow JF. 1985. Trace organic compounds in rain-II. Gas scavenging of neutral organic compounds. *Atmos Environ* 19:1609-1617.
- Lindstrom K, et al. 1982. Psychological prognosis of diagnosed chronic organic solvent intoxication. *Neurobehav Toxicol Tetraol* 4:581-588.
- *Ling S, Lindsay WA. 1971. Perchloroethylene bums (Letter). *Br Med J* 3:115.
- Lock S, Kacew S, eds. 1988. *General principles in pediatric pharmacology and toxicology. Toxicologic and Pharmacologic Principles in Pediatrics*. New York, NY: Hemisphere Publishing Corporation, 1-16.

8. REFERENCES

- Lorenz H, Weber E, Omlor A, et al. 1990. Nachweis von Himschädigungen durch tetrachlorethen (Evidence of cerebral lesions caused by tetrachloroethylene). *Zbl Arbeitsmed* 40:355-364. [German]
- *Ludwig HR, Meister MV, Roberts DR, et al. 1983. Worker exposure to perchloroethylene in the commercial dry cleaning industry. *Am Ind Hyg Assoc J* 44:600-605.
- *Lukaszewski T. 1979. Acute tetrachloroethylene fatality. *Clin Toxicol* 15:411-415.
- *Lurker PA, Clark CS, Elia VJ. 1982. Atmospheric release of chlorinated organic compounds from the activated sludge process. *J Water Pollut Control Fed* 54:1566-1573.
- Lynge E. 1994. Danish cancer registry as a resource for occupational research. *J Occup Med* 36(11):1169-1173.
- *Lynge E, Carstensen B, Andersen O. 1995. Primary liver cancer and renal cell carcinoma in laundry and dry cleaning workers in Denmark. *Stand J Work Environ Health* 21:293-295.
- *Lynge E, Myg E, L. 1990. Primary liver cancer among women in laundry and dry cleaning work in Denmark. *Stand J Work Environ Health* 16:108-112.
- *Ma T-H, Sandhu SS, Peng Y, et al. 1992. Synergistic and antagonistic effects on genotoxicity of chemicals commonly found in hazardous waste sites *Mutation Research* 270:71-77.
- *MacMahon B. 1986. Comment. *Journal of the American Statistical Association* 81(395):597-599.
- MacPhail RC, Berman E, Elder JA. 1995. A multidisciplinary approach to toxicological screening: IV. Comparison of results. *J Tox Environ Health*. 45:211-220.
- *Magara Y, Furuichi T. 1986. Environmental pollution by trichloroethylene and tetrachloroethylene: A nationwide survey. *Dev Toxicol Environ Sci* 12:231-243.
- Mahieu P, Laurwerys R, Geubel A, et al. 1986. Acute intoxication by chlorinated solvents report of 3 observations. *Acta Clin Belg* 41:241-247.
- *Makide Y, Tominaga T, Rowland FS. 1979. Gas chromatographic analysis of halogenated hydrocarbons in air over Japan. *Chemistry Letters*, 355-358.
- *Marth E. 1987. Metabolic changes following oral exposure to tetrachloroethylene in subtoxic concentrations. *Arch Toxicol* 60:293-299.
- Materna BL. 1985. Occupational exposure to perchloroethylene in the dry cleaning industry. *Am Ind Hyg Assoc J* 46:268-273.
- Matter-Mueller C, Gujer W, Giger W. 1981. Transfer of volatile substances from water to the atmosphere. *Water Res* 15:1271-1279.
- *Mattsson JL, Albee RR, Yano BL, et al. 1992. Neurotoxicologic examination of rats exposed to 1,1,2,2-tetrachloroethylene vapor for 13 weeks. The Dow Chemical Company, sponsored by Halogenated Solvents Industry Alliance.

8. REFERENCES

- *Mazzullo M, Grilli S, Lattanzi G, et al. 1987. Evidence of DNA binding activity of perchloroethylene. *Res Commun Chem Pathol Pharmacol* 58:215-235.
- McCarty PL, Roberts PV. 1987. In-situ biotransformation of chlorinated solvents in groundwater. New Orleans, LA: American Chemical Society Division of Environmental Chemistry Preprint, 27:129-132.
- *McConnell G, Ferguson DM, Pearson CR. 1975. Chlorinated hydrocarbons and the environment. *Endeavour* 34:13-18.
- *McDonald AD, Armstrong B, Cherry NM, et al. 1986. Spontaneous abortion and occupation. *J Occup Med* 28:1232-1238.
- McDonald AD, McDonald JC, Armstrong B, et al. 1987. Occupation and pregnancy outcome. *Br J Ind Med* 44:521-526.
- *McDougal JN, Jepson GW, Clewell HJ III, et al. 1990. Dermal absorption of organic chemical vapors in rats and humans. *Fundam Appl Toxicol* 14:299-308.
- McKone TE. 1987. Human exposure to volatile organic compounds in household tap water: The indoor inhalation pathway. *Environ Sci Technol* 21:1194-1201.
- McLaughlin JK, et al. 1987. Occupational risks for renal cancer in Sweden. *Br J Ind Med* 44:119-123.
- *Meckler LC, Phelps DK. 1966. Liver disease secondary to tetrachloroethylene exposure: A case report. *J Am Med Assoc* 197:662-663.
- Mede A, Szakmary E, Ungvary G. 1989. Postnatal significance and reevaluation of minor and major anomalies found at day 21 of gestation. 17th Conference of the European Teratology Society, Budapest, Hungary, September 4-7, 1989. *Teratology* 40:290-291.
- *Mennear J, Maronpot R, Boorman G, et al. 1986. Toxicologic and carcinogenic effects of inhaled tetrachloroethylene in rats and mice. *Dev Toxicol Environ Sci* 12:201-210.
- *Meyer BR, Fischbein A, Rosenman K, et al. 1984. Increased urinary enzyme excretion in workers exposed to nephrotoxic chemicals. *Am J Med* 76:989-998.
- *Michael LC, Erickson MD, Parks SP, et al. 1980. Volatile environmental pollutants in biological matrices with a headspace purge technique. *Anal Chem* 52:1836-1841.
- Mieure JP. 1980. Determining volatile organics in water. *Environ Sci Technol* 14:930-935.
- *Milde G, Nerger M, Mergler R. 1988. Biological degradation of volatile chlorinated hydrocarbons in groundwater. Meeting on Groundwater Microbiology: Problems and Biological Treatment held at the International Association on Water Pollution Research and Control Symposium, Kuopio, Finland, August 4-6, 1987. *Water Sci Technol* 20:67-73.
- *Monster AC. 1986. Biological monitoring of chlorinated hydrocarbon solvents. *J Occup Med* 28:583-588.

8. REFERENCES

- *Monster AC. 1988. Biological markers of solvent exposure. *Arch Environ Health* 43:90-93.
- Monster AC, Boersma G. 1975. Simultaneous determination of trichloroethylene and metabolites in blood and exhaled air by gas chromatography. *Int Arch Occup Environ Health* 35:155-163.
- *Monster AC, Smolders JFJ. 1984. Tetrachloroethylene in exhaled air of persons living near pollution sources. *Int Arch Occup Environ Health* 53:331-336.
- *Monster AC, Boersma G, Steenweg H. 1979. Kinetics of tetrachloroethylene in volunteers: Influence of exposure concentration and work load. *Int Arch Occup Environ Health* 42:303-309.
- *Monster AC, Regouin-Peeters W, Van Schijndel A, et al. 1983. Biological monitoring of occupational exposure to tetrachloroethene. *Stand J Work Environ Health* 9:273-281.
- *Morgan B. 1969. Dangers of perchloroethylene (Letter). *Br Med J* 2:513.
- *Moser VC, Cheek BM, MacPhail RC. 1995. A multidisciplinary approach to toxicological screening: III. Neurobehavioral toxicity. *J Tox Environ Health* 45:173-210.
- *Moslen MT, Reynolds ES, Szabo S. 1977. Enhancement of the metabolism and hepatotoxicity of trichloroethylene and perchloroethylene. *Biochem Pharmacol* 26:369-375.
- Motohashi Y, Miyazaki Y, Takano T. 1993. Assessment of behavioral effects of tetrachloroethylene using a set of time-series analyses. *Neurotoxicology and Teratology* 15:3-10.
- *Motwani JN, Popp SA, Johnson GM, et al. 1986. Field screening techniques developed under the Superfund program. Management of Uncontrolled Hazardous Waste Sites, Hazardous Materials Control Research Institute, Washington, DC, December 1-3, 1986, Silver Spring, MD, 105-109.
- *Murakami K, Horikawa K. 1995. The induction of micronuclei in mice hepatocytes and reticulocytes by tetrachloroethylene. *Chemosphere* 31(7):3733-3739.
- *Murray AJ, Riley JP. 1973. Occurrence of some chlorinated aliphatic hydrocarbons in the environment. *Nature* 242:37-38.
- *Mutti A, Alinovi R, Bergamaschi E, et al. 1992. Nephropathies and exposure to perchloroethylene in dry-cleaners. *Lancet* 340(8813):189-193.
- Mutti A, et al. 1981. Organic-solvents and chronic glomerulonephritis: A cross-sectional study with negative findings for aliphatic and alicyclic C-C hydrocarbons. *J Appl Toxicol* 1:224-226.
- *Mutti A, Franchini I. 1987. Toxicity of metabolites to dopaminergic systems and the behavioral effects of organic solvents. *Br J Ind Med* 44:721-723.
- *Nakatsuka H, Watanabe T, Takeuchi Y, et al. 1992. Absence of blue-yellow color vision loss among workers exposed to toluene or tetrachloroethylene, mostly at levels below occupational exposure limits. *Int Arch Occup Environ Health* 64:113-117.

8. REFERENCES

- *Narotsky MG, Kavlock RJ. 1995. A multidisciplinary approach to toxicological screening: II. Development toxicity. *J Tox Environ Health* 45:145-171.
- *NAS. 1980. Drinking water and health. Volume 3. Washington, DC: National Academy of Science, Safe Drinking Water Committee, National Academy Press.
- *NAS/NRC. 1989. Biologic markers in reproductive toxicology. National Academy of Sciences/National Research Council. Washington, DC: National Academy Press, 15-35.
- *NATICH. 1994. Database report on state, local, and EPA air toxics activities. National Air Toxics Information Clearinghouse. US Environmental Protection Agency, Office of Air Quality Planning and Standards. Washington, DC.
- *NCI. 1977. Bioassay of tetrachloroethylene for possible carcinogenicity. National Cancer Institute. U.S. Department of Health, Education, and Welfare, Public Health Service, National Institutes of Health, DHEW Publ (NIH) 77-813.
- *Neely WB, Branson DR, Blau GE. 1974. Partition coefficients to measure bioconcentration potential of organic chemicals in fish. *Environ Sci Technol* 8:1113-1115
- *Nelson BK, Taylor BJ, Setzer JV, et al. 1980. Behavioral teratology of perchloroethylene in rats. *J Environ Pathol Toxicol* 3:233-250.
- *Newcomb PA, Carbone PP. 1992. The health consequences of smoking: Cancer. *Med Clin North Am* 76(2):305-331.
- *Nicoara S, Culea M, Paliobroda N, et al. 1994. Volatile organic chemical pollutants in laboratory indoor air. *Indoor Environment* 3(3):83-86.
- Niedzielski JJ, Schram RM, Phelps TJ, et al. 1989. A total-recycle expanded-bed bioreactor design which allows direct headspace sampling of volatile chlorinated aliphatic-compounds. *J Microbial Methods* 10:215-223.
- *NIOSH. 1978. Current Intelligence Bulletin 20. Tetrachloroethylene (perchloroethylene). Washington, DC: U.S. Department of Health, Education, and Welfare, National Institute for Occupational Safety and Health (NIOSH) publication no. 78-112. January 20, 1978, 1-2.
- *NIOSH. 1980. Teratogenic-mutagenic risk of workplace contaminants: Trichloroethylene, perchloroethylene, and carbon disulfide. Report to National Institute for Occupational Safety and Health, Cincinnati, OH by Litton Bionetics, Inc., Kensington, MD. Publication no. 82-185075.
- *NIOSH. 1994a. Documentation for immediately dangerous to life or health concentrations (IDLHs). National Institute for Occupational Safety and Health. Cincinnati, OH. PB94195047.
- *NIOSH. 1994b. NIOSH manual of analytical methods. Third edition. Second supplement. Cincinnati, OH: NIOSH, U.S. Department of Health and Human Services, National Institute for Occupational Safety and Health, NIOSH method no. 1003.

8. REFERENCES

- *NIOSH. 1994c. NIOSH pocket guide to chemical hazards. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, 300-301, 342.
- *NOES. 1990. National Occupational Exposure Survey (1981-1983). U.S. Department of Health and Human Services, National Institute for Occupational Safety and Health, Cincinnati, OH. July 1, 1990.
- *NTD. 1995. The National Trade Database: Import and export data for tetrachloroethylene. U.S. Department of Commerce, Economics and Statistics Administration.
- *NTP. 1986. National Toxicology Program--technical report series no. 311. Toxicology and carcinogenesis studies of tetrachloroethylene (perchloroethylene) (CAS No. 127- 18-4) in F344/N rats and B6C3F₁ mice (inhalation studies). Research Triangle Park, NC: U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health, NIH publication no. 86-2567.
- *Odum J, Green T, Foster JR, et al. 1988. The role of trichloroacetic acid and peroxisome proliferation in the differences in carcinogenicity of perchloroethylene in the mouse and rat. *Toxicol Appl Pharmacol* 92:103-112.
- *Ogata M, Takasuka Y, Tomokuni K, et al. 1971. Excretion of organic chlorine compounds in the urine of persons exposed to vapours of trichloroethylene and tetrachloroethylene. *Br J Ind Med* 28:386-390.
- *Ohio River Valley Sanitation Commission. 1980. Valley Water Sanitation Commission. Assessment of water quality conditions. Ohio River Mainstem 1978-79. Cincinnati, OH.
- OHM/TADS. 1990. Oil and hazardous materials/Technical Assistance Data System. Baltimore, MD: Chemical Information Systems, Inc. September, 1986.
- *Ohtsuki T, Sato K, Koizumi A, et al. 1983. Limited capacity of humans to metabolize tetrachloroethylene. *Int Arch Occup Environ Health* 5 1:381-390.
- *Okouchi S. 1986. Volatilization coefficient for stripping trichloroethylene, 1,1,1-trichloroethylene, and tetrachloroethylene from water. *Water Sci Technol* 181137-138.
- *Olsen GW, Heam S, Cook RR, et al. 1989. Mortality experience of a cohort of Louisiana USA chemical workers. *J Occup Med* 31:32-34.
- *Olsen J, Hemminki K, Ahlborg G, et al. 1990. Low birthweight, congenital malformations, and spontaneous abortions among dry cleaning workers in Scandinavia. *Stand J Work Environ Health* 16:163-168.
- *Olson MJ, Johnson JT, Reidy CA. 1990. A comparison of male rat and human urinary proteins: Implications for human resistance to hyaline droplet nephropathy. *Toxicol Appl Pharmacol* 102:524-536.
- *Opdam JJG, Smolders JFJ. 1986. Alveolar sampling and fast kinetics of tetrachloroethene in man. I. Alveolar Sampling. *Br J Ind Med* 43:814-824.

8. REFERENCES

- *Oppelt ET. 1987. Incineration of hazardous waste. A critical review. *J Air Pollut Control Assoc* 37:558-586.
- *OSHA. 1993. Air contaminants. Occupational Safety and Health Administration. Federal Register 58(124). 29 CFR 1910.
- *Otson R, Williams DT. 1982. Headspace chromatographic determination of water pollutants. *Anal Chem* 54:942-946
- *Otson R, Williams DT, Bothwell PD. 1982. Volatile organic compounds in water at thirty Canadian potable water treatment facilities. *J Assoc Off Anal Chem* 65: 1370-1374.
- *Otto D, Hudnell K, Boyes W. 1988. Electrophysiological measures of visual and auditory function as indices of neurotoxicity. *Toxicology* 49:205-218.
- Pagga U. 1987. Biodegradation of substances at low concentrations. *J Water Wastewater Res* 20:101-107.
- *Page GW. 1981. Comparison of groundwater and surface water for patterns and levels of contamination by toxic substances. *Environ Sci Technol* 15:1475-1481.
- *Pankow JF, Rosen ME. 1988. Determination of volatile compounds in water by purging directly to a capillary column with whole column cryotrapping. *Environ Sci Technol* 22:398-405.
- *Park KS, Sorenson DL, Sims JL. 1988. Volatilization of wastewater trace organics in slow rate land treatment systems. *Hazardous Waste and Hazardous Materials* 5:219-229.
- Parker GS, Rosen SL. 1981. Woburn: Cancer incidence and environmental hazards 1969-1978. Atlanta, GA: Massachusetts Department of Health, PHS-CDC. EPI-80-37-2.
- *Parsons F, Barrio-Lage G, Rice R. 1985. Biotransformation of chlorinated organic solvents in static microcosms. *Environ Toxicol Chem* 4:739-742.
- *Parsons F, Wood PR, Demarco J. 1984. Transformation of tetrachloroethene and trichloroethene in microcosms and groundwater. *J Am Water Works Assoc* 76:56-59.
- *Pate1 R, Janakiraman N, Johnson R, et al. 1973. Pulmonary edema and coma from perchloroethylene. *J Am Med Assoc* 223:1510.
- *Pearson CR, McConnell G. 1975. Chlorinated C1 and C2 hydrocarbons in the marine environment. *Proc R Soc Lond [Biol]* 189:305-332.
- *Peers AM. 1985. Method 5: Determination of trichloroethylene and tetrachloroethylene in air. International Agency for Research on Cancer Science Publication 68:205-211.
- *Pegg DG, Zempel JA, Braun WH, et al. 1979. Disposition of (¹⁴C) tetrachloroethylene following oral and inhalation exposure in rats. *Toxicol Appl Pharmacol* 5 1:465-474.

8. REFERENCES

- *Pekari K, Aitio A. 1985a. Method 29: Determination of trichloroacetic acid in urine. International Agency for Research on Cancer Science Publication 68:467-472.
- *Pekari K, Aitio A. 1985b. Method 30: Determination of 2,2,2-trichloroethanol in urine. International Agency for Research on Cancer Science Publication 68:473-478.
- Pellizzari ED, Bunch JE. 1979. Ambient air carcinogenic vapors: Improved sampling and analytical techniques and field studies. EPA 600/2-79-08 1.
- Pellizzari ED, Hartwell TD, Harris BS, et al. 1982. Purgeable organic compounds in mother's milk. Bull Environ Contam Toxicol 28:322-328.
- Pellizzari ED, Hartwell TD, Perritt RL, et al. 1986. Comparison of indoor and outdoor residential levels of volatile organic chemicals in five USA geographical areas. Environ Int 12:619-624.
- *Pereira MA. 1996. Carcinogenic activity of dichloroacetic acid and trichloroacetic acid in the liver of female B6C3F1 mice. Fund Appl Toxicol 31: 192-199.
- *Pezzagno G, Imbriani M, Ghittori S, et al. 1988. Urinary concentration, environmental concentration, and respiratory uptake of some solvents: Effect of the Work Load. Am Ind Hyg Assoc J 49:546-552.
- *Piet GJ, Morra CHF, DeKruif HAM. 1981. The behavior of organic micropollutants during passage through the soil. Quality of Groundwater, Proceedings of an International Symposium. Studies in Environ Science 17:557-564.
- *Plumb RH, Jr. 1987. A comparison of groundwater monitoring data from CERCLA and RCRA sites. Ground Water Monit Rev 7:94-100.
- *Pocklington DW. 1992. Determination of tetrachloroethylene - Results of a collaborative study and the standardized method. Journal of the Am Oil Chem Soc 69(8):789-793.
- Pohland FG, Schaffer TR, Yaris S, et al. 1987. Fate of selected organic pollutants during landfill disposal operations. AD-A180117.
- Popp W, Muller G, Baltes-Schmitz B, et al. 1992. Concentrations of tetrachloroethene in blood and trichloroacetic acid in urine of workers and neighbors of dry cleaning shops. Int Arch Occup Environ Health 63:393-395.
- *Prentice, RL. 1986. Comment. Journal of the American Statistical Association 81(395):600-601.
- *Price PJ, Hassett CM, Mansfield JI. 1978. Transforming activities of trichloroethylene -and proposed industrial alternatives. *In vitro* 14:290-293.
- *Price RG, Taylor SA, Crutcher E, et al. 1995. The assay of laminin fragments in serum and urine as an indicator of renal damage induced by toxins. Toxicol Let 77:313-318.
- Rampy LW, Quast JF, Balmer MF, et al. 1978. Results of a long-term inhalation toxicity study. Perchloroethylene in rats. Toxicology Research Laboratory, Health and Environmental Research, The Dow Chemical Company, Midland, MI. Unpublished.

8. REFERENCES

Rampy LW, Quast JF, Leong BKJ, et al. 1978. Results of long-term inhalation toxicity studies on rats of 1,1,1-trichloroethane and perchloroethylene formulations. Proceedings of the First International Congress on Toxicology 562.

*Ramsey JD, Flanagan RJ. 1982. Detection and identification of volatile organic compounds in blood by headspace gas chromatography as an aide to the diagnosis of solvent abuse. *J Chromatogr* 240:423-444.

*Rae HV, Brown DR. 1993. A physiologically based pharmacokinetic assessment of tetrachloroethylene in groundwater for a bathing and showering determination. *Risk Anal* 13(1):37-49.

*Rae VR, Levy K, Lustik M. 1993. Logistic regression of inhalation toxicities of perchloroethylene-Application in noncancer risk assessment. *Regulatory Toxicology and Pharmacology* 18:233-247.

*Rasmussen RA, Harsch DE, Sweany PH, et al. 1977. Determination of atmospheric halocarbons by a temperature-programmed gas chromatographic freezeout concentration method. *J Air Pollut Control Assoc* 27:579-581.

*Ratnoff WD, Gress RE. 1980. The familial occurrence of polycythemia Vera: Report of a father and a son, with consideration of the possible etiologic role of exposure to organic solvents, including tetrachloroethylene. *Blood* 56:233-236.

*Rea WJ, Pan Y, Laseter JL, et al. 1987. Toxic volatile organic hydrocarbons in chemically sensitive patients. *Clin Ecol* 5:70-74.

Reichert D. 1983. Biological actions and interactions of tetrachloroethylene. *Mutat Res* 123:411-429.

*Reinhard M, Goodman NL, Barker JF. 1984. Occurrence and distribution of organic chemicals in two landfill leachate plumes. *Environ Sci Technol* 18:953-961.

*Reinhardt CF, Mullin LS, Maxfield ME. 1973. Epinephrine-induced cardiac arrhythmia potential of some common industrial solvents. *J Occup Med* 15:953-955.

*Reitz, RH, Gargas ML, Mendrala AL, et al. 1996. In vivo and in vitro studies of perchloroethylene metabolism for physiologically based pharmacokinetic modeling in rats, mice, and humans. *Toxicol Appl Phamacol* 136:289-306.

Reitz RH, Quast JF, Schumann AM, et al. 1980. Nonlinear pharmacokinetic parameters need to be considered in high-dose/low-dose extrapolation. *Arch Toxicol Suppl* 3:79-94.

*Riihimaki V, Pfaffli, P. 1978. Percutaneous absorption of solvent vapors in man. *Stand J Work Environ Health* 4:73-85.

Robb EV, McGaughey JF, Sykes AL, et al. 1986. Recovery of principal organic hazardous constituents and products of incomplete combustion from a volatile organic sampling train. Environmental Protection Agency, Research Triangle Park, NC, Air and Energy Engineering Research Lab. EPA/600/7-86/1025.

8. REFERENCES

- *Roberts PV, Dandliker PG. 1983. Mass transfer of volatile organic contaminants from aqueous solution to the atmosphere during surface aeration. *Environ Sci Technol* 17:484-489.
- *Roberts PV, Goltz MN, Mackay DM. 1986. A natural gradient experiment on solute transport in a sand aquifer. 3. Retardation estimates and mass balance for organic solutes. *Water Resources Research* 22(13):2047-2058.
- *Rogan WJ. 1986. Comment. *Journal of the American Statistical Association* 81(395):602-603.
- *Rosengren LE, Kjellstrand P, Haglid KG. 1986. Tetrachloroethylene: Levels of DNA and S-100 in the gerbil CNS after chronic exposure. *Neurobehav Toxicol Teratol* 8:201-206.
- *Rowe VK, McCollister DD, Spencer HC, et al. 1952. Vapor toxicity of tetrachloroethylene for laboratory animals and human subjects. *AMA Arch Ind Hyg Occup Med* 5:566-579.
- *Ruder AM, Ward EM, Brown DP. 1994. Cancer mortality and female and male dry cleaning workers. *J Occup Environ Med* 36(8):867-874.
- *Sahel GV, Clark TP. 1984. Volatile organic compounds as indicators of municipal solid waste leachate contamination. *Waste Management and Research* 2:119-130.
- *Saisho. 1994. Bioaccumulation of volatile chlorinated hydrocarbons in blue mussel, *Mytilus edulus*, and killfish, *Oryzias latipes*. *Japanese Journal of Toxicology and Environmental Health* 40:274-278.
- *Saland G. 1967. Accidental exposure to perchloroethylene. *N Y State J Med* 67:2359-2361.
- *Sallmen M, Lindbohm ML, Kyyronen P, et al. 1995. Reduced fertility among women exposed to organic solvents. *Am J Ind Med* 27:699-713.
- *Sandground JH. 1941. Coma following medication with tetrachloroethylene. *J Am Med Assoc* 117:440-441.
- SANSS. 1987. Chemical Information System (CIS). Structure and Nomenclature Search System.
- *Sauer TC Jr. 1981. Volatile organic compounds in open ocean and coastal surface waters. *Organic Geochemistry* 3:91-101.
- *Savolainen H, Pfaffli P, Tengen M, et al. 1977. Biochemical and behavioral effects of inhalation exposure to tetrachloroethylene and dichloromethane. *J Neuropathol Exp Neurol* 36:941-949.
- *Sax NI, Lewis RJ, eds. 1987. *Hawley's condensed chemical dictionary*. New York, NY: Van Nostrand Reinhold, 11:886.
- *Schlichting LM, Wright PFA, Stacey NH. 1992. Effects of tetrachloroethylene on hepatic and splenic lymphocytotoxic activities in rodents. *Toxicology and Industrial Health* 8(5):255-266.
- *Schreiber JS. 1993. Predicted infant exposure to tetrachloroethylene in human breast milk. *Risk Analysis* 13(5):515-524.

8. REFERENCES

- *Schreiber JS, House S, Prohonic E, et al. 1993. An investigation of indoor air contamination in residences above dry cleaners. *Risk Anal* 13(3):335-344.
- *Schultz B, Kjeldsen P. 1986. Screening of organic matter in leachates from sanitary landfills using gas chromatography combined with mass spectrometry. *Water Res* 20:965-970.
- *Schumann AM, Quast JF, Watanabe PG. 1980. The pharmacokinetics and macromolecular interactions of perchloroethylene in mice and rats as related to oncogenicity. *Toxicol Appl Pharmacol* 55:207-219.
- *Schwarzenbach RP, Giger W, Hoehn E, Schneider JK. 1983. Behavior of organic compounds during infiltration of river water to ground water. Field studies. *Environmental Science and Technology* 17:472-479.
- Schwarzenbach RP, Westall J. 1981. Transport of nonpolar organic compounds from surface water to groundwater. Laboratory sorption studies. *Environ Sci Technol* 15:1360-1367.
- *Schwarzenbach RP, Molnar-Kubica E, Giger W, et al. 1979. Distribution, residence time, and fluxes of tetrachloroethylene and 1,2-dichlorobenzene in Lake Zurich, Switzerland. *Environ Sci Technol* 13:1367-1373.
- *Schwetz BA, Leong BKJ, Gehring PJ. 1975. The effect of maternally inhaled trichloroethylene, perchloroethylene, methyl chloroform, and methylene chloride on embryonal and fetal development in mice and rats. *Toxicol Appl Pharmacol* 32:84-96.
- *Seeber A. 1989. Neurobehavioral toxicity of long-term exposure to tetrachloroethylene. *Neurotoxicol Teratol* 11:579-583.
- *Seidel HJ, Weber L, Barthel E. 1992. Hematological toxicity of tetrachloroethylene in mice. *Arch Toxicol* 66:228-230.
- *Seiji K, Inoue O, Jin C, et al. 1989. Dose-excretion relationship in tetrachloroethylene-exposed workers and the effect of tetrachloroethylene co-exposure on trichloroethylene metabolism. *Am J Ind Med* 16:675-684.
- *Seiji K, Jin C, Watanabe T, et al. 1990. Sister chromatid exchanges in peripheral lymphocytes of workers exposed to benzene, trichloroethylene, or tetrachloroethylene, with references to smoking habits. *Int Arch Occup Environ Health* 62: 171-176.
- *Seip HM, Alstad J, Carlberg GE, et al. 1986. Measurement of mobility of organic compounds in soils. *Sci Total Environ* 50:87-101.
- Seppalainen AM, Antti-Poika M. 1983. Time course of electrophysiological findings for patient with solvent poisoning. *Stand J Work Environ Health* 9:15-24.
- *Shipman AJ, Whim BP. 1980. Occupational exposure to trichloroethylene in metal cleaning processes and to tetrachloroethylene in the dry cleaning industry in the U.K. *Ann Occup Hyg* 23:197-204.

8. REFERENCES

- *Singh HB, Lillian D, Appleby A, et al. 1975. Atmospheric formation of carbon tetrachloride from tetrachloroethylene. *Environ Lett* 10:253-256.
- *Singh HB, Salas LJ, Shigeishi H, et al. 1979. Atmospheric distributions, sources and sinks of selected halocarbons, hydrocarbons, SF₆ and N₂O. Research Triangle Park, NC: Environmental Protection Agency, Environmental Sciences Research Laboratory, Office of Research and Development, 24-33. EPA 600/3-79-107.
- Singh HB, Salas LJ, Smith AJ, et al. 1981. Measurements of some potentially hazardous organic chemicals in urban environments. *Atmos Environ* 15:601-612.
- *Singh HB, Salas LJ, Stiles RE. 1982. Distribution of selected gaseous organic mutagens and suspect carcinogens in ambient air. *Environ Sci Technol* 16:872-880.
- Sittig M. 1985. *Handbook of Toxic and Hazardous Chemicals and Carcinogens*. Park Ridge, NJ: Noyes Publications, 843-845.
- *Skender L, Karacic V, Bosner B, et al. 1993. Assessment of exposure to trichloroethylene and tetrachloroethylene in the population of Zagreb, Croatia. *Int Arch Occup Environ Health* 65:S163-S165.
- *Skender L, Karacic V, Bosner B, et al. 1994. Assessment of urban population exposure to trichloroethylene and tetrachloroethylene by means of biological monitoring. *Arch Environ Health* 49(6):445-451.
- *Skender LJ, Karacic V, Prpic-Majic D. 1991. A comparative study of human levels of trichloroethylene and tetrachloroethylene after occupational exposure. *Archives of Environmental Health* 46(3):174-178.
- *Skender L, Karacic V, Prpic-Majic D. 1987. Occupational exposure indicators to tetrachloroethylene in a dry cleaning shop. In: *Occu Environ Chem Hazard Proc Int Symp, Biochem Cell Indices Hum Toxic Occup Environ Med* 192:6.
- Smith LR, Dragun J. 1984. Degradation of volatile chlorinated aliphatic priority pollutants in groundwater. *Environment International* 10:291-298.
- *Smith EM, Miller ER, Woolson RF, et al. 1985. Bladder cancer risk among laundry workers, dry cleaners, and others in chemically related occupations. *J Occup Med* 27:295-297.
- *Solet D, Robins TG. 1991. Renal function in dry cleaning workers exposed to perchloroethylene. *American Journal of Industrial Medicine* 20:601-614.
- Solet D, Robins TG, Sampaio C. 1990. Perchloroethylene exposure assessment among dry cleaning workers. *Am Ind Hyg Assoc J* 51(10):566-574
- *Soni M, Nomiya H, Nomiya K. 1990. Chronic inhalation effects of tetrachloroethylene on hepatic and renal microsomal electron transport components and 6aminolevulinic acid dehydratase in rats. *Toxicology Letters* 54:207-213.

8. REFERENCES

- Spee T. 1986. Evaluation of an ISO draft proposal for sampling and analysis of chlorinated hydrocarbon solvent vapors in workplace atmospheres. *Am Ind Hyg Assoc J* 47:27-36.
- *Spencer AB, Earnest GS, Kovein RJ. 1995. In-depth survey report: Exposures during the spotting procedure in a commercial dry cleaner at Widmer's Dry Cleaning, Cincinnati, Ohio. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Report No. ECTB 201-13b. 27 pages.
- Spicer CW, Holdren MW, Slivon LE, et al. 1986. Intercomparison of sampling techniques for toxic organic compounds in indoor air. Proceedings of the 1986 EPA/APCA Symposium on Measurement of Toxic Air Pollutants, Raleigh, NC, April 27-30, 1986, Environmental Monitoring Systems Laboratory, U.S. Environmental Protection Agency, EPA Report No. 600/9-86-013, 45-60.
- *Spirtas R, Stewart PA, Lee JS, et al. 1991. Retrospective cohort mortality study of workers at an aircraft maintenance facility. I. Epidemiological results. *Br. J. Ind. Med.* 48(8):515-530.
- *SRI. 1995. Directory of Chemical Producers, United States of America. SRI International, Menlo Park, California, 794.
- *Staples CA, Werner AF, Hoogheem TJ. 1985. Assessment of priority pollutant concentrations in the United States using STORET database. *Environ Toxicol Chem* 4:131-142.
- *Stemhagen A, Slade J, Altman R et al. 1983. Occupational risk factors and liver cancer: A retrospective case-control study of primary liver cancer in New Jersey. *Am J Epidemiol* 117:443-454.
- Stetter JR, Jurs PC, Rose SL. 1986. Detection of hazardous gases and vapors: Pattern recognition analysis of data from an electrochemical sensor array. *Anal Chem* 58:860-866.
- *Stewart RD. 1969. Acute tetrachloroethylene intoxication. *The Journal of the American Medical Association* 208:1490-1492.
- *Stewart RD, Baretta ED, Dodd HC, et al. 1970. Experimental human exposure to tetrachloroethylene. *Arch Environ Health* 20:224-229.
- *Stewart RD, Dodd HC. 1964. Absorption of carbon tetrachloride, trichloroethylene, tetrachloroethylene, methylene chloride, and 1,1,1-trichloroethane through the human skin. *Am Ind Hyg Assoc J* 25:439-446.
- *Stewart RD, Erley DS, Schaffer AW, et al. 1961a. Accidental vapor exposure to anesthetic concentrations of a solvent containing tetrachloroethylene. *Indust Med Surg* 30:327-330.
- *Stewart RD, Gay HH, Erley DS, et al. 1961b. Human exposure to tetrachloroethylene vapor: Relationship of expired air and blood concentrations to exposure and toxicity. *Arch Environ Health* 2:516-522.
- *Stewart RD, Hake CL, Forster HV, et al. 1981. Tetrachloroethylene: Development of a biologic standard for the industrial worker by breath analysis. Cincinnati, OH: National Institute for Occupational Safety and Health. Contract no. HSM 99-72-84. NIOSH-MCOW-ENVM-PCE-74-6. NTIS No. PB82-152166

8. REFERENCES

- *Stewart RD, Hake CL, Wu A, et al. 1977. Effects of perchloroethylene/druug interaction on behavior and neurological function. Final report. National Institute for Occupational Safety and Health, Washington, DC. PB83-17460.
- *Story DL, Meierhenry EF, Tyson CA, et al. 1986. Differences in rat liver enzyme-altered foci produced by chlorinated aliphatics and phenobarbital. *Toxicol Ind Health* 2:351-362.
- *Strachan WMJ, Edwards CJ. 1984. Organic pollutants in Lake Ontario. In: Nriagu JO, Simmons MS, eds. *Toxic contaminants in the Great Lakes*. New York, NY: John Wiley and Sons, 239-264.
- *Stutz DR, Ulin S. 1992. *Hazardous materials injuries: A handbook for pre-hospital care*. Third edition. Beltsville, MD: Bradford Communications Corporations, 286-287.
- *Su C, Goldberg ED. 1976. Environmental concentration and fluxes of some halocarbons. In: Windom HL, et al., eds. *Marine Pollutant Transfer*, Lexington, MA, 353-374.
- *Suber RL. 1989. Clinical pathology for toxicologists. In: Hayes AW, ed. *Principles and methods of toxicology*. 2nd ed. New York, NY: Raven Press, 485-519.
- *Sullivan DA, Jones AD, Williams JG. 1985. Results of the U.S. Environmental Protection Agency's Air Toxics Analysis in Philadelphia. In: *Proceedings of the 78th Annual Meeting of the Air Pollution Control Association*. Vol. 2, 1-15.
- *Swan SH, Robins JM. 1986. Comment. *Journal of the American Statistical Association* 81(395):604-608.
- *Swann RL, Laskowski DA, McCall PJ, et al. 1983. A rapid method for the estimation of the environmental parameters octanol/water partition coefficient, soil sorption constant, water to air ratio, and water solubility. *Res Rev* 85:17-28.
- *Swenberg JA, Short B, Borghoff S, et al. 1989. The comparative pathobiology of a α -2 μ globulin nephropathy. *Toxicol Appl Pharmacol* 97:35-46.
- *Tabak HH, Quave SA, Mashni CI, et al. 1981. Biodegradability studies with organic priority pollutant compounds. *J Water Pollut Control Fed* 53:1503-1518.
- *Thomas KW, Pellizzari ED, Perritt RL. 1991. Effect of dry-cleaned clothes on tetrachloroethylene levels in indoor air, personal air, and breath for residents of several New Jersey homes. *Journal of Exposure Analysis and Environmental Epidemiology* 1(4):475-490.
- *Thomas RG. 1982. Volatilization from water. In: Lyman WJ, Reehl WF, Rosenblatt DH, eds. *Handbook of chemical property estimation methods: Environmental behavior of organic compounds*. New York, NY: McGraw-Hill, 1-2, 13, 25.
- *Thompson KM, Evans JS. 1993. Workers' breath as a source of perchloroethylene (pert) in the home. *Journal of Exposure Analysis and Environmental Epidemiology* 3(4):417-430.
- *Tichenor BA, Sparks LE, Jackson MD, et al. 1990. Emissions of perchloroethylene from dry cleaned fabrics. *Atmospheric Environment* 24A(5):1219-1229.

8. REFERENCES

- *Tinston DJ. 1995. Perchloroethylene: Multigeneration inhalation study in the rat. Report No: CTL/P/4097. Sponsored by Halogenated Solvents Industry Alliance, HSIA/90/0002.
- *Travis CC, White RK, Arms AD. 1989. A physiologically based pharmacokinetic approach for assessing the cancer risk of tetrachloroethylene. In: Paustenbach DJ ed. The Risk Assessment of Environmental and Human Health Hazards: A Textbook of Case Studies. John Wiley & Sons, New York 769-796.
- Travis CC, Hattemer-Frey HA, ARMS AD. 1988. Relationships between dietary intake of organic chemicals and their concentrations in human adipose tissue and breast milk. Arch Environ Contam Toxicol 17:473-478.
- *TRI93. 1995. Toxic Chemical Release Inventory. National Library of Medicine, National Toxicology Information Program, Bethesda, MD.
- *TRI88. 1990. Toxic Chemical Release Inventory. National Library of Medicine, National Toxicology Information Program, Bethesda, MD.
- *Tsulaya VR, Bonashevskaya TI, Zykova VV, et al. 1977. [The toxicological characteristics of some chlorinated hydrocarbons]. Gig Sanit 8:50-53. (Russian)
- *Tsuruta H. 1975. Percutaneous absorption of organic solvents. (1) Comparative study of the *in vivo* percutaneous absorption of chlorinated solvents in mice. Ind Health 13:227-236.
- *Tsuruta H. 1977. Percutaneous absorption of organic solvents. (2) A method for measuring the penetration rate of chlorinated solvents through excised rat skin. Ind Health 15:131-139.
- *Tsuruta H. 1989. Skin absorption of organic solvent vapors in nude mice *in vivo*. Ind Health 27:37-47.
- *Tu AS, Murray TA, Hatch KM, et al. 1985. *In vitro* transformation of BALB/C-3T3 cell by chlorinated ethanes and ethylenes. Cancer Lett 28:85-92.
- *Urano K, Murata C. 1985. Adsorption of principal chlorinated organic compounds on soil. Chemosphere 14:293-299.
- Vahdat N. 1987. Permeation of protective clothing materials by methylene chloride and perchloroethylene. Am Ind Hyg Assoc J 48:646-651.
- *Valencia R, Mason JM, Woodruff RC, et al. 1985. Chemical mutagenesis testing in *Drosophila* III. Results of 48 coded compounds tested for the National Toxicology Program. Environ Mutagen 7:325-348.
- *Vamvakas S, Herkenhoff M, Dekant W, et al. 1989. Mutagenicity of tetrachloroethene in the Ames test: Metabolic activation by conjugation with glutathione. J Biochem Toxicol 4:21-27.
- Van Der Gulden JWJ, Zielhuis GA. 1989. Reproductive hazards related to perchloroethylene. A review. Int Arch Occup Environ Health 61:235-242.

8. REFERENCES

- *Van Duuren BL, Goldschmidt BM, Loewengart G, et al. 1979. Carcinogenicity of halogenated olefinic and aliphatic hydrocarbons in mice. *J Natl Cancer Inst* 63:1433-1439.
- *Van Rillaer W, Beemaert H. 1989. Determination of residual tetrachloroethylene in olive oil by headspace-gas chromatography. *Z Lebensm Unters Forsch* 188:221-222.
- *Vartianinen T, Pukkala E, Rienoja T, et al. 1993. Population exposure to tri- and tetrachloroethene and cancer risk: Two cases of drinking water pollution. *Chemosphere* 27(7):1171-1181.
- *Veith GD, Macek KJ, Petrocelli SR, et al. 1980. An evaluation of using partition coefficients and water solubility to estimate bioconcentration factors for organic chemicals in fish. In: Eaton JG, Parrish PR, Hendricks AC, eds. *Aquatic Toxicology*. Am Soc Test Mater, ASTM STP 707, 116-129.
- *Vemmer H, Rohleder K, Wiczorek H, et al. 1984. [Perchloroethylene in the tissues of fattening pigs]. *Landwirtsch Forsch* 37:36-43. (German)
- *Verschueren K. 1983. *Handbook of environmental data on organic chemicals*. New York, NY: Van Nostrand Reinhold, 1076-1080.
- Vessel ES, Beam AG. 1962. Variations in the lactic dehydrogenase of vertebrate erythrocytes, *J Gen Physiol* 45:553-564.
- *Viau C, Bernard A, Lauwerys R, et al. 1987. A cross-sectional survey of kidney function in refinery employees. *Am J Ind Med* 11:177-187.
- *Vogel TM, McCarty PL. 1985. Biotransformation of tetrachloroethylene to trichloroethylene, dichloroethylene, vinyl chloride and carbon dioxide under methanogenic conditions. *Appl Environ Microbiol* 49:1080-1083.
- *Vyskocil A, Emminger S, Tejral J. 1990. Study on kidney function in female workers exposed to perchloroethylene. *Human & Experimental Toxicology* 9:377-380.
- *WAC. 1985. Groundwater quality. Chapter NR 140. Wisconsin Administrative Code. Department of Natural Resources.
- *WAC. 1988. Control of hazardous pollutants. Chapter NR 445. Wisconsin Administrative Code. Department of Natural Resources.
- *Wakeham SG, Davis AC, Karas JL. 1983. Mesocosm experiments to determine the fate and persistence of volatile organic compounds in coastal seawater. *Environ Sci Technol* 17:611-617.
- *Wakeham SG, Davis AC, Witt RT, et al. 1980. Tetrachloroethylene contamination of drinking water by vinyl-coated asbestos-cement pipe. *Bull Environ Contam Toxicol* 25:639-645.
- *Wallace LA. 1986. Personal exposures, indoor and outdoor air concentrations and exhaled breath concentrations of selected volatile organic compounds measured for 600 residents of New Jersey, North Dakota, North Carolina, and California. *Toxicol Environ Chem* 12:215-236.

8. REFERENCES

- *Wallace LA, Pellizzari ED, Hartwell TD, et al. 1985. Personal exposures, indoor-outdoor relationships, and breath levels of toxic air pollutants measured for 355 persons in New Jersey. *Atmos Environ* 19:1651-1661.
- *Wallace LA, Pellizzari ED, Hartwell TD, et al. 1986a. Concentrations of 20 volatile organic compounds in the air and drinking water of 350 residents of New Jersey compared with concentrations in their exhaled breath. *J Occup Med* 28:603-608.
- *Wallace LA, Pellizzari ED, Hartwell TD, et al. 1986b. Results from the first three seasons of the TEAM study: personal exposures, indoor-outdoor relationships, and breath levels of toxic air pollutants measured for 355 persons in New Jersey. In: Kopfler PC, Craun G, eds. *Environmental Epidemiology*. Lewis Publ. 181-200.
- *Wallace LA, Pellizzari ED, Hartwell TD. 1986c. Total exposure assessment methodology (TEAM) study: Personal exposures, indoor-outdoor relationships, and breath levels of volatile organic compounds in New Jersey. *Environ Int* 12:369-387.
- *Wallace LA, Pellizzari ED, Hartwell TD, et al. 1989. The influence of personal activities on exposure to volatile organic compounds. *Environ Res* 50:37-55.
- *Wallace LA, Pellizzari ED, Leaderer B, et al. 1987. Emissions of volatile organic compounds from building materials and consumer products. *Atmos Environ* 21:385-393.
- *Wallace LA, Pellizzari ED, Sheldon L, et al. 1986d. The total exposure assessment methodology (TEAM) study: Direct measurements of personal exposures through air and water for 600 residents of several U.S. cities. In: Cohen Y, ed. *Pollutants in a Multimedia Environment*. New York, NY: Plenum Publishing Corp, 289-315.
- *Walles SAS. 1986. Induction of single-strand breaks in DNA of mice by trichloroethylene and tetrachloroethylene. *Toxicol Lett* 31:31-35.
- Walter H, Selby FW. 1966. Lactic acid dehydrogenase isoenzymes of buffy coat cells and erythrocytes from different species. *Nature* 212:613-614.
- *Wang S, Karlsson J-E, Kyrklund T, et al. 1993. Perchloroethylene-induced reduction in glial and neuronal cell marker proteins in rat brain. *Pharmacology and Toxicology* 72:273-278.
- *Ward RC, Travis CC, Hetrick DM, et al. 1988. Pharmacokinetics of tetrachloroethylene. *Toxicol Appl Pharmacol* 93:108-117.
- *Warner HP, Cohen JM, Ireland JC. 1987. Determination of Henry's Law Constants of selected priority pollutants. EPA/600/D-87/229, NTIS PB87-212684.
- *Wayne LG, Orcutt JA. 1960. The relative potentials of common organic solvents as precursors of eye-irritants in urban atmospheres. *J Occup Med* 2:383-388.
- *Weant GE, McCormick GS. 1984. Nonindustrial sources of potentially toxic substances and their applicability to source apportionment methods. EPA-450/4-84-003. NTIS PB84-231232, 34, 84.

8. REFERENCES

- *Webber T, Brown HS. 1993. Exposure to tetrachloroethylene via contaminated drinking water pipes in Massachusetts: A predictive model. *Archives of Environmental Health* 48(5):293-297.
- *Weiss NS. 1995. Cancer in relation to occupational exposure to perchloroethylene. *Cancer Causes Control* 6:257-266.
- *Wenzel DG, Gibson RD. 1951. A study of the toxicity and anthelmintic activity of n-butylidene chloride. *J Pharmacol* 3:169-176.
- *Westrick JJ, Mello JW, Thomas RF. 1984. The groundwater supply survey. *J Am Water Works Assoc* 76:52-59.
- *Whitmore AS. 1986. Comment. *J Am Statistical Assoc* 81(395):609-610.
- *WHO. 1993. *Guidelines for Drinking Water Quality*. 2nd ed. World Health Organization, Geneva, Switzerland, 63-64, 175.
- *WHO. 1987. *Tetrachloroethylene health and safety guide*. World Health Organization, Distribution and Sales 1211 Geneva 27, Switzerland, 32.
- *Wiedmann TO, Guthner B, Class TJ. 1994. Global distribution of tetrachloroethene in the troposphere: Measurements and modeling. *Environmental Science and Technology* 28:2321-2329.
- *Williams GM, Shimada T. 1983. The evaluation for genotoxicity of several halogenated solvents, *Environ Mutagen* 5 :447.
- *Wilson JT, Enfield CG, Dunlap WJ, et al. 1981. Transport and fate of selected organic pollutants in a sandy soil. *J Environ Qual* 10:501-506.
- *Wilson JT, McNabb JF, Balkwill DL, et al. 1983a. Enumeration and characterization of bacteria indigenous to a shallow water-table aquifer. *Ground Water* 21:134-142.
- *Wilson JT, McNabb JF, Wilson RH, et al. 1983b. Biotransformation of selected organic pollutants in groundwater. *Dev Ind Microbiol* 24:225-233.
- *Windham GC, Shusterman D, Swan S, et al. 1991. Exposure to organic solvents and adverse pregnancy outcome. *Am J of Ind Med* 20:241-259.
- Windholz M. 1983. *The Merck Index*. 10th ed. Merck and Co., Rahway, NJ, 1315.
- *Wright WH, Bozicevick J, Gordon LS. 1937. Studies on oxyuriasis. V. Therapy with-single doses of tetrachloroethylene. *J Am Med Assoc* 109:570-573.
- *Yagi O, Uchiyama H, Iwasaki K. 1992. Biodegradation rate of chloroethylene in soil environment. *Water Science and Technology* 25:419-424.
- *Yllner S. 1961. Urinary metabolites of ¹⁴C-tetrachloroethylene in mice. *Nature* 191:820.

8. REFERENCES

*Yuskus LR. 1984. Reducing tetrachloroethylene in vinyl-lined asbestos-cement pipes through cementmotor linings. *Journal of the American Water Works Association*, 76-81.

*Zielhuis GA, Gijsen R, Van Der Gulden JWJ. 1989. Menstrual disorders among dry cleaning workers. *Scand J Work Environ Health* 15:238.

*Ziglio G, Beltramelli G, Pegliaso F. 1984. A procedure for determining plasmatic trichloroacetic acid in human subjects exposed to chlorinated solvents at environmental levels. *Arch Environ Contam Toxicol* 13:129-134.

*Zytner RG, Biswas N, Bewtra JK. 1989a. Adsorption and desorption of perchloroethylene in soils, peat moss, and granular activated carbon. *Canadian Journal of Civil Engineering* 16:798-806.

*Zytner RG, Biswas N, Bewtra JK. 1989b. Volatilization of perchloroethylene from stagnant water and soil. In: Bell JM, ed, *Proceedings of the 43rd Industrial Waste Conference*, Purdue University, May 10-12, 1988. Chelsea, MI: Lewis Publishers, Inc. 101-108.

