5. List of References

Abbott BD, Birnbaum LS. 1989. Cellular alterations and enhanced induction of cleft palate after coadministration of retinoic acid and TCDD. Toxicol Appl Pharmacol 99:287–301.

*Abbott BD, Perdew GH, Birnbaum LS. 1994. Ah receptor in embryonic mouse palate and effects of TCDD on receptor expression. Toxicol Appl Pharmacol 126:16–25.

Adami H-O, Lipworth L, Titus-Ernstoff L, et al. 1995. Organochlorine compounds and estrogen-related cancers in women. Cancer Causes Control 6:551–566.

Andersen ME, Birnbaum LS, Barton HA, et al. 1997a. Regional hepatic CYP1A1 and CYP1A2 induction with 2,3,7,8-tetrachlorodibenzo-*p*-dioxin evaluated with a multicompartment geometric model of hepatic zonation. Toxicol Appl Pharmacol 144:145–155.

Andersen ME, Eklund CR, Mills JJ, et al. 1997b. A multicompartment geometric model of the liver in relation to regional induction of cytochrome P450s. Toxicol Appl Pharmacol 144:135–144.

Anderson H, Falk C, Fiore B, et al. 1996. Consortium for the health assessment of Great Lakes sportfish consumption. Toxicol Ind Health 12(3–4):369–373.

*Anderson HA, Falk C, Hanrahan L, et al. 1998. Profiles of Great Lakes critical pollutants: A sentinel analysis of human blood and urine. Environ Health Perspect Suppl 106(5):279–289.

*Andersson PL, Blom A, Johannisson A, et al. 1999. Assessment of PCBs and hydroxylated PCBs as potential xenoestrogens: *In vitro* studies based on MCF-7 cell proliferation and induction of vitellogenin in primary culture of rainbow trout hepatocytes. Arch Environ Contam Toxicol 37:145–150.

*Andrews JE, Courtney KD, Stead AG, et al. 1989. Hexachlorobenzene-induced hyperthyroidism and osteosclerosis in rats. Fundam Appl Toxicol 12:242–251.

*Angus WGR, Mousa MA, Vargas VM, et al. 1997. Inhibition of L-aromatic amino acid decarboxylase by polychlorinated biphenyls. Neurotoxicology 18:857–868.

*Arcaro KF, Yi L, Seegal RF, et al. 1999. 2,2',6,6'-Tetrachlorobiphenyl is estrogenic in vitro and in vivo. J Cell Biochem 72:94–102.

*Arnold DL, Bryce F, Karpinski K, et al. 1993b. Toxicological consequences of Aroclor 1254 ingestion by female Rhesus (*macaca mulatta*) monkeys. Part 1B. Prebreeding phase: Clinical and analytical laboratory findings. Food Chem Toxicol 31(11):811–824.

*Arnold DL, Bryce F, McGuire PF, et al. 1995. Toxicological consequences of Aroclor 1254 ingestion by female Rhesus (*macaca mulatta*) monkeys. Part 2. Reproduction and infant findings. Food Chem Toxicol 33:457–474.

*Cited in text

*Arnold DL, Bryce F, Stapley R, et al. 1993a. Toxicological consequences of Aroclor 1254 ingestion by female Rhesus (*macaca mulatta*) monkeys. Part 1A. Prebreeding phase: Clinical health findings. Food Chem Toxicol 31(11):799–810.

*Arnold DL, Nera EA, Stapley R, et al. 1996a. Prevalence of endometriosis in Rhesus (*macaca mulatta*) monkeys ingesting PCB (Aroclor 1254): Review and evaluation. Fundam Appl Toxicol 31(1):42–55.

Arnold DL, Stapley R, Bryce F, et al. 1998b. A multigeneration study to ascertain the toxicological effects of Great Lakes salmon fed to rats: Study overview and design. Reg Toxicol Pharmacol 27:S1–S7.

Arnold SF, Klotz DM, Collins BM, et al. 1996b. Synergistic activation of estrogen receptor with combinations of environmental chemicals. Science 272:1489–1492.

Asplund L, Svensson B, Eriksson U, et al. 1994. Polychlorinated biphenyls, 1,1,1-trichloro-2,2bis(p-chlorophenyl)ethane (p,p'-DDT) and 1,1-dichloro-2,2-bis(p-chlorophenyl)-ethylene (p,p'-DDE) in human plasma related to fish consumption. Arch Environ Health 49:477–486.

*Astroff B, Safe S. 1990. 2,3,7,8-Tetrachlorodibenzo-*p*-dioxin as an antiestrogen: Effect on rat uterine peroxidase activity. Biochem Pharmacol 39:485–488.

ATSDR. 1992a. Toxicological profile for chlorinated dibenzofurans (CDFs). Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

*ATSDR. 1992b. Public health assessment guidance manual. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

*ATSDR. 1996. Toxicological Profile for Hexachlorobenzene. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

*ATSDR. 1998. Toxicological Profile for Chlorinated Dibenzo-p-Dioxins: December 1998. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

*ATSDR. 1999a. Toxicological Profile for DDT, DDE, DDD: August, 1999: Preliminary draft for Public Comment. Atlanta, GA: Agency for Toxic Substances and Disease Registry, Public Health Service, U.S. Department of Health and Human Services.

*ATSDR. 1999b. Toxicological Profile for Mercury. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

*ATSDR. 2000. Toxicological profile for polychlorinated biphenyls. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

*ATSDR. 2001a. Draft guidance manual for the assessment of joint toxic action of chemical mixtures. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

*ATSDR. 2001b. Draft guidance manual for preparation of an interaction profile. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

ATSDR. 2001c. Interaction profile on Persistent Chemicals Found in Breast Milk: Chlorinated Dibenzop-dioxins, Hexachlorobenzene, p,p'-DDE, and Polychlorinated Biphenyls. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

*Aulerich RJ, Ringer RK. 1977. Current status of PCB toxicity to mink, and effect on their reproduction. Arch Environ Contam Toxicol 6:279–292.

*Aulerich RJ, Bursian SJ, Breslin WJ, et al. 1985. Toxicological manifestations of 2,4,5, 2',4',5'-, 2,3,6,2',3',6'-, and 3,4,5,3',4',5'-hexachlorobiphenyl and Aroclor 1254 in mink. J Toxicol Environ Health 15:63–79.

*Aulerich RJ, Bursian SJ, Evans MG, et al. 1987. Toxicity of 3,4,5,3',4',5'-hexachlorobiphenyl to mink. Arch Environ Contam Toxicol 16:53–60.

*Backlin BM, Bergman A. 1995. Histopathology of postpartum placental sites in mink (*mustela vison*) exposed to polychlorinated biphenyls or fractions thereof. APMIS 103(12):843–54.

*Backlin BM, Madej A, Forsberg M. 1997. Histology of ovaries and uteri and levels of plasma progesterone, oestradiol-17beta and oestrone sulphate during the implantation period in mated and gonadotrophin-releasing hormone-treated mink (mustela vison) exposed to polychlorinated biphenyls. J Appl Toxicol 17(5):297–306.

*Backlin BM, Persson E, Jones CJ, et al. 1998. Polychlorinated biphenyl (PCB) exposure produces placental vascular and trophoblastic lesions in the mink (*mustela vison*): a light and electron microscopic study. APMIS 106(8):785–99.

*Bager Y, Kato Y, Kenne K, et al. 1997. The ability to alter the gap junction protein expression outside GST-P positive foci in liver of rats was associated to the tumour promotion potency of different polychlorinated biphenyls. Chem Biol Interact 103(3):199–212.

*Banerjee BD, Ray A, Pasha ST. 1996. A comparative evaluation of immunotoxicity of DDT and its metabolites in rats. Indian J Exp Biol 34:517–522.

*Bannister R, Davis D, Zacherewski T, et al. 1987. Aroclor 1254 as a 2,3,7,8-tetrachlorodibenzo*p*-dioxin antagonist: Effects on enzyme induction and immunotoxicity. Toxicology 46:29–42.

*Barnett JB, Barfield L, Walls R, et al. 1987. The effect of in utero exposure to hexachlorobenzene on the developing immune response of BALB/C mice. Toxicol Lett 39:263–274.

*Barsotti DA, Marlar RJ, Allen JR. 1976. Reproductive dysfunction in Rhesus monkeys exposed to low levels of polychlorinated biphenyls (Aroclor 1248). Food Cosmet Toxicol 14:99–103.

*Battershill JM. 1994. Review of the safety assessment of polychlorinated biphenyls (PCBs) with particular reference to reproductive toxicity. Hum Exp Toxicol 13:581–597.

*Bemis JC, Seegal RF. 1999. Polychlorinated biphenyls and methylmercury act synergistically to reduce rat brain dopamine content *in vitro*. Environ Health Perspect 107:879–885.

*Bergman A, Athanasiadou M, Bergek S, et al. 1992. PCB and PCB methyl sulfones in mink treated with PCB and various PCB fractions. Ambio 21(8):570–576.

*Biegel L, Harris M, Davis D, et al. 1989b. 2,2',4,4',5,5'-Hexachlorobiphenyl as a 2,3,7,8-tetrachlordibenzo-*p*-dioxin antagonist in C57BL/6J mice. Toxicol Appl Pharmacol 97:561–571.

*Biegel L, Howie L, Safe S. 1989a. Polychlorinated biphenyl (PCB) congeners as 2,3,7,8-TCDD antagonists: Teratogenicity studies. Chemosphere 19(1–6):955–958.

*Bignert A, Olsson M, Persson W, et al. 1998. Temporal trends of organochlorines in Northern Europe, 1967–1995. Relation to global fractionation, leakage from sediments and international measures. Environ Pollut 99:177–198.

Birnbaum LS. 1994. The mechanism of dioxin toxicity: Relationship to risk assessment. Environ Health Perspect Suppl 102(9):157–167.

*Birnbaum LS, Weber H, Harris MW, et al. 1985. Toxic interaction of specific polychlorinated biphenyls and 2,3,7,8-tetrachlorodibenzo-*p*-dioxin: Increased incidence of cleft palate in mice. Toxicol Appl Pharmacol 77:292–302.

*Bitman J, Cecil HC. 1970. Estrogenic activity of DDT analogs and polychlorinated biphenyls. J Agric Food Chem 18:1108–1112.

*Bjerke DL, Sommer RJ, Moore RW, et al. 1994. Effects of *in utero* and lactational 2,3,7,8-tetrachlorodibenzo-*p*-dioxin exposure on responsiveness of the male rat reproductive system to testosterone stimulation in adulthood. Toxicol Appl Pharmacol 127:250–257.

*Bowerman WW, Best DA, Grubb TG, et al. 2000. Assessment of environmental endocrine disruptors in bald eagles of the Great Lakes. Chemosphere 1569–1574.

*Boyages SC. 2000. The neuromuscular system and brain in hypothyroidism. In: Braverman LE, Utiger RD, eds. Werner & Ingbar's the thyroid: A fundamental and clinical text. Eighth edition. Philadelphia, PA: Lippincott Williams & Wilkins, 804–810.

*Brandt I, Bergman A. 1987. PCB methyl sulphones and related compounds: Identification of target cells and tissues in different species. Chemosphere 8/9:1671–1676.

*Brezner E, Terkel J, Perry AS. 1984. The effect of Aroclor 1254 (PCB) on the physiology of reproduction in the female rat-I. Comp Biochem Physiol 77:65–70.

*Brouwer A, Morse DC, Lans MC, et al. 1998. Interactions of persistent environmental organohalogens with the thyroid hormone system: Mechanisms and possible consequences for animal and human health. Toxicol Ind Health 14(1-2):59-84.

*Brown AP, Gamey PE. 1995. Neutrophil degranulation and superoxide production induced by polychlorinated biphenyls are calcium dependent. Toxicol Appl Pharmacol 13:198–205.

Brown NM, Manzolillo PA, Zhang J-X, et al. 1998. Prenatal TCDD and predisposition to mammary cancer in the rat. Carcinogenesis 19:1623–1629.

*Buck GM, Sever LE, Mendola P, et al. 1997. Consumption of contaminated sportfish from Lake Ontario and time-to-pregnancy. Am J Epidemiol 146(11):949–954.

Burbacher TM, Mohamed MK, Mottett NK. 1988. Methylmercury effects on reproduction and offspring size at birth. Reprod Toxicol 1(4):267–278.

*Byrne JJ, Carbone JP, Hanson EA. 1987. Hypothyroidism and abnormalities in the kinetics of thyroid hormone metabolism in rats treated chronically with polychlorinated biphenyl and polybrominated biphenyl. Endocrinology 121:520–527.

*Byrne JJ, Carbone JP, Pepe MG. 1988. Suppression of serum adrenal cortex hormones by chronic low-dose polychlorobiphenyl or polybromobiphenyl treatments. Arch Environ Contam Toxicol 17:47–53.

Cantoni C, Comi G. 1997. Changes in the concentrations of pesticide residues in foods and in human tissues between 1960 and 1996. Outlook On Agriculture 26(1):47–52.

Carthew P, Smith AG. 1994. Pathological mechanisms of hepatic tumour formation in rats exposed chronically to dietary hexachlorobenzene. J Appl Toxicol 14(6):447–452.

*Chauhan KR, Kodavanti PRS, McKinney JD. 2000. Assessing the role of *ortho*-substitution on polychlorinated biphenyl binding to transthyretin, a thyroxine transport protein. Toxicol Appl Pharmacol 162:10–21.

*Cheek AO, Kow K, Chen J, et al. 1999. Potential mechanisms of thyroid disruption in humans: Interaction of organochlorine compounds with thyroid receptor, transthyretin, and thyroid-binding globulin. Environ Health Perspect 107(4):273–278.

*Choksi NY, Kodavanti PRS, Tilson HA, et al. 1997. Effects of polychlorinated biphenyls (PCBs) on brain tyrosine hydroxylase activity and dopamine synthesis in rats. Fundam Appl Toxicol 39:76–80.

*Chu I, Poon R, Yagminas A, et al. 1998. Subchronic toxicity of PCB 105 (2,3,3',4,4'-pentachlorobiphenyl) in rats. J Appl Toxicol 18:285–292.

*Chu I, Velleneuve DC, Tagimnas A, et al. 1994. Subchronic toxicity of 3,3',4,4',5-pentachlorobiphenyl in the rat. Fundam Appl Toxicol 22:457–468.

Chu I, Villeneuve DC, Valli VE, et al. 1984. Toxicological response and its reversibility in rats fed Lake Ontario or Pacific coho salmon for 13 weeks. J Environ Sci Health B 19(8–9):713–731.

*Chu I, Villeneuve DC, Yagminas A, et al. 1995. Toxicity of PCB 77 (3,3',4,4'-tetrachlorobiphenyl) and PCB 118 (2,3',4,4',5-pentachlorobiphenyl) in the rat following subchronic dietary exposure. Fundam Appl Toxicol 26:282–292.

*Chu I, Villeneuve DC, Yagminas A, et al. 1996a. Toxicity of 2,2',4,4',5,5'-hexachlorobiphenyl in rats: Effects following a 90-day oral exposure. J Appl Toxicol 16(2):121–128.

*Chu I, Villeneuve DC, Yagminas A, et al. 1996b. Toxicity of 2,4,4'-trichlorobiphenyl in rats following 90-day dietary exposure. J Toxicol Environ Health 49(3):301–318.

*Cogliano VJ. 1998. Assessing the cancer risk from environmental PCBs. Environ Health Perspect 106(6):317–323.

*Collins WT, Capen CC. 1980. Fine structural lesions and hormonal alterations in thyroid glands of perinatal rats exposed *in utero* and by the milk to polychlorinated biphenyls. Am J Pathol 99:125–142.

*Collins WT, Capen CC, Kasza L, et al. 1977. Effect of polychlorinated biphenyl (PCB) on the thyroid gland of rats: Ultrastructural and biochemical investigations. Am J Pathol 89:119–136.

*Connor K, Ramamoorthy K, Moore M, et al. 1997. Hydroxylated polychlorinated biphenyls (PCBs) as estrogens and antiestrogens: Structure-activity relationships. Toxicol Appl Pharmacol 145:111–123.

*Connor K, Safe S, Jefcoate CR, et al. 1995. Structure-dependent induction of CYP2B by polychlorinated biphenyl congeners in female Sprague-Dawley rats. Biochem Pharmacol 50(11):1913–1920.

*Cooke PS, Zhao Y-D, Hansen LG. 1996. Neonatal polychlorinated biphenyl treatment increases adult testis size and sperm production in the rat. Toxicol Appl Pharmacol 136:112–117.

*Corey DA, Juarez de Ku LM, Bingman VP, et al. 1996. Effects of exposure to polychlorinated biphenyl (PCB) from conception on growth, and development of endocrine, neurochemical, and cognitive measures in 60 day old rats. Growth Dev Aging 60:131–143.

*Courtney KD. 1979. Hexachlorobenzene (HCB): A review. Environ Res 20:225–266.

Courval JM, De Hoog JV, Stein AD, et al. 1997. Sport caught fish consumption and conception failure in Michigan andlers. Montreal, Quebec, Canada: Health Conference 1997 : Great Lakes and St. Lawrence.

*Courval JM, DeHoog JV, Stein AD, et al. 1999. Sport-caught fish consumption and conception delay in licensed Michigan anglers. Environ Res (A) 80:S183–S188.

Couture LA, Abbott BD, Birnbaum LS. 1990. A critical review of the developmental toxicity and teratogenicity of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin: Recent advances toward understanding the mechanism. Teratology 42:619–627.

Crane JL. 1996. Carcinogenic human health risks associated with consuming contaminated fish from five Great Lakes areas of concern. J Great Lakes Res 22(3):653–668.

Crisp TM, Clegg ED, Cooper RL, et al. 1998. Environmental endocrine disruption: An effects assessment and analysis. Environ Health Perspect Suppl 106(1):11–56.

*Daly HB. 1991. Reward reductions found more aversive by rats fed environmentally contaminated salmon. Neurotoxicol Teratol 13:449–453.

*Danse IR, Jaeger RJ, Kroger MK, et al. 1997. Position paper of the American Council on Science and Health: Public health concerns about environmental polychlorinated biphenyls (PCBs). Ecotoxicol Environ Saf 38:71–84.

*Dar E, Kanarek MS, Anderson HA, et al. 1992. Fish consumption and reproductive outcomes in Green Bay, Wisconsin. Environ Res 59:189–201.

*Darnerud PO, Morse D, Klasson-Wehler E, et al. 1996. Binding of a 3,3', 4,4'-tetrachlorobiphenyl (CB-77) metabolite to fetal transthyretin and effects on fetal thyroid hormone levels in mice. Toxicology 106(1–3):105–114.

Daston GP, Gooch JW, Breslin WJ, et al. 1997. Environmental estrogens and reproductive health: A discussion of the human and environmental data. Reprod Toxicol 11(4):465–481.

*Davis D, Safe S. 1989. Dose-response immunotoxicities of commercial polychlorinated biphenyls (PCBs) and their interaction with 2,3,7,8-tetrachlorodibenzo-*p*-dioxin. Toxicol Lett 48:35–43.

*Davis D, Safe S. 1990. Immunosuppressive activities of polychlorinated biphenyls in C57BL/6N mice: Structure-activity relationships as Ah receptor agonists and partial antagonists. Toxicology 63:97–111.

*DeCaprio AP, McMartin DM, O'Keefe PW, et al. 1986. Subchronic oral toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin in the guinea pig: Comparisons with a PCB-containing transformer fluid pyrolysate. Fundam Appl Toxicol 6:454–463.

*De Haan LHJ, Halfwerk S, Hovens SEL, et al. 1996. Inhibition of intercellular communication and induction of ethoxyresorufin-*O*-deethylase activity by polychlorobiphenyls, -dibenzo-*p*-dioxins and -dibenzofurans in mouse hepa1c1c7 cells. Environ Toxicol Pharmacol 1:27–37.

*DenBesten C, Bennik MH, Bruggeman I, et al. 1993. The role of oxidative metabolism in hexachlorobenzene-induced porphyria and thyroid hormone homeostasis: A comparison with pentachlorobenzene in a 13-week feeding study. Toxicol Appl Pharmacol 119:181–194.

DenBesten C, Bennik MMH, Van Iersel M, et al. 1994. Comparison of the urinary metabolite profiles of hexachlorobenzene and pentachlorobenzene in the rat. Chem Biol Interact 90:121–137.

*Den Tonkelaar EM, Verschuuren HG, Bankovska J, et al. 1978. Hexachlorobenzene toxicity in pigs. Toxicol Appl Pharmacol 43:137–145.

DeRosa CT, Johnson BL. 1996. Strategic elements of ATSDR's Great Lakes Human Health Effects Research Program. Toxicol Ind Health 12(3–4):315–325.

*Desaulniers D, Poon R, Phan W, et al. 1997. Reproductive and thyroid hormone levels in rats following 90-day dietary exposure to PCB 28 (2,4,4'-trichlorobiphenyl) or PCB 77 (3,3'4,4'-tetrachlorobiphenyl). Toxicol Ind Health 13(5):627–638.

de Solla SR, Bishop CA, Van Der Kraak G, et al. 1998. Impact of organochlorine contamination on levels of sex hormones and external morphology of common snapping turtles (*Chelydra serpentina serpentina*) in Ontario, Canada. Environ Health Perspect 106(5):253–260.

*De Vito MJ, Birnbaum LS. 1994. Toxicology of dioxins and related chemicals. In: Schecter A, ed. Dioxins and health. New York, NY: Plenum Press, 139–162.

Dickman MD, Leung KMC. 1998. Mercury and organochlorine exposure from fish consumption in Hong Kong. Chemosphere 37(5):991–1015.

*Durkin P. 1995. Development of mixtures assessment methods: Guidelines for application of the binary weight-of-evidence methodology. Submitted to The Kevric Company, Inc., Silver Spring, MD. SERA TR 95-018-001a.

Elliott JE, Norstrom RJ, Smith GEJ. 1996. Patterns, trends, and toxicological significance of chlorinated hydrocarbon and mercury contaminants in bald eagle eggs from the Pacific coast of Canada, 1990–1994. Arch Environ Contam Toxicol 31:354–367.

*EPA. 1988. Recommendations for and documentation of biological values for use in risk assessment. Cincinnati, OH: U.S. Environmental Protection Agency. PB88-179874.

*EPA. 1996. PCBs: Cancer dose-response assessment and application to environmental mixtures. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development, National Center for Environmental Assessment. EPA/600/P-96/001F.

*EPA. 1997a. Addition of dioxin and dioxin-like compounds: Modification of polychlorinated biphenyls (PCBs) listing; toxic chemical release reporting; community right-to-know. Washington, DC: U.S. Environmental Protection Agency. Federal Register. 62FR 24887.

*EPA. 1997b. Health Effects Assessment Summary Tables: FY 1997 Update. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development, Office of Emergency and Remedial Response. NTIS PB97-921199.

*Eriksson P, Norberg A. 1986. The effects of DDT, DDOH-palmitic acid, and a chlorinated paraffin on muscarinic receptors and the sodium-dependent choline uptake in the central nervous system of immature mice. Toxicol Appl Pharmacol 85:121–127.

*Eriksson P, Ahlbom J, Frederiksson A. 1992. Exposure to DDT during a defined period in neonatal life induces permanent changes in brain muscarinic receptors and behaviour in adult mice. Brain Res 582:277–281.

*Eriksson P, Nilsson-Hakansson L, Nordberg.A, et al. 1990. Neonatal exposure to DDT and its fatty acid conjugate: Effects on cholinergic and behavioural variables in the adult mouse. Neurotoxicology 11:345–354.

*Expert Panel. 1994. Interpretive review of the potential adverse effects of chlorinated organic chemicals on human health and the environment: Chapter 5: Polychlorinated biphenyls. Regul Toxicol Pharmacol 20(1):S187–S307.

Fan F, Rozman KK. 1995. Short- and long-term biochemical effects of 2,3,7,8-tetrachlorodibenzo*p*-dioxin in female Long-Evans rats. Toxicol Lett 75:209–216. *Feeley MM, Jordan SA. 1998. Dietary and tissue residue analysis and contaminant intake estimations in rats consuming diets composed of Great lakes salmon: A multigeneration study. Regul Toxicol Pharmacol 27:S8–S17.

*Feeley MM, Jordan SA, Gilman AP. 1998. The Health Canada Great Lakes Multigeneration Studysummary and regulatory considerations. Regul Toxicol Pharmacol 27:S90–S98.

*Fein GG, Jacobson JL, Jacobson SW, et al. 1984. Prenatal exposure to polychlorinated biphenyls: Effects on birth size and gestational age. J Pediatr 105(2):315–320.

*Fischer LJ, Seegal RF, Gamey PE, et al. 1998. Symposium overview: Toxicity of non-coplanar PCBs. Toxicol Sci 41:49–61.

*Fitzhugh O. 1948. Use of DDT insecticides on food products. Ind Eng Chem 40:704–705.

Foster WG. 1995. The reproductive toxicology of Great Lakes contaminants. Environ Health Perspect Suppl 103(9):63–69.

*Franklin MR, Phillips JD, Kushner JP. 1997. Cytochrome P450 induction, uroporphyrinogen decarboxylase depression, porphyin accumulation and excretion, and gender influence in a 3-week rat model of porphyria cutanea tarda. Toxicol Appl Pharmacol 147:289–299.

*Gabliks J, Al-Zubaidy T, Askari E. 1975. DDT and immunological responses. Arch Environ Health 30:81–84.

*Gamey PE, Sirosis JE, Denison M, et al. 1993. Neutrophil function after exposure to polychlorinated biphenyls *in vitro*. Environ Health Perspect 101:430–434.

*Gao X, Son D-S, Terranova PF, et al. 1999. Toxic equivalency factors of polychlorinated dibenzo-*p*-dioxins in an ovulation model: Validation of the toxic equivalency concept for one aspect of endocrine disruption. Toxicol Appl Pharmacol 157:107–116.

*Gao X, Terranova PF, Rozman KK. 2000. Effects of polychlorinated dibenzofurans, biphenyls, and their mixture with dibenzo-*p*-dioxins on ovulation in the gonadotropin-primed immature rat: Support for the toxic equivalency concept. Toxicol Appl Pharmacol 163:115–124.

*Gierthy JF, Acaro KF, Floyd M. 1997. Assessment of PCB estrogenicity in a human breast cancer cell line. Chemosphere 34(5–7):1495–1505.

*GLWQB. 1985. Report to the International Joint Commission: Report on Great Lakes water quality. Kingston, Ontario: Great Lakes Water Quality Board.

*Goldey ES, Crofton KM. 1998. Thyroxine replacement attentuates hypothyroxinemia, hearing loss, and motor deficits following developmental exposure to Aroclor 1254 in rats. Toxicol Sci 45:94–105.

*Goldey ES, Kehn LS, Lau C, et al. 1995. Development exposure to polychlorinated biphenyls (Aroclor 1254) reduces circulating thyroid hormone concentrations and causes hearing deficits in rats. Toxicol Appl Pharmacol 135:77–88.

Goldstein JA, Friesen M, Linder RE, et al. 1977. Effects of pentachlorophenol on hepatic drugmetabolizing enzymes and porphyria related to contamination with chlorinated dibenzo-*p*-dioxins and dibenzofurans. Biochem Pharmacol 26:1549–1557.

Goldstein JA, Friesen M, Scotti TM, et al. 1978. Assessment of the contribution of chlorinated dibenzo*p*-dioxins and dibenzofurans to hexachlorobenzene-induced toxicity, porphyria, changes in mixed function oxygenases, and histopathological changes. Toxicol Appl Pharmacol 46:633–649.

Goldstein JA, Linko P, Hahn ME, et al. 1986. Structure-activity relationships of chlorinated benzenes as inducers of hepatic cytochrome P-450 isozymes in the rat. In: Morris CR, Cabral JRP, ed. Hexachlorobenzene: Proceedings of an international symposium. Lyon, France: International Agency for Research on Cancer, 519–526.

Gopalaswamy UV, Nair CKK. 1992. DNA binding and mutagenecity of lindane and its metabolites. Bull Environ Contam Toxicol 49:300–305.

*Gralla EJ, Fleischman YK, Luthra YK, et al. 1977. Toxic effects of hexachlorobenzene after daily administration to beagle dogs for one year. Toxicol Appl Pharmacol 40:227–239.

Gray LE. 1998. Xenoendocrine disrupters: Laboratory studies on male reproductive effects. Toxicol Lett 102-103:331–335.

Gray LE, Ostby J. 1998. Effects of pesticides and toxic substances on behavioral and morphological reproductive development: Endocrine versus nonendocrine mechanisms. Toxicol Ind Health 14:159–183.

*Gray LE, Kelce WR, Monosson E, et al. 1995. Exposure to TCDD during development permanently alters reproductive function in male Long Evans rats and hamsters: Reduced ejaculated and epididymal sperm numbers and sex accessory gland weights in offspring with normal androgenic status. Toxicol Appl Pharmacol 131:108–118.

Gray LE, Ostby JS, Kelce WR. 1994. Developmental effects of an environmental antiandrogen: The fungicide vinclozolin alters sex differentiation of the male rat. Toxicol Appl Pharmacol 129:46–52.

*Gray LE, Ostby JS, Kelce WR. 1997. A dose-response analysis of the reproductive effects of a single gestational dose of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin in male Long Evans hooded rat offspring. Toxicol Appl Pharmacol 146:11–20.

*Gray LE Jr, Ostby J, Marshall R, et al. 1993. Reproductive and thyroid effects of low-level polychlorinated biphenyl (Aroclor 1254) exposure. Fundam Appl Toxicol 20(3):288–294.

Gray LE, Ostby J, Wolf C, et al. 1998. The value of mechanistic studies in laboratory animals for the prediction of reproductive effects in wildlife: Endocrine effects on mammalian sexual differentiation. Environ Toxicol Chem 17(1):109–118.

Grimalt JO, Sunyer J, Moreno V, et al. 1994. Risk excess of soft-tissue sarcoma and thyroid cancer in a community exposed to airborne organochlorinated compound mixtures with a high hexachlorobenzene content. Int J Cancer 56:200–203.

Grimvall E, Rylander L, Nilsson-Ehle P, et al. 1997. Monitoring of polychlorinated biphenyls in human blood plasma: Methodological developments and influence of age, lactation, and fish consumption. Arch Environ Contam Toxicol 32:329–336.

Guttes S, Failing K, Neumann K, et al. 1998. Chlororganic pesticides and polychlorinated biphenyls in breast tissue of women with benign and malignant beast disease. Arch Environ Contam Toxicol 35:140–147.

*Haake JM, Safe S, Mayura K, et al. 1987. Aroclar 1254 as an antagonist of the teratogenicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin. Toxicol Lett 38:299–306.

Haghdoost NR, Newman LM, Johnson EM. 1997. Multiple chemical exposures: Synergism vs. individual exposure levels. Reprod Toxicol 11(1):9–27.

*Hahn ME, Goldstein JA, Linko P, et al. 1989. Interaction of hexachlorobenzene with the receptor for 2,3,7,8-tetrachlorodibenzo-*p*-dioxin *in vitro* and *in vivo*. Arch Biochem Biophys 270(1):344–355.

*Hany J, Lilienthal H, Sarasin A, et al. 1999. Developmental exposure of rats to a reconstituted PCB mixture or Aroclor 1254: Effects on organ weights, aromatase activity, sex hormone levels, and sweet preference behavior. Toxicol Appl Pharmacol 158:231–243.

*Hansen H, De Rosa CT, Pohl H, et al. 1998. Public health challenges posed by chemical mixtures. Environ Health Perspect Suppl 106(6):1271–1280.

*Hansen LG. 1998. Stepping backward to improve assessment of PCB congener toxicities. Environ Health Perspect Suppl 106(1):171–189.

*Hansen LG, Li MH, Saeed A, et al. 1995. Environmental polychlorinated biphenyls: Acute toxicity of landfill soil extract to female prepubertal rats. Arch Environ Contam Toxicol 29:334–343.

Hardell L, Van Bavel B, Lindstrom G, et al. 1996. Higher concentrations of specific polychlorinated biphenyl congeners in adipose tissue from non-Hodgkin's lymphoma patients compared with controls without a malignant disease. Int J Oncol 9:603–608.

*Harper N, Connor K, Safe S. 1993a. Immunotoxic potencies of polychlorinated biphenyl (PCB), dibenzofuran (PCDF) and dibenzo-*p*-dioxin (PCDD) congeners in C57BL/6 and DBA/2 mice. Toxicology 80:217–227.

*Harper N, Howie L, Conner K, et al. 1993b. Immunosuppressive effects of highly chlorinated biphenyls and diphenyl ethers on T-cell dependent and independent antigens in mice. Toxicology 85:123–135.

Harris M, Zacharewski T, Astroff B, et al. 1989. Characterization of 6-methyl-1,3,8-trichlorobenzofuran (MCDF) as a 2,3,7,8-TCDD antagonist in male rats: Induction of monooxygenases. Chemosphere 19:769–772.

*Hayes WJ, Durham WF, Cueto C, et al. 1956. The effect of known repeated oral doses of chlorophenothane (DDT) in man. J Am Med Assoc 162(9):890–897.

*Hemming H, Bager Y, Flodstrom S, et al. 1995. Liver tumour promoting activity of 3,4,5,3',4'-pentachlorobiphenyl and its interaction with 2,3,7,8-tetrachlorodibenzo-*p*-dioxin. Eur J Pharmacol 292:241–249.

*Hemming H, Flodstrom S, Warngard L, et al. 1993. Relative tumour promoting activity of three polychlorinated biphenyls in rat liver. Eur J Pharmacol 248(2):163–174.

*Hertzler DR. 1990. Neurotoxic behavioral effects of Lake Ontario salmon diets in rats. Neurotoxicol Teratol 12:139–143.

*Hicks HE. 1996. The Great Lakes: A Historical Overview. Toxicol Indus Health 12:303–313.

Hicks HE, De Rosa CT, Cibulas W. 1998. U.S. Agency for Toxic Substances and Disease Registry: Great Lakes Human Health Effects Research Program. J Clean Technol Environ Toxicol Occup Med 7(4):375–396.

*Hirano M, Mitsumori K, Maita k, et al. 1986. Further carcinogenicity study on methylmercury chloride in ICR mice. Jap J Vet Sci 48(1)127–135.

*Hong R, Taylor K, Abonour R. 1989. Immune abnormalities associated with chronic TCDD exposure in rhesus. Chemosphere 18:1785–1797.

*Hood A, Hashmi R, Klaassen CD. 1999. Effects of microsomal enzyme inducers on thyroid-follicular cell proliferation, hyperplasia, and hypertrophy. Toxicol Appl Pharmacol 160:163–170.

*Hori M, Kondo H, Ariyoshi N, et al. 1997. Changes in the hepatic glutathione peroxidase redox system produced by coplanar polychlorinated biphenyls in Ah-responsive and -less-responsive strains of mice: mechanism and implications for toxicity. Environ Toxicol Pharmacol 3:267–275.

*Huisman M, Koopman-Esseboom C, Fidler V, et al. 1995a. Perinatal exposure to polychlorinated biphenyls and dioxins and its effect on neonatal neurological development. Early Hum Dev 41:111–127.

*Huisman M, Koopman-Essesboom C, Lanting CI, et al. 1995b. Neurological condition in 18-month-old children perinatally exposed to polychlorinated biphenyls and dioxins. Early Hum Dev 43:165–176.

*Hultman P, Enestrom S. 1992. Dose-response studies in murine mercury-induced autoimmunity and immune-complex disease. Toxicol Appl Pharmacol 113:199–208.

*Hultman P, Johansson U, Turley SJ, et al. 1994. Adverse immunological effects and autoimmunity induced by dental amalgam and alloy in mice. FASEB J 8:1183–1190.

*IARC. 1997. Polychlorinated dibenzo-*para*-dioxins. Lyons, France: International Agency for Research on Cancer. http://193.51.164.11/htdocs/monographs/vol69/dioxin.html.

*IRIS. 2001. Integrated Risk Information System (IRIS). Cincinnati, OH: National Center for Environmental Assessment. http://www.epa.gov/iris/. February 22, 2001.

*Ilback N-G. 1991. Effects of methyl mercury exposure on spleen and blood natural killer (NK) cell activity in the mouse. Toxicology 67:117–124.

*Ilback –G, Sundberg J, Oskarsson A. 1991. Methyl mercury exposure via placenta and milk impairs natural killer (NK) cell function in newborn rats. Toxicol Lett 58:149–158.

Iverson F, Mehta R, Hierlihy L, et al. 1998. Microsomal enzyme activity, glutathione s-transferaseplacental form expression, cell proliferation, and vitamin A stores in livers of rats consuming Great Lakes salmon. Regul Toxicol Pharmacol 27:S76–S89.

*Jacobson JL, Jacobson SW. 1996. Intellectual impairment in children exposed to polychlorinated biphenyls in utero. N Engl J Med 335(11):783–789.

*Jacobson JL, Jacobson SW, Humphrey HEB. 1990a. Effects of in utero exposure to polychlorinated biphenyls and related contaminants on cognitive functioning in young children. J Pediatr 116(1):38–45.

*Jacobson JL, Jacobson SW, Humphrey HEB. 1990b. Effects of exposure to PCBs and related compounds on growth and activity in children. Neurotoxicol Teratol 12:319–326.

*Jacobson JL, Jacobson SW, Schwartz PM, et al. 1984. Prenatal exposure to an environmental toxin: A test of the multiple effects model. Dev Psychol 20(4):523–532.

*Jacobson SW, Fein GG, Jacobson JL, et al. 1985. The effect of intrauterine PCB exposure on visual recognition memory. Child Dev 56:853–860.

*Jansen HT, Cooke PS, Porcelli J, et al. 1993. Estrogenic and antiestrogenic actions of PCBs in the female rat: In vitro and in vivo studies. Reprod Toxicol 7:237–248.

Johnson BL, Hicks HE, Jones DE, et al. 1998. Public health implications of persistent toxic substances in the Great Lakes and St. Lawrence Basins. J Great Lakes Res 24(2):698–722.

*Jones CJ, Backlin BM, Stoddart RW, et al. 1997. Environmental pollutants as aetiological agents in female reproductive pathology: Placental glycan expression in normal and polychlorinated biphenyl (PCB) - exposed mink (*mustela vison*). Placenta 18(8):689–699.

Jordan SA, Feeley MM. 1999. PCB congener patterns in rats consuming diets containing Great Lakes salmon: Analysis of fish, diets, and adipose tissue. Environ Res A 80:S207–S212.

*Juarez de Ku LM, Sharma-Stokkermans M, Meserve LA. 1994. Thyroxine normalizes polychlorinated biphenyl (PCB) dose-related depression of choline acetyltransferase (ChAT) activity in hippocampus and basal forebrain of 15-day-old rats. Toxicology 94:19–30.

*Kasza L, Collins WT, Capen CC, et al. 1978. Comparative toxicity of polychlorinated biphenyl and polybrominated biphenyl in the rat thyroid gland: Light and electron microscopic alterations after subacute dietary exposure. J Environ Pathol Toxicol 1:587–599.

Kavlock RJ, Perreault SD. 1994. Multiple chemical exposure and risks of adverse reproductive function and outcome. In: Yang RSH, ed. Toxicology of chemical mixtures: Case studies, mechanisms, and novel approaches. San Diego, CA: Academic Press, 245–297.

*Kelce WR, Lambright CR, Gray LE, et al. 1997. Vinclozolin and *p*,*p*'-DDE alter androgen-dependent gene expression: *In vivo* confirmation of an androgen receptor-mediated mechanism. Toxicol Appl Pharmacol 142:192–200.

*Kelce WR, Stone CR, Laws SC, et al. 1995. Persistent DDT metabolite *p*,*p*-DDE is a potent androgen receptor antagonist. Nature 375:581–585.

*Kerkvliet NI. 1994. Immunotoxicology of dioxins and related chemicals. In: Schecter A, ed. Dioxins and health. New York,NY: Plenum Press, 199–225.

*Kihlstrom JE, Olsson M, Jensen SJ, et al. 1992. Effects of PCB and different fractions of PCB on the reproduction of the mink (*mustela vison*). Ambio 21(8):563–569.

*Kimbrough RD. 1997. Studies in children: Polychlorinated biphenyls, dibenzo-p-dioxins and dibenzofurans. Health Environ Dig 10(7):61–63.

*Kociba RJ, Keyes DG, Beyer JE, et al. 1978. Results of a two-year chronic toxicity and oncogenicity study of 2,3,7,8-tetrachlorodibenzo-p-dioxin in rats. Toxicol Appl Pharmacol 46:279–303.

*Kodavanti PRS, Tilson HA. 1997. Structure-activity relationships of potentially neurotoxic PCB congeners in the rat. Neurotoxicology 18(2):425–442.

*Kodavanti PRS, Shin D-S, Tilson HA, et al. 1993. Comparative effects of two polychlorinated biphenyl congeners on calcium homeostasis in rat cerebellar granule cells. Toxicol Appl Pharmacol 123:97–106.

*Kodavanti PRS, Ward TR, McKinney JD, et al. 1995. Increased [³H]phorbol ester binding in rat cerebellar granule cells by polychlorinated biphenyl mixtures and congeners: Structure-activity relationships. Toxicol Appl Pharmacol 130:140–148.

*Kodavanti PRS, Ward TR, McKinney JD, et al. 1996. Increased [³H]phorbol ester binding in rat cerebellar granule cells and inhibition of ⁴⁵Ca²⁺ sequestration in rat cerebellum by polychlorinated diphenyl ether congeners and analogs: Structure-activity relationships. Toxicol Appl Pharmacol 138:251–261.

*Kohn MC, Sewall CH, Lucier GW, et al. 1996. A mechanistic model of effects of dioxin on thyroid hormones in the rat. Toxicol Appl Pharmacol 165:29–48.

*Koopman-Esseboom C, Morse DC, Weisglas-Kuperus N, et al. 1994. Effects of dioxins and polychlorinated biphenyls on thyroid hormone status of pregnant women and their infants. Pediatr Res 36(4):468–473.

*Koopman-Esseboom C, Weisglas-Kuperus N, de Ridder MAJ, et al. 1996. Effects of polychlorinated biphenyl/dioxin exposure and feeding type on infants' mental and psychomotor development. Pediatrics 97(5):700–706.

*Korach KS, Sarver P, Chae K, et al. 1988. Estrogen receptor-binding activity of polychlorinated hydroxybiphenyls: Conformationally restricted structural probes. Mol Pharmacol 33:120–126.

*Kostyniak PJ, Stinson C, Greizerstein HB, et al. 1999. Relation of Lake Ontario fish consumption, lifetime lactation, and parity to breast milk polychlorobiphenyl and pesticide concentrations. Environ Res A80:S166–S174.

*Kramer VJ, Helferich WG, Bergman A, et al. 1997. Hydroxylated polychlorinated biphenyl metabolites are anti-estrogenic in a stably transfected human breast adenocarcinoma (MCF7) cell line. Toxicol Appl Pharmacol 144:363–374.

*Krishnan V, Safe S. 1993. Polychlorinated biphenyls (PCBs), dibenzo-p-dioxins (PCDDs), and dibenzofurans (PCDFs) as antiestrogens in MCF-7 human breast cancer cells: Quantitative structure-activity relationships. Toxicol Appl Pharmacol 120(1):55–61.

*Larsen BR, Turrio-Baldassarri L, Nilsson T, et al. 1994. Toxic PCB congeners and organochlorine pesticides in Italian human milk. Ecotoxicol Environ Saf 28:1–13.

*Laug E, Nelson A, Gitzhugh O, et al. 1950. Liver cell alternation and DDT storage in the fat of the rat induced by dietary levels of 1 to 50 ppm DDT. J Pharmacol Exp Ther 98:268.

*Lecavalier PR, Chu I, Villeneuve D, et al. 1994. Combined effects of mercury and hexachlorobenzene in the rat. J Environ Sci Health B 29(5):951–961.

*Leonzio C, Fossi MC, Casini S. 1996b. Porphyrins as biomarkers of methylmercury and PCB exposure in experimental quail. Bull Environ Contam Toxicol 56:244–250.

*Leonzio C, Monaci F, Fossi MC, et al. 1996a. Multiresponse biomarker evaluation of interactions between methylmercury and Arochlor 1260 in quail. Ecotoxicology 5:365–376.

*Li M-H, Hansen LG. 1996b. Enzyme induction and acute endocrine effects in prepubertal female rats receiving environmental PCB/PCDF/PCDD mixtures. Environ Health Perspect 104(7):712–722.

*Li M-H, Hansen LG. 1996a. Responses of prepubertal female rats to environmental PCBs with high and low dioxin equivalencies. Fundam Appl Toxicol 33:282–293.

*Li M-H, Hansen LG. 1997. Consideration of enzyme and endocrine interactions in the risk assessment of PCBs. Rev Toxicol 1 :71–156.

*Li M-H, Rhine C, Hansen LG. 1998. Hepatic enzyme induction and acute endocrine effects of 2,3,3',4',6'-pentachlorobiphenyl in prepubertal female rats. Arch Environ Contam Toxicol 35:97–103.

*Li SMA, Denomme MA, Leece B, et al. 1989. Hexachlorobenzene: Biochemical effects and synergistic toxic interactions with 2,3,7,8-tetrachlorodibenzo-*p*-dioxin. Toxicol Environ Chem 22:215–227.

*Li X, Rozman KK. 1995. Subchronic effects of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) and their reversibility in male Sprague-Dawley rats. Toxicology 97:133–140.

Liljegren G, Hardell L, Lindstrom G, et al. 1998. Case-control study on breast cancer and adipose tissue concentrations of congener specific polychlorinated biphenyls, DDE and hexachlorobenzene. Eur J Cancer Prev 7:135–140.

*Linder RE, Gaines TB, Kimbrough RD. 1974. The effect of polychlorinated biphenyls on rat reproduction. Food Cosmet Toxicol 12:63–77.

*Linko P, Yeowell HN, Gasiewicz TA, et al. 1986. Induction of cytochrome P-450 isozymes by hexachlorobenzene in rats and aromatic hydrocarbon (Ah)-responsive mice. J Biochem Toxicol 1(2):95–107.

*Loeffler IK, Peterson RE. 1999. Interactive effects of TCDD and *p*,*p*'-DDE on male reproductive tract development in *in utero* and lactationally exposed rats. Toxicol Appl Pharmacol 154:28–39.

*Longcope C. 2000. The male and female reproductive systems in hypothyroidism. In: Braverman LE, Utiger RD, eds. Werner & Ingbar's the thyroid: A fundamental and clinical text. Eighth edition. Philadelphia, PA: Lippincott Williams & Wilkins, 824–827.

Longnecker MP, Rogan WJ, Lucier G. 1997. The human health effects of DDT (dichlorodiphenyltrichloroethane) and PCBs (polychlorinated biphenyls) and an overview of organochlorines in public health. Ann Rev Public Health 18:211–244.

*Lonky E, Reihman J, Darvill T, et al. 1996. Neonatal behavioral assessment scale performance in humans influenced by maternal consumption of environmentally contaminated Lake Ontario fish. J Great Lakes Res 22(2):198–212.

*Lundberg C. 1974. Effect of DDT on Cytochrome P-450 and oestrogen-dependent functions in mice. Environ Physiol Biochem 4:200–204.

*Lundkvist U. 1990. Clinical and reproductive effects of Clophen A50 (PCB) administered during gestation on pregnant guinea pigs and their offspring. Toxicology 6:249–257.

Luster MI, Hong LH, Osborne R, et al. 1986. 1-Amino-3,7,8-trichlorodibenzo-p-dioxin: A specific antagonist for TCDD-induced myelotoxicity. Biochem Biophys Res Commun 139(2):747–756.

MacLuskey NJ, Brown TJ, Schantz S, et al. 1998. Hormonal interactions in the effects of halogenated aromatic hydrocarbons on the developing brain. Toxicol Ind Health 14(1–2):185–208.

*Mariussen E, Anderson JM, Fonnum F. 1999. The effect of polychlorinated biphenyls on the uptake of dopamine and other neurotransmitters into rat brain synaptic vesicles. Toxicol Appl Pharmacol 161:274–282.

*Marty MS, Atchison WD. 1997. Pathways mediating Ca^{2+} entry in rat cerebellar granule cells following *in vitro* exposure to methyl mercury. Toxicol Appl Pharmacol 147:319–330.

*Mayes BA, McConnell EE, Neal BH, et al. 1998. Comparative carcinogenicity in Sprague-Dawley rats of the polychlorinated biphenyl mixtures Aroclors 1016, 1242, 1254, and 1260. Toxicol Sci 41(1):62–76.

*Mendola P, Buck GM, Sever LE, et al. 1997. Consumption of PCB-contaminated freshwater fish and shortened menstrual cycle length. Am J Epidemiol 146(11):955–960.

*Mendola P, Buck GM, Vena JE, et al. 1995. Consumption of PCB-contaminated sportfish and risk of spontaneous fetal death. Environ Health Perspect 103(5):498–502.

Mendoza CE, Shields JB. 1976. Effects of hexachlorobenzene on malathion LD50 and on cholinesterase and carboxylesterase activities in organs of the suckling albino rat. Toxicol Appl Pharmacol 35:447–453.

*Mergler D, Belanger S, Larribe F, et al. 1998. Preliminary evidence of neurotoxicity associated with eating fish from the upper St. Lawrence River Lakes. Neurotoxicology 19(4–5):691–702.

*Mitsumori k, Hirano M, Ueda H, et al. 1990. Chronic toxicity and carcinogenicity of methylmercury chloride in B6C3F1 mice. Fundam Appl Toxicol 14:179–190.

*Mohamed M, Burbacher T, Mottett N. 1987. Effects of methyl mercury on testicular functions in Macaca fascicularis monkeys. Pharmacol Toxicol 60(1):29–36.

*Moore M, Mustain M, Daniel K, et al. 1997. Antiestrogenic activity of hydroxylated polychlorinated biphenyl congeners identified in human serum. Toxicol Appl Pharmacol 142:160–168.

*Morrissey RE, Harris MW, Diliberto JJ, et al. 1992. Limited PCB antagonism of TCDD-induced malformations in mice. Toxicol Lett 60:19–25.

*Morse DC, Wehler EK, Wesseling W, et al. 1996. Alterations in rat brain thyroid hormone status following pre- and postnatal exposure to polychlorinated biphenyls (Aroclor 1254). Toxicol Appl Pharmacol 136:269–279.

Mumtaz M, Colman J. 1992. The risk assessment of chemical mixtures: Fine tuning the hazard index. In: Dodd DE, Clewell HJ, Mattie DR eds. Proceedings of the 1992 conference on toxicology: Applications of advances in toxicology to risk assessment. Wright-Patterson Air Force Base, Ohio: Armstrong Laboratory.

Mumtaz MM, Durkin PR. 1992. A weight-of-evidence approach for assessing interactions in chemical mixtures. Toxicol Ind Health 8: 377–406.

Mumtaz MM, De Rosa CT, Durkin PR. 1994. Approaches and challenges in risk assessments of chemical mixtures. In: Yang RSH, ed. Toxicology of chemical mixtures: Case studies, mechanisms and novel approaches. New York, NY: Academic Press, 565–597.

Mumtaz MM, Poirier KA, Colman JT. 1997. Risk assessment for chemical mixtures: Fine-tuning the hazard index approach. J Clean Technol Environ Toxicol Occup Med 6(2):189–204.

*NCI. 1978. Bioassays of DDT, TDE, and p,p'-DDE for possible carcinogenicity. Bethesda, MD: Carcinogenesis Testing Program, National Cancer Institute. Publication No. 78-1386.

*Nesaretnam K, Darbre P. 1997. 3,5,3',5'-Tetrachlorobiphenyl is a weak oestrogen agonist *in vitro* and *in vivo*. J Steroid Biochem Mol Biol 62(5/6):409–418.

*Nesaretnam K, Corcoran D, Dils RR, et al. 1996. 3,4,3',4'-Tetrachlorobiphenyl acts as an estrogen *in vitro* and *in vivo*. Mol Endocrinol 923–936.

*Ness DK, Schantz SL, Mostagian J, et al. 1993. Perinatal exposure to two polychlorinated biphenyl congeners: Histological effects on the thyroid gland. Toxicologist 13(1):358.

*NTP. 1982. Carcinogenesis bioassay of 2,3,7,8-tetrachlorodibenzo-p-dioxin (CAS no. 1746-01-6) in Osborne-Mendel rats and B6C3F1 mice (gavage study). Bethesda, MD: Carcinogenesis Testing Program, National Cancer Institute, National Institute of Health. Research Triangle Park, NC: National Toxicology Program. DHHS publication no. 82-1765.

*Okey AB, Riddick DS, Harper PA. 1994. The Ah receptor: Mediator of the toxicity of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD) and related compounds. Toxicol Lett 70:1–22.

*Omara FO, Brochu C, Flipo D, et al. 1997. Immunotoxicity of environmentally relevant mixtures of polychlorinated aromatic hydrocarbons with methyl mercury on rat lymphocytes in vitro. Environ Toxicol Chem 16(3):576–581.

*Omara FO, Flipo D, Brochu C, et al. 1998. Lack of suppressive effects of mixtures containing low levels of methylmercury (MeHg), polychlorinated dibenzo-*p*-dioxins (PCDDS), polychlorinated dibenzofurans (PCDFS), and aroclor biphenyls (PCBS) on mixed lymphocyte reaction, phagocytic, and natural killer cell activities of rat leukocytes in vitro. J Toxicol Environ Health A54:561–577.

*Pappas BA, Murtha SJE, Park GAS, et al. 1998. Neurobehavioral effects of chronic ingestion of Great Lakes chinook salmon. Regul Toxicol Pharmacol 27:S55–S67.

*Parham FM, Portier CJ. 1998. Using structural information to create physiologically based pharmacokinetic models for all polychlorinated biphenyls. Toxicol Appl Pharmacol 151:110–116.

Patandin S, Koopman-Esseboom C, De Ridder MAJ, et al. 1998. Effects of environmental exposure to Polychlorinated biphenyls and dioxins on birth size and growth in Dutch children. Pediatr Res 44(4):538–545.

*Patandin S, Lanting CI, Mulder PGH, et al. 1999. Effects of environmental exposure to polychlorinated biphenyls and dioxins on cognitive abilities in Dutch children at 42 months of age. J Pediatr 134(1):33–41.

*Patnode KA, Curtis LR. 1994. 2,2',4,4',5,5'- and 3,3',4,4',5,5'-Hexachlorobiphenyl alteration of uterine progesterone and estrogen receptors coincides with embryotoxicity in mink (*mustela vison*). Fundam Appl Toxicol 127:9–18.

*Poland A, Knutson JC. 1982. 2,3,7,8-Tetrachlorodibenzo-p-dioxin and related halogenated aromatic hydrocarbons. Examinations of the mechanisms of toxicity. Ann Rev Pharmacol Toxicol 22:517–554.

*Porterfield SP, Hendry LB. 1998. Impact of PCBs on thyroid hormone directed brain development. Toxicol Ind Health 14:103–120.

*Preston BD, Miller EC, Miller JA. 1985. The activities of 2,2',5,5'-tetrachlorobiphenyl, its 3,4-oxide metabolite, and 2,2',4,4'-tetrachlorobiphenyl in tumor induction and promotion assays. Carcinogenesis 6(3):451–453.

*Price SC, Ozalp S, Weaver R, et al. 1988. Thyroid hyperactivity caused by hypolipodaemic compounds and polychlorinated biphenyls: The effect of coadministration in the liver and thyroid. Arch Toxicol Suppl 12:85–92.

*Provost TL, Juarez De Ku LM, Zender C, et al. 1999. Dose- and age-dependent alterations in choline acetyltransferase (ChAT) activity, learning and memory, and thyroid hormones in 15- and 30-day old rats exposed to 1.25 or 12.5 ppm polychlorinated biphenyl (PCB) beginning at conception. Prog Neuro-Psychopharmacol Biol Psychiat 23:915–928.

*Renner G. 1980. Toxicities of combinations of pentachloronitrobenzene with mercuric chloride or cadmium chloride, and hexachlorobenzene with mercuric chloride administered to rats. Xenobiotica 10(7–8):551–556.

*Restum JC, Bursian SJ, Giesy JP, et al. 1998. Multigenerational study of the effects of consumption of PCB-contaminated carp from Saginaw Bay, Lake Huron, on mink. 1. Effects on mink reproduction, kit growth and survival, and selected biological parameters. J Toxicol Environ Health, A 54:343–375.

*Rice DC. 1997. Effect of postnatal exposure to a PCB mixture in monkeys on multiple fixed intervalfixed ratio performance. Neurotoxicol Teratol 19(6):429–434.

*Rice DC. 1998. Effects of postnatal exposure of monkeys to a PCB mixture on spatial discrimination reversal and DRL performance. Neurotoxicol Teratol 20(4):391–400.

*Rice DC. 1999. Behavioral impairment produced by low-level postnatal PCB exposure in monkeys. Environ Res 80:S113–S121.

*Rice DC, Hayward S. 1997. Effects of postnatal exposure to a PCB mixture in monkeys on nonspatial discrimination reversal and delayed alternation performance. Neurotoxicology 18(2):479–494.

*Rice DC, Hayward S. 1999. Effects of exposure to 3,3',4,4',5-pentachlorobiphenyl (PCB 126) throughout gestation and lactation on behavior (concurrent random interval-random interval and progressive ratio performance) in rats. Neurotoxicol Teratol 21(6):679–687.

Riddick DS, Huang Y, Harper PA, et al. 1994. 2,3,7,8-tetrachlorodibenzo-*p*-dioxin versus 3-methylcholanthrene: Comparative studies of Ah receptor binding, transformation, and induction of CYP1A1. J Biol Chem 269(16):12118–12128.

*Rier SE, Martin DC, Bowman RE, et al. 1993. Endometriosis in Rhesus monkeys (*Macata mulatta*) following chronic exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin. Fundam Appl Toxicol 21:433–441.

*Risebrough RW, Anderson DW. 1975. Some effects of DDE and PCB on mallards and their eggs. J Wildl Manage 39(3):508–513.

*Robertson LW, Gupta RC. 2000. Metabolism of polychlorinated biphenyls (PCBs) generates electrophiles and reactive oxygen species that damage DNA. In: Williams GM, Aruoma OI, eds. Molecular drug metabolism and toxicology. OICA International, 1–19.

*Rogan WJ, Gladen BC, McKinney JD, et al. 1986b. Neonatal effects of transplacental exposure to PCBs and DDE. J Pediatr 109:335–341.

*Rogan WJ, Gladen BC, McKinney JD, et al. 1986a. Polychlorinated biphenyls (PCBs) and dichlorophenyl dichloroethene (DDE) in human milk: Effects of maternal factors and previous lactation. Am J Public Health 76:172–177.

*Rogan WJ, Gladen BC, McKinney JD, et al. 1987. Polychlorinated biphenyls (PCBs) and dichlorodiphenyl dichloroethene (DDE) in human milk: Effects on growth, morbidity, and duration of lactation. Am J Public Health 77:1294–1297.

*Roman BL, Pollenz RS, Peterson RE. 1998a. Responsiveness of the adult male rat reproductive tract to 2,3,7,8-tetrachlorodibenzo-*p*-dioxin exposure: Ah receptor and ARNT expression, CY1A1 induction, and Ah receptor down-regulation. Toxicol Appl Pharmacol 150:228–239.

*Roman BL, Sommer RJ, Shinomiya K, et al. 1995. *In utero* and lactational exposure of the male rat to 2,3,7,8-tetrachlorodibenzo-*p*-dioxin: Impaired prostrate growth and development without inhibited androgen production. Toxicol Appl Pharmacol 134:241–250.

*Roman BL, Timms BG, Prins GS, et al. 1998b. *In utero* and lactational exposure of the male rat to 2,3,7,8-tetrachlorodibenzo-*p*-dioxin impairs prostrate development: 2. Effects on growth and cytodifferentiation. Toxicol Appl Pharmacol 150:254–270.

*Ross PS, De Swart RL, Reijnders PJH, et al. 1995. Contaminant-related suppression of delayed-type hypersensitivity and antibody responses in harbor seals fed herring from the Baltic Sea. Environ Health Perspect 103(2):162–167.

Rozman K, Mueller W, Coulston F, et al. 1977. Long-term feeding study of hexachlorobenzene in rhesus monkeys: (Preliminary communication). Chemosphere 2/3:81–84.

Rozman K, Rozman T, Greim H. 1981. Enhanced fecal elimination of stored hexachlorobenzene from rats and rhesus monkeys by hexadecane or mineral oil. Toxicology 22:33–44.

*Rylander L, Hagmar L. 1995. Mortality and cancer incidence among women with a high consumption of fatty fish contaminated with persistent organochlorine compounds. Scand J Work Environ Health 21:419–426.

*Rylander L, Hagmar L. 1999. Anthropometric and psychometric examinations of conscripts born to mothers with a high intake of fish contaminated with persistent organochlorines. In: Dioxin 99: International Symposium on halogenated environmental organic pollutants and POPs. Organohalogen Compounds 44:413–416.

*Rylander L, Stromberg U, Dyremark E, et al. 1998a. Polychlorinated biphenyls in blood plasma among Swedish female fish consumers in relation to low birth weight. Am J Epidemiol 147(5):493–502.

*Rylander L, Stromberg U, Hagmar L. 1995. Decreased birthweight among infants born to women with a high dietary intake of fish contaminated with persistent organochlorine compounds. Scand J Work Environ Health 21:368–375.

*Rylander L, Stromberg U, Hagmar L. 1996. Dietary intake of fish contaminated with persistent organochlorine compounds in relation to low birthweight. Scand J Work Environ Health 22:260–266.

*Rylander L, Stromberg U, Hagmar L. 1998b. Lowered birthweight among infants born to women with high intake of fish contaminated with persistent organochlorine compounds. In: Organohalogen Compounds 38:275–277.

*Safe S. 1984. Polychlorinated biphenyls (PCBs) and polybrominated biphenyls (PBBs): Biochemistry, toxicology, and mechanism of action. CRC Crit Rev Toxicol 13:319–395.

*Safe S. 1990. Polychlorinated biphenyls (PCBs), dibenzo-p-dioxins (PCDDs), dibenzofurans (PCDFs), and related compounds: Environmental and mechanistic considerations which support the development of toxic equivalency factors (TEFs). Crit Rev Toxicol 21:51–88.

Safe SH. 1994. Polychlorinated biphenyls (PCBs): Environmental impact, biochemical and toxic responses, and implications for risk assessment. Crit Rev Toxicol 24(2):87–149.

*Safe S. 1998a. Limitations of the toxic equivalency factor approach for the risk assessment of TCDD and related compounds. Teratogenesis Carcinog Mutagen 17:285–304.

*Safe S. 1998b. Development validation and problems with the toxic equivalency factor approach for risk assessment of dioxins and related compounds. J Anim Sci 76:134–141.

*Safe SH. 1999. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) and related environmental antiestrogens: Characterization and mechanism of action. In: Naz RK, ed. Endocrine disruptors: Effects on male and female reproductive systems. Boca Raton, FL: CRC Press, 187–221.

Safe SH, Gaido K. 1998. Phytoestrogens and anthropogenic estrogenic compounds. Environ Toxicol Chem 17(1):119–126.

*Sager DB, Girard DM. 1994. Long-term effects on reproductive parameters in female rats after translactational exposure to PCBs. Environ Res 66(1):52–76.

*Sager D, Girard D, Nelson D. 1991. Early postnatal exposure to PCBs: Sperm function in rats. Environ Toxicol Chem 10:737–746.

*Sager DB, Shih-Schroeder W, Girard D. 1987. Effect of early postnatal exposure to polychlorinated biphenyls (PCBs) on fertility in male rats. Bull Environ Contam Toxicol 38:946–953.

*Schantz SL, Gardiner JC, Gasior DM, et al. 1999. Motor function in aging Great Lakes fisheaters. Environ Res 80:546–556.

*Schantz SL, Seo B-W, Wong PW, et al. 1997. Long-term effects of developmental exposure to 2,2',3,5',6-pentachlorobiphenyl (PCB 95) on locomotor activity, spatial learning and memory and brain ryanodine binding. Neurotoxicology 18(2):457–468.

*Schantz SL, Sweeney AM, Gardiner JC, et al. 1996. Neuropsychological assessment of an aging population of Great Lakes fisheaters. Toxicol Ind Health 12(3–4):403–417.

Schecter A, Ryan JJ, Papke O. 1998. Decrease in levels and body burden of dioxins, dibenzofurans, PCBs, DDE, and HCB in blood and milk in a mother nursing twins over a thirty-eight month period. Chemosphere 37(9–12):1807–1816.

Schielen P, Den Besten C, Vos JG, et al. 1995. Immune effects of hexachlorobenzene in the rat: Role of metabolism in a 13-week feeding study. Toxicol Appl Pharmacol 131:37–43.

*Schuur AG, Bergman A, Brouwer A, et al. 1999. Effects of pentachlorophenol and hydroxylated polychlorinated biphenyls on thyroid hormone conjugation in a rat and a human hepatoma cell line. Toxicol in Vitro 13:417–425.

*Schuur AG, Cenjin PH, van Toor H, et al. 1998a. Effect of Aroclor 1254 on thyroid hormone sulfation in fetal rats. In: Johansson N, Bergman A, Broman D, et al., ed. Organohalogen compounds. Akademitryck, Edsbruk:, Vol. 37, 249–252.

*Schuur AG, van Leeuwen-Bol I, Jong WMC, et al. 1998b. *In vitro* inhibition of thyroid hormone sulfation by polychloribiphenylols: Isozyme specificity and inhibition kinetics. Toxicol Sci 45:188–194.

Schwab BW. 1999. The TEF approach for hexachlorobenzene. Environ Health Perspect 107(4):A183–A184.

Seegal RF. 1996a. Can epidemiological studies discern subtle neurological effects due to perinatal exposure to PCBs? Neurotoxicol Teratol 18(3):251–254.

*Seegal RF. 1996b. Epidemiological and laboratory evidence of PCB-induced neurotoxicity. Crit Rev Toxicol 26(6):709–737.

*Seegal RF. 1999. Are PCBs the major neurotoxicant in Great Lakes salmon? Environ Res 80:S38–S45.

*Seegal RF, Brosch K, Bush, B et al. 1989. Effects of Aroclor 1254 on dopamine and norepinephrine concentrations in pheochromocytoma (PC-12) cells. Neurotoxicology 10:757–764.

*Seegal RF, Brosch KO, Okoniewski RJ. 1997. Effects of *in utero* and lactational exposure of the laboratory rat to 2,4,2',4'- and 3,4,3',4'-tetrachlorobiphenyl on dopamine function. Toxicol Appl Pharmacol 146:95–103.

*Seegal RF, Bush B, Shain W. 1990. Lightly chlorinated *ortho*-substituted PCB congeners decrease dopamine in nonhuman primate brain and in tissue culture. Toxicol Appl Pharmacol 106:136–144.

*Seegal RF, Pappas BA, Park GAS. 1998. Neurochemical effects of consumption of Great Lakes salmon by rats. Regul Toxicol Pharmacol 27:S68–S75.

*Seiler P, Fischer B, Lindenau A, et al. 1994. Effects of persistent chlorinated hydrocarbons on fertility and embryonic development in the rabbit. Human Reprod 9:1920–1926.

*Seo BW, Meserve LA. 1995. Effects of maternal ingestion of Aroclor 1254 (PCB) on the developmental pattern of oxygen consumption and body temperature in neonatal rats. Bull Environ Contam Toxicol 55:22–28.

*Shain W, Bush B, Seegal R. 1991. Neurotoxicity of polychlorinated biphenyls: Structure-activity relationship of individual congeners. Toxicol Appl Pharmacol 111:33–42.

*Shenker BJ, Berthold P, Rooney C, et al. 1993. Immunotoxic effects of mercuric compounds on human lymphocytes and monocytes. III. Alterations in B-cell function and viability. Immunopharmacol Immunotoxicol 15(1):87–112.

*Silberhorn EM, Glauert HP, Robertson LW. 1990. Carcinogenicity of polyhalogenated biphenyls: PCBs and PBBs. Crit Rev Toxicol 20:440–496.

*Silkworth JB, Grabstein EM. 1982. Polychlorinated biphenyl immunotoxicity: Dependence on isomer planarity and the Ah gene complex. Toxicol Appl Pharmacol 65:109–115.

Sinclair PR, Gorman N, Sinclair JF, et al. 1995. Ascorbic acid inhibits chemically induced uroporphyria in ascorbate-requiring rats. Hepatology 22(2):565–572.

*Sinclair PR, Walton HS, Gorman N, et al. 1997. Multiple roles of polyhalogenated biphenyls in causing increases in cytochrome P450 and uroporphyrin accumulation in cultured hepatocytes. Toxicol Appl Pharmacol 147:171–179.

*Smialowicz RJ, DeVito MJ, Riddle MM, et al. 1997. Opposite effects of 2,2',4,4',5,5'-hexachlorobiphenyl and 2,3,7,8-tetrachlorodibenzo-p-dioxin on the antibody response to sheep erythrocytes in mice. Fundam Appl Toxicol 37:141–149.

Smith AG, Carthew P, Francis JE, et al. 1993. Enhancement by iron of hepatic neoplasia in rats caused by hexachlorobenzene. Carcinogenesis 14(7):1381–1387.

*Smith AG, Dinsdale D, Cabral JRP, et al. 1987. Goitre and wasting induced in hamsters by hexachlorobenzene. Arch Toxicol 60:343–349.

*Smith AG, Francis JE, Carthew P. 1990a. Iron as a synergist for hepatocellular carcinoma induced by polychlorinated biphenyls in *Ah*-responsive C57BL/10ScSn mice. Carcinogenesis 11(3):437–444.

*Smith AG, Francis JE, Green JA, et al. 1990b. Sex-linked hepatic uroporphyria and the induction of cytochromes P450IA in rats caused by hexachlorobenzene and polyhalogenated biphenyls. Biochem Pharmacol 40(9):2059–2068.

Sonnenschein C, Soto AM. 1998. An updated review of environmental estrogen and androgen mimics and antagonists. J Steroid Biochem Mol Biol 65:143–150.

*Stack AS, Altman-Hamamdzic S, Morris PJ, et al. 1999. Polychlorinated biphenyl mixtures (Aroclors) inhibit LPS-induced murine splenocyte proliferation in vitro. Toxicology 139:137–154.

*Stahl BU, Kettup A, Rozman K. 1992. Comparative toxicity of four chlorinated dibenzo-*p*-dioxins (CDDs) and their mixture: Part I: Acute toxicity and toxic equivalency factors (TEFs). Arch Toxicol 66:471–477.

*Stewart P, Darvill T, Lonky E, et al. 1999. Assessment of prenatal exposure to PCBs from maternal consumption of Great Lakes fish: An analysis of PCB patter and concentration. Environ Res 80:S87–S96.

*Stewart P, Pagano J, Sargent D, et al. 2000a. Effects of Great Lakes fish consumption on brain PCB pattern, concentration, and progressive-ratio performance. Environ Res 82:18–32.

*Stewart P, Reihman J, Lonky E, et al. 2000b. Prenatal PCB exposure and neonatal behavioral assessment scale (NBAS) performance. Neurotoxicol Teratol 22:21–29.

*Sufit RL, Hodach R, Arends R, et al. 1986. Decreased conduction velocity and pseudomyotonia in hexachlorobenzene-fed rats. In: Hexachlorobenzene: Proceedings of an international symposium. IARC Sci Publ 77:361–362..

*Svensson B-G, Hallberg T, Nilsson A, et al. 1994. Parameters of immunological competence in subjects with high consumption of fish contaminated with persistent organochlorine compounds. Int Arch Occup Environ Health 65:351–358.

Svensson B-G, Nilsson A, Hansson M, et al. 1991. Exposure to dioxins and dibenzofurans through the consumption of fish. N Engl J Med 324:8–12.

*Svensson B-G, Nilsson A, Jonsson E, et al. 1995. Fish consumption and exposure to persistent organochlorine compounds, mercury, selenium and methylamines among Swedish fishermen. Scand J Work Environ Health 21:96–105.

Swain WR. 1991. Effects of organochlorine chemicals on the reproductive outcome of humans who consumed contaminated Great Lakes fish: An epidemiologic consideration. J Toxicol Environ Health 33:587–639.

*Swanson GM, Ratcliffe HE, Fischer LJ. 1995. Human exposure to polychlorinated biphenyls (PCBs): A critical assessment of the evidence for adverse health effects. Regul Toxicol Pharmacol 21:136–150.

*Takabatake E, Fujita M, Sawa Y. 1980. Combined effects of polychlorinated biphenyls and methylmercury on hepatic microsomal monooxygenases and the hepatotoxic action of bromobenzene. J Pharmacobiodyn 3:463–469.

*Tanimura T, Ema M, Kihara T. 1980. Effects of combined treatment with methylmercury and polychlorinated biphenyls (PCBs) on the development of mouse offspring. In: Persuad TVN, ed. Neural and behavioral teratology. Baltimore, MD: University Park Press, 163–198.

*Tilson HA, Kodavanti PRS. 1997. Neurochemical effects of polychlorinated biphenyls: An overview and identification of research needs. Neurotoxicology 18(3):727–744.

*Tilson HA, Kodavanti PRS. 1998. The neurotoxicity of polychlorinated biphenyls. Neurotoxicology 19(4–5):517–526.

*Tilson HA, Kodavanti PRS, Mundy WR, et al. 1998. Neurotoxicity of environmental chemicals and their mechanism of action. Toxicol Lett 102–103:631–635.

*Tithof PK, Contreras ML, Gamey PE. 1995. Aroclor 1242 stimulates the production of inositol phosphates in polymorphonuclear neutrophils. Toxicol Appl Pharmacol 131:136–143.

Toraason M. 1999. 8-Hydroxydeoxyguanosine as a biomarker of workplace exposures. Biomarkers 4(1):3–26.

Tryphonas H. 1995. Immunotoxicity of PCBs (Aroclors) in relation to Great Lakes. Environ Health Perspect Suppl 103(9):35–46.

*Tryphonas L, Arnold DL, Zawidzka Z, et al. 1986a. A pilot study in adult Rhesus monkeys (*m.mullata*) treated with Aroclor 1254 for two years. Toxicol Pathol 14:1–10.

*Tryphonas L, Charbonneau S, Tryphonas H, et al. 1986b. Comparative aspects of Aroclor 1254 toxicity in adult Cynomolgus and Rhesus monkeys: A pilot study. Arch Environ Contam Toxicol 15:159–169.

Tryphonas H, Fournier M, Lacroix F, et al. 1998a. Effects of Great Lakes fish consumption on the immune system of Sprague-Dawley rats investigated during a two-generation reproductive study: II. Quantitative and functional aspects. Regul Toxicol Pharmacol 27:S40–S54.

*Tryphonas H, Hayward S, O'Grady L, et al. 1989. Immunotoxicity studies of PCB (Aroclor 1254) in the adult Rhesus (macaca mulatta) monkey-preliminary report. Int J Immunopharmacol 11(2):199–206.

Tryphonas H, Luster MI, Schiffman G, et al. 1991. Effect of chronic exposure of PCB (Aroclor 1254) on specific and nonspecific immune parameters in the Rhesus (*macaca mulatta*) monkey. Fundam Appl Toxicol 16:773–786.

*Tryphonas H, McGuire P, Fernie S, et al. 1998b. Effects of Great Lakes fish consumption on the immune system of Sprague-Dawley rats investigated during a two-generation reproductive study: I. Body and organ weights, food consumption, and hematological parameters. Regul Toxicol Pharmacol 27:S28–S39.

Unkila M, Pohjanvirta R, MacDonald E, et al. 1994. Dose response and time course of alterations in tryptophan metabolism by 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD) in the most TCDD-susceptible and the most TCDD-resistant rat strain: Relationship with TCDD lethality. Toxicol Appl Pharmacol 128:280–292.

*van Birgelen APJM. 1998. Hexachlorobenzene as a possible major contributer to the dioxin activity of human milk. Environ Health Perspect 106(11):683–688.

*van Birgelen APJM, Fase KM, van der Kolk J, et al. 1996. Synergistic effect of 2,2',4,4',5,5'-hexachlorobiphenyl and 2,3,7,8-tetrachlorodibenzo-*p*-dioxin on hepatic porphyrin levels in the rat. Environ Health Perspect 104(5):550–557.

*Van Birgelen APJM, Van Der Kolk J, Fase KM, et al. 1994a. Toxic potency of 2,3,3',4,4',5-hexachlorobiphenyl relative to and in combination with 2,3,7,8-tetrachlorodibenzo-*p*-dioxin in a subchronic feeding study in the rat. Toxicol Appl Pharmacol 126:202–213.

*Van Birgelen APJM, Van Der Kolk J, Fase KM, et al. 1994b. Toxic potency of 3,3',4,4',5-pentachlorobiphenyl relative to and in combination with 2,3,7,8-tetrachlorodibenzo-*p*-dioxin in a subchronic feeding study in the rat. Toxicol Appl Pharmacol 127:209–221.

*Van Birgelen APJM, Van der Kolk J, Fase KM, et al. 1995. Subchronic dose-response study of 3,4,7,8-tetrachlorodibenzo-p-dioxin in female Sprague-Dawley rats. Toxicol Appl Pharmacol 132:1–13.

*van Birgelen APJM, van der Kolk J, Poiger H, et al. 1992. Interactive effects of 2,2',4,4',5,5'-hexachlorobiphenyl and 2,3,7,8-tetrachlorodibenzo-p-dioxin on thyroid hormone, Vitamin A, and Vitamin K metabolism in the rat. Chemosphere 25:7–10.

*Van den Berg M, Birnbaum L, Bosveld ATC, et al. 1998. Toxic equivalency factors (TEFs) for PCBs, PCDDs, PCDFs for humans and wildlife. Environ Health Perspect 106(12):775–792.

*Van den Berg M, De Jongh J, Poiger H, et al. 1994. The toxicokinetics and metabolism of polychlorinated dibenzo-*p*-dioxins (PCDDs) and dibenzofurans (PCDFs) and their relevance for toxicity. Crit Rev Toxicol 24(1):1–74.

Van den Berg M, de Vroom E, Olie K, et al. 1986a. Bioavailability of PCDDs and PCDFs on fly ash after semi-chronic oral ingestion by guinea pig and Syrian golden hamster. Chemosphere 15(4):519–533.

Van den Berg M, Van Greevenbroek M, Olie X, et al. 1986b. Bio-availability of PCDDs and PCDFs on fly ash after semi-chronic oral ingestion by the rat. Chemosphere 13(4):509–513.

*van der Burght ASAM, Clijsters PJ, Horbach GJ, et al. 1999. Structure-dependent induction of CYP1A by polychlorinated biphenyls in hepatocytes of Cynomolgus monkeys (*Macaca fascicularis*). Toxicol Appl Pharmacol 155:13–23.

*van der Kolk J, van Birgelen APJM, Poiger H, et al. 1992. Interactions of 2,2',4,4',5,5'-hexachlorobiphenyl and 2,3,7,8-tetrachlorodibenzo-p-dioxin in a subchronic feeding study in the rat. Chemosphere 25(12):2023–2027.

*van der Plas SA, Haag-Gronlund M, Scheu G, et al. 1999. Induction of altered hepatic foci by a mixture of dioxin-like compounds with and without 2,2',4,4',5,5'-hexachlorobiphenyl in female Sprague-Dawley rats. Toxicol Appl Pharmacol 156:30–39.

*VanRaaij JAGM, Frijters CMG, Van Den Berg KJ. 1993. Hexachlorobenzene-induced hypothyroidism: Involvement of different mechanisms by parent compound and metabolite. Biochem Pharmacol 46(8):1385–1391.

*Vena JE, Buck GM, Kostyniak P, et al. 1996. The New York angler cohort study: Exposure characterization and reproductive and developmental health. Toxicol Ind Health 12:327–334.

*Verschuuren HG, Kroes R, Den Tonkelaar EM, et al. 1976. Toxicity of methylmercury chloride in rats. III. Long-term toxicity study. Toxicology 6:107–123.

Villeneuve DC, Valli VE, Norstrom RJ, et al. 1981. Toxicological response of rats fed Lake Ontario or Pacific coho salmon for 28 days. J Environ Sci Health B 16(6):649–689.

*Viluksela M, Stahl BU, Birnbaum LS, et al. 1998a. Subchronic/chronic toxicity of a mixture of four chlorinated dibenzo-*p*-dioxins in rats. I. Design, general observations, hematology, and liver concentrations. Toxicol Appl Pharmacol 151:57–69.

*Viluksela M, Stahl BU, Birnbaum LS, et al. 1998b. Subchronic/chronic toxicity of a mixture of four chlorinated dibenzo-*p*-dioxins in rats. II. Biochemical effects. Toxicol Appl Pharmacol 151:70–78.

*Vonier PM, Crain DA, McLachlan JA, et al. 1996. Interaction of environmental chemicals with the estrogen and progesterone receptors from the oviduct of the American alligator. Environ Health Perspect 104(12):1318–1322.

*Vos JG. 1986. Immunotoxicity of hexachlorobenzene. In: Morris CR, Cabral JRP, eds. Hexachlorobenzene: Proceedings of an international symposium. Lyon,France: International Agency for Research on Cancer, 347–356.

Vos RME, Snoek MC, Van Berkel WJH, et al. 1988. Differential induction of rat hepatic glutathione *s*-transferase isoenzymes by hexachlorobenzene and benzyl isothiocyanate: Comparison with induction by phenobarbital and 3-methylcholanthrene. Biochem Pharmacol 37(6):1077–1082.

Weber H, Harris MW, Haseman JK, et al. 1985. Teratogenic potency of TCDD and TCDD-TCDF combinations in C57BL/6N mice. Toxicol Lett 26:159–167.

Weber LWD, Lebofsky M, Stahl BU, et al. 1992a. Comparative toxicity of four chlorinated dibenzop-dioxins (CDDs) and their mixture: Part II: Structure-activity relationships with inhibition of hepatic phosphoenolpyruvate carboxykinase, pyruvate carboxylase, and γ -glutamyl transpeptidase activities. Arch Toxicol 66:478–483.

Weber LWD, Lebofsky M, Stahl BU, et al. 1992b. Comparative toxicity of four chlorinated dibenzo*p*-dioxins (CDDs) and their mixture: Part III: Structure-activity relationship with increased plasma tryptophan levels, but no relationship to hepatic ethoxyresorufin *o*-deethylase activity. Arch Toxicol 66:484–488.

*Weber LWD, Lebofsky M, Stahl BU, et al. 1995. Correlation between toxicity and effects on intermediary metabolism in 2,3,7,8-tetrachlorodibenzo-p-dioxin-treated male C257BL/6J and DBA/2J mice. Toxicol Appl Pharmacol 131:155–162.

*Welsch F. 1985. Effects of acute or chronic polychlorinated biphenyl ingestion on maternal metabolic homeostasis and on the manifestations of embryotoxicity caused by cyclophosphamide in mice. Arch Toxicol 57:104–113.

Wildhaber ML, Schmitt CJ. 1996. Estimating aquatic toxicity as determined through laboratory tests of Great Lakes sediments containing complex mixtures of environmental contaminants. Environ Monit Assess 41:255–289.

*Wolfle D. 1998. Interactions between 2,3,7,8-TCDD and PCBs as tumor promoters: Limitations of TEFs. Teratogen Carcinog Mutagen 17:217–224.

*Wong PW, Pessah IN. 1996. *Ortho*-substituted polychlorinated biphenyls alter calcium regulation by a ryanodine receptor-mediated mechanism: Structural specificity toward skeletal- and cardiac-type microsomal calcium release channels. Mol Pharmacol 49:740–751.

*Wong PW, Pessah IN. 1997. Noncoplanar PCB 95 alters microsomal calcium transport by an immunophilin FKBP 12-dependent mechanism. Mol Pharmacol 51:693–702.

*Wong PW, Joy RM, Albertson TE, et al. 1997. *Ortho*-substituted 2,2',3,5',6-pentachlorobiphenyl (PCB 95) alters rat hippocampal ryanodine receptors and neuroplasticity *in vitro*: Evidence for altered hippocampal function. Neurotoxicology 18(2):443–456.

*Wren CD, Hunter DB, Leatherland JF, et al. 1987a. The effects of polychlorinated biphenyls and methylmercury, singly and in combination, on mink. I: Uptake and toxic responses. Arch Environ Contam Toxicol 16:441–447.

*Wren CD, Hunter DB, Leatherland JF, et al. 1987b. The effects of polychlorinated biphenyls and methylmercury, singly and in combination on mink. II: Reproduction and kit development. Arch Environ Contam Toxicol 16:449–454.

*Zoeller RT, Dowling ALS, Vas AA. 2000. Developmental exposure to polychlorinated biphenyls exerts thyroid hormone-like effects on the expression of RC3/neurogranin and myelin basic protein messenger ribonucleic acids in the developing rat brain. Endocrinology 141(1):181–189.