

CHAPTER 8. REFERENCES

- *ACGIH. 1986. Documentation of the threshold limit values and biological exposure indices. 5th ed. American Conference of Governmental Industrial Hygienists, Inc., Cincinnati, OH.
- *ACGIH. 2001. Bis(chloromethyl)ether. TLVs and BEIs based on the documentation of the threshold limit values for chemical substances and physical agents and biological exposure indices. Cincinnati, OH: American Conference of Governmental Industrial Hygienists.
- *ACGIH. 2016. Bis(chloromethyl)ether. TLVs and BEIs based on the documentation of the threshold limit values for chemical substances and physical agents and biological exposure indices. Cincinnati, OH: American Conference of Governmental Industrial Hygienists, May 11, 2017.
- *AIHA. 2015. Current ERPG Values (2015). Fairfax, VA: American Industrial Hygiene Association. <https://www.aiha.org/get-involved/AIHAGuidelineFoundation/EmergencyResponsePlanningGuidelines/Documents/2015%20ERPG%20Levels.pdf>. March 22, 2016.
- *Albert RE, Sellakumar AR, Laskin S, et al. 1982. Gaseous formaldehyde and hydrogen chloride induction of nasal cancer in the rat. *J Nat Cancer Inst* 68:597-603.
- *Albert RE, Pasternack BS, Shore RE, et al. 1975. Mortality patterns among workers exposed to chloromethyl ethers- a preliminary report. *Environ Health Perspect* 2:109-214.
- *Anderson D, Styles JA. 1978. The bacterial mutation test. *Br J Cancer* 37:924-930.
- *ASTM. 1987. Bis(chloromethyl) ether (bis CME) in workplace atmospheres (gas chromatography - mass spectrometry - Method D 347679). 1987 Annual book of ASTM standards, Vol. 11.03, Atmospheric analysis; occupational health and safety. American Society for Testing and Materials, Philadelphia, PA: 259-264.
- *ATSDR. 1989. Decision guide for identifying substance-specific data needs related to toxicological profiles; Notice. Agency for Toxic Substances and Disease Registry. *Fed Regist* 54(174):37618-37634.
- *ATSDR. 2017. Bis(2-Chloroethyl)Ether. Full SPL data. Substance priority list (SPL) resource page. Agency for Toxic Substances and Disease Registry, Centers for Disease Control and Prevention. <http://www.atsdr.cdc.gov/SPL/resources/index.html>. October 6, 2017.
- *Barnes DG, Dourson M. 1988. Reference dose (RfD): Description and use in health risk assessments. *Regul Toxicol Pharmacol* 8(4):471-486.
- *Barnes D, Bellin J, DeRosa C, et al. 1987. Reference dose (RfD): Description and use in health risk assessments. Volume I, Appendix A: Integrated risk information system supportive documentation. Washington, DC: US Environmental Protection Agency, Office of Health and Environmental Assessment. EPA600886032a.
- *Brothers RF. 1989. Written communication (May 5) to Director, Office of External Affairs, Agency for Toxic Substances and Disease Registry, regarding manufacture of BCME. Director, Issues Development, Eastman Kodak Company.
- Bruner F, Bertonic G, Severni C. 1978. Gas chromatographic determination of bis(chloromethyl) ether emissions in the atmosphere of industrial plants. *Anal Chem* 50:53.
- *Callahan MA, Slimak MW, Gabel NW, et al. 1979. Water-related environmental fate of 129 priority pollutants. Vol. II. Halogenated aliphatic hydrocarbons, halogenated ethers, monocyclic aromatics, phthalate esters, polycyclic aromatic hydrocarbons, nitrosamines, and miscellaneous compounds. Washington, DC: U.S. Environmental Protection Agency. EPA 440479029b. PB80-2043816.
- Chovil AC. 1979. Occupational lung cancer and smoking: A review in the light of current theories of carcinogenesis. *Chem Manuf Assoc J* 121:548-555.

* Cited in text

8. REFERENCES

- *Clewell HJ, Andersen ME. 1985. Risk assessment extrapolations and physiological modeling. *Toxicol Ind Health* 1(4):111-131.
- *CLPSD. 1988. Contract laboratory program statistical database. Viar and Company, Alexandria VA: September 6, 1988.
- *Cole P, Merletti F. 1980. Chemical agents and occupational cancer. *J Environ Pathol* 3:399-417.
- *Collier L. 1972. Determination of bis-chloromethyl ether at the ppb level in air samples by high-resolution mass spectrometry. *Environ Sci Technol* 6:930-932.
- *Collingwood KW, Pasternack BS, Shore RE. 1987. Industry-wide study of respiratory cancer in chemical workers exposed to chloromethyl ether. *J Nat Cancer Inst* 78:1127-1135.
- *Cupitt LT. 1980. Fate of toxic and hazardous materials in the air environment. Research Triangle Park, NC: U.S. Environmental Sciences Research Laboratory. PB8022194b.
- *DOE. 2016a. Table 3: Protective Action Criteria (PAC) Rev. 29 based on applicable 60-minute AEGLs, ERPGs, or TEELs. The chemicals are listed by CASRN. May 2016. Oak Ridge, TN: U.S. Department of Energy. https://sp.eota.energy.gov/pac/teel/Revision_29_Table3.pdf. February 28, 2017.
- *DOE. 2016.b Protective Action Criteria (PAC) with AEGLs, ERPGs, & TEELs: Rev. 29 for Chemicals of Concern - May 2016. Oak Ridge, TN: U.S. Department of Energy. <https://energy.gov/ehss/protective-action-criteria-pac-aegls-erpgs-teels-rev-29-chemicals-concern-may-2016>. February 28, 2017.
- *DeFonso LR, Kelton SC. 1976. Lung cancer following exposure to chloromethyl methyl ether. *Arch Environ Health* 31:125-130.
- *Drew R, Laskin S, Kuschner M, et al. 1975. Inhalation carcinogenicity of alpha halo ethers. *Arch Environ Health* 30:61-69.
- EPA. 1975. Investigation of selected potential environmental contaminants, haloethers. Washington, DC: U.S. Environmental Protection Agency. NTIS PB246356.
- *EPA. 1979. Toxic Substances Control Act. Chemical substances inventory. Washington, DC: U.S. Environmental Protection Agency, Office of Toxic Substances.
- *EPA. 1980a. An exposure and risk assessment for bis(chloromethyl) ether. Washington, DC: U.S. Environmental Protection Agency, Office of Water Regulations and Standards (Interim Draft).
- *EPA. 1980b. U.S. Environmental Protection Agency. Hazardous waste; identification and listing; final and interim rules. *Fed Regist* 45:3384-33133.
- EPA. 1980c. Guidelines and methodology used in the preparation of health effect assessment chapters of the consent decree water criteria documents. U.S. Environmental Protection Agency. *Fed Regist* 45:79347-79357.
- EPA. 1980d. Ambient water quality criteria document for haloethers. Washington, DC: U.S. Environmental Protection Agency, Office of Water Regulations and Standards. EPA 440580050. NTIS PB81117616.
- *EPA. 1985. U.S. Environmental Protection Agency. Part II. Notification requirements; reportable quantity adjustments; final rule and proposed rule. *Fed Regist* 50:13456-13522.
- *EPA. 1987a. Health and environmental effects document for haloethers. Cincinnati, OH: U.S. Environmental Protection Agency, Office of Health and Environmental Assessment. ECAOCING014.
- *EPA 1987b. U.S. Environmental Protection Agency. Part II. Hazardous substances, reportable quantity adjustments; proposed rules. *Fed Regist* 52:8140-8171.
- *EPA. 1987c. U.S. Environmental Protection Agency. Part II. Extremely hazardous substances list and threshold planning quantities; emergency planning and release notification requirements; final rule. *Fed Regist* 52:13378-13410.
- *EPA. 1988. Integrated risk information system (IRIS). Printout for bis(chloromethyl) ether. Washington, DC: U.S. Environmental Protection Agency. November.
- *EPA. 2005. Toxic chemical release inventory reporting forms and instructions: Revised 2004 version. Section 313 of the Emergency Planning and Community Right-to-Know Act (Title III of the

8. REFERENCES

- Superfund Amendments and Reauthorization Act of 1986). U.S. Environmental Protection Agency, Office of Environmental Information. EPA260B05001.
- *EPA. 2009. National primary drinking water regulations. Washington, DC: Office of Ground Water and Drinking Water, U.S. Environmental Protection Agency. EPA 816-F-09-0004. https://www.epa.gov/sites/production/files/2016-06/documents/npwdr_complete_table.pdf. February 28, 2017.
- *EPA. 2012. Drinking water standards and health advisories. Washington, DC: Office of Water, U.S. Environmental Protection Agency. EPA 822-S-12-001. <https://www.epa.gov/sites/production/files/2015-09/documents/dwstandards2012.pdf>. May 11, 2017.
- *EPA. 2016. Acute Exposure Guideline Levels (AEGs) values. U.S. Environmental Protection Agency. <https://www.epa.gov/aegl/access-acute-exposure-guideline-levels-aegls-values#chemicals>. February 28, 2017.
- Evans HJ. 1982. Cytogenetic studies on industrial populations exposed to mutagens. In: Bridges BA, Butterworth BE, Weinstein IB, eds. Indicators of genotoxic exposure. Banbury Report 13:325-336.
- *FDA. 2013. Everything added to food in the United States (EAFUS). Washington, DC: U.S. Food and Drug Administration. <http://www.accessdata.fda.gov/scripts/fcn/fcnavigation.cfm?rpt=eafuslisting>. February 28, 2017.
- *Figuroa WC, Razkowski R, Weiss W. 1973. Lung cancer in chloromethyl methyl ether workers. *N Engl J Med* 288:1096-1097.
- *Fishbein L. 1979. Potential halogenated industrial carcinogenic and mutagenic chemicals. III. Alkane halides, alkanols and ethers. *Sci Total Environ* 11:223-257.
- *Frankel LS, McCallum KS, Collier L. 1974. Formation of bis(chloromethyl) ether from formaldehyde and hydrogen chloride. *Environ Sci Technol* 8:356-359.
- Frost JK, Gupta PK, Erozan YS, et al. 1973. Pulmonary cytologic alterations in toxic environmental inhalation. *Human Pathol* 4:521-536.
- *FSTRAC. 1988. Summary of state and federal drinking water standards and guidelines. Federal-State Toxicology and Regulatory Alliance Committee. March, 1988.
- *Gargus JL, Reese WH, Rutter HA. 1969. Induction of lung adenomas in newborn mice by bis(chloromethyl) ether. *Toxicol Appl Pharmacol* 15:9296.
- Gilbert D, Eschenroeder A, Goyer M, et al. 1979. An exposure and risk assessment for bis(chloromethyl) ether. Interim draft. Washington, DC: U.S. Environmental Protection Agency, Office of Water Regulations and Standards.
- Gold LS, Backman GM, Hooper NK, et al. 1987. Ranking the potential carcinogenic hazards to workers from exposures to chemicals that are tumorigenic in rodents. *Environ Health Perspect* 76:211-219.
- Gosselin RE. 1984. Clinical toxicology of commercial products. 5th edition. Baltimore, MD: The Williams and Wilkins Co., 11-183.
- *Gowers DS, DeFonso LR, Schaffer P, et al. 1993. Incidence of respiratory cancer among workers exposed to chloromethyl-ethers. *Am J Epidemiol* 137(1):31-42.
- *HSDB. 1988. Hazardous Substances Data Base - computer printout for bis(chloromethyl) ether. Bethesda, MD: National Library of Medicine, National Toxicology Information Program.
- *IARC. 1982. IARC Monographs on the evaluation of the carcinogenic risk of chemicals to humans. Chemicals, industrial processes and industries associated with cancer in humans. IARC Monographs, Volumes 1 to 29, Supplement 4. Lyon, France: International Agency for Research on Cancer.
- *IARC. 2012. Bis(chloromethyl)ether and chloromethyl methyl ether. IARC Monographs on the evaluation of carcinogenic risks to humans. Volume 100F. Chemical agents and related occupations. Lyon, France: International Agency for Research on Cancer. <http://monographs.iarc.fr/ENG/Monographs/vol100F/mono100F-25.pdf>. May 10, 2017.

8. REFERENCES

- *IARC. 2017. Agents classified by the IARC Monographs , Volumes 1–118. Lyon, France: International Agency for Research on Cancer.
http://monographs.iarc.fr/ENG/Classification/List_of_Classifications.pdf. May 10, 2017.
- *IRIS. 2002. Bis(chloromethyl)ether (BCME); CASRN 542-88-1. Chemical assessment summary. Integrated Risk Information System. Washington, DC: U.S. Environmental Protection Agency. May 11, 2017.
- *Kallos GJ. 1981. Oxygen-induced response enhancement in determination of bis(chloromethyl) ether by gas chromatography with ⁶³Ni electron capture detection. *Anal Chem* 53: 963-965.
- *Krishnan K, Anderson ME, Clewell HJ, et al. 1994. Physiologically based pharmacokinetic modeling of chemical mixtures. In: Yang RSH, ed. *Toxicology of chemical mixtures. Case studies, mechanisms, and novel approaches*. San Diego, CA: Academic Press, 399-437.
- *Krost KJ, Pellizzari ED, Walburn SG, et al. 1982. Collection and analysis of hazardous organic emissions. *Anal Chem* 54: 810-817.
- *Kuschner M, Laskin S, Drew RT, et al. 1975. Inhalation carcinogenicity of alpha halo ethers. *Arch Environ Health* 30:73-77.
- *Laskin S, Kuschner M, Drew RT, et al. 1971. Tumors of the respiratory tract induced by inhalation of bis(chloromethyl) ether. *Arch Environ Health* 23:135-136.
- Lemen RA. 1986. Occupationally induced lung cancer epidemiology. In: Merchant JA, ed. *Occupational respiratory diseases*. U.S. Department of Health and Human Services, 629-656.
- *Lemen RA, Johnson WM, Wagoner JK, et al. 1976 A cytologic observations and cancer incidence following exposure to BCME. *Ann NY Acad Sci* 271:71-80.
- *Leong BK, MacFarland HN, Reese WH. 1971. Induction of long adenomas by chronic inhalation of bis(chloromethyl) ether. *Arch Environ Health* 22:663-666.
- *Leong BKJ, Kociba RI, Jersey GC. 1981. A lifetime study of rats and mice exposed to vapors of bis(chloromethyl) ether. *Toxicol Appl Pharmacol* 58:269-281.
- *Mabey WR, Smith JH, Podoll RT, et al. 1982. Aquatic fate process data for organic priority pollutants. Washington, DC: U.S. Environmental Protection Agency, Office of Water Regulations and Standards. EPA440481014. PB87169090.
- *Maher KV, DeFonso IX. 1987. Respiratory cancer among chloromethyl ether workers. *J Nat Cancer Inst* 78:839-843.
- Martin H, Worthing CR, eds. 1974. *Pesticide manual. Basic information on the chemicals used as active components of pesticides*. 4th ed. Worcester, England: British Crop Protection Council, 177.
- McCallum RI, Woolley V, Petrie A. 1983. Lung cancer associated with chloromethyl methyl ether manufacture: an investigation at two factories in the United Kingdom. *Brit J Ind Med* 40:384-389.
- *Mueller G, Norpoth K, Travenius SZM. 1981. Quantitative determination of bis(chloromethyl) ether (BCME) in the ppb range using portable air sample collectors. *Int Arch Occup Environ Health* 48:325-329.
- *NAS/NRC. 1989. Report of the oversight committee. *Biologic markers in reproductive toxicology*. Washington, DC: National Academy of Sciences, National Research Council, National Academy Press, 15-35.
- *NIOSH. 1972a. Field survey of Dow Chemical Company chloromethyl ether facilities, Midland, MI. Cincinnati, OH: National Institute for Occupational Safety and Health. PB82151598.
- *NIOSH. 1972b. Field survey of Rohm and Haas Company Bridesburg Facilities Philadelphia, PA. Cincinnati, OH: National Institute for Occupational Safety and Health. PB82162561.
- NIOSH. 1973. Hazard review of bis(chloromethyl) ether. Cincinnati, OH: National Institute for Occupational Safety and Health. PB87210779.
- NIOSH. 1974. Survey of Burlington Industries, Inc. Burlington House Finishing Plant, Form Fabrics Plant, Durham Domestic Plant, Brookneal Finishing Plant. PB82151077.
- *NIOSH. 1977. *Manual of Analytical Methods*. Vol. 1. Method 213. Cincinnati, OH: National Institute for Occupational Safety and Health.

8. REFERENCES

- *NIOSH. 1986. NIOSH recommendations for occupational safety and health standards. Atlanta, GA: U.S. Department of Health and Human Services, National Institute for Occupation Safety and Health. September, 1986.
- *NIOSH. 2016. bis-Chloromethyl ether. NIOSH pocket guide to chemical hazards. Atlanta, GA: National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention. <https://www.cdc.gov/niosh/npg/npgd0128.html>. May 11, 2017.
- *NLM. 1988. National Library of Medicine - chemline database printout for bis(chloromethyl) ether. August, 1988.
- *Norpoth K, Mueller G, Zilius Z, et al. 1981. Sensitive spectrophotometric determination of carcinogenic alpha chloro ethers. *Int Arch Occup Environ Health* 49:151-155.
- NTP. 1985. National Toxicology Program. Fourth annual report on carcinogens. Summary 1985. Washington, DC: U.S. Department of Health and Human Services.
- *NTP. 2016. Bis(chloromethyl) ether and technical-grade chloromethyl methyl ether. Report on Carcinogens, Fourteenth Edition. CASRN Index in MS Excel. Research Triangle Park, NC: U.S. Department of Health and Human Services, Public Health Service, National Toxicology Program. https://ntp.niehs.nih.gov/ntp/roc/content/profiles/bis_chloromethyl_ether.pdf. May 11, 2017.
- *OSHA. 1974. Occupational Safety and Health Administration, Department of Labor. Occupational Safety and Health Standards. Carcinogens. *Fed Regist* 39:3756-3796.
- *OSHA. 2016. Subpart Z - Toxic and hazardous substances. Section 1910.1003 - 13 Carcinogens. Occupational Safety and Health Standards. Code of Federal Regulations 29 CFR 1910.1003 <https://www.gpo.gov/fdsys/pkg/CFR-2016-title29-vol6/pdf/CFR-2016-title29-vol6-sec1910-1003.pdf>. May 10, 2017.
- *Pasternack BBS, Shore R, Albert RE. 1977. Occupational exposure to chloromethyl ethers. *J Occup Med* 19:741-746.
- Pepelko WE. 1984. Experimental respiratory carcinogenesis in small laboratory animals. *Environ Res* 33:144-188.
- *Ress JF. 1977. Tight control prevents exposure to bis(chloromethyl) ether. *Occupational Health and Safety*. May/June 1977:40-43.
- *Reznik G, Wagner HH, Atay Z. 1977. Lung cancer following exposure to bis(chloromethyl) ether: A case report. *J Environ Pathol Toxicol* 1:105-111.
- *Roe FJC. 1985. Chloromethylation: Three lung cancer deaths in young men. *The Lancet* 2:268.
- *Sakabe H. 1973. Lung cancer due to exposure to bis(chloromethyl) ether. *Ind Health* 11:145-148.
- Sax NI, Lewis RJ, eds. 1987. *Hawley's condensed chemical dictionary*. 11th edition. New York, NY: Van Nostrand Reinhold Co., 149.
- *Sellakumar AR, Snyder CA, Solomon JJ, et al. 1985. Carcinogenicity of formaldehyde and hydrogen chloride in rats. *Toxicol Appl Pharmacol* 81:401-406.
- *Sittig M. 1985. *Handbook of toxic and hazardous chemicals and carcinogens*. Park Ridge, NJ: Noyes Publications, 133-135.
- Sram RJ, Samkova I, Hola N. 1983. High-dose ascorbic prophylaxis in workers occupationally exposed to halogenated ethers. *J Hyg Epidemiol Microbial Immunol* 27:305-318.
- *Thiess AM, Hey W, Zeller H. 1973. Zur Toxikologie von Dichlordimethylaether-Verdacht auf kanzerogene Wirkung such beim Menschen. *Zentralbl Arbeitsmed* 23:97-102.
- Thomas DC, Whittemore AS. 1988, Methods for testing interactions, with applications to occupational exposures, smoking, and lung cancer. *Am J Ind Med* 13:131-147.
- *Tou JC, Kallos GJ. 1974. Kinetic study of the stabilities of chloromethyl methyl ether and bis(chloromethyl) ether in humid air. *Anal Chem* 46:1866-1869.
- *Tou JC, Westover LB, Sonnabend LF. 1974. Kinetic studies of bis(chloromethyl)ether hydrolysis by mass spectrometry. *J Phys Chem* 78:1096-1098.
- *Travenius SZM. 1982. Formation and occurrence of bis(chloromethyl) ether and its prevention in the chemical industry. *Scand J Work Environ Health* 8 (Suppl. 3):1-86.

8. REFERENCES

- *TRI16 2017. TRI explorer: Providing access to EPA's toxics release inventory data. Washington, DC: Office of Information Analysis and Access. Office of Environmental Information. U.S. Environmental Protection Agency. Toxics Release Inventory. <http://www.epa.gov/triexplorer/>. October 18, 2017.
- *Union Carbide. 1968. Summary of acute toxicity and irritancy studies of bis(chloromethyl)ether. Danbury, CT: Union Carbide Corporation. Rpt 31-85; 1968.
- *USITC. 1987. Synthetic organic chemicals. United States production and sales. 1986. Washington, DC: U.S. International Trade Commission. Pub. 2009.
- Van Duuren BL. 1969. Carcinogenic epoxides, lactones, and halo-ethers and their mode of action. *Ann NY Acad Sci* 163:633-651.
- Van Duuren BL. 1980. Prediction of carcinogenicity based on structure, chemical reactivity and possible metabolic pathways. *J Environ Pathol Toxicol* 3:11-34.
- Van Duuren BL. 1989. Comparison of potency of human carcinogens: vinyl chloride, chloromethyl methyl ether and bis(chloromethyl)ether. *Environ Res* 49:143-151.
- Van Duuren BL, Van Duuren SB. 1988. Chemistry, reactivity and carcinogenicity of chloro ethers. In: Politzer P, Martin FJ, eds. *Chemical carcinogens*. New York, NY: Elsevier, 114-176.
- *Van Duuren BL, Goldschmidt BM, Katz C, et al. 1968. Alpha-haloethers: A new type of alkylating carcinogen. *Arch Environ Health* 16:472-476.
- *Van Duuren BL, Sivak A, Goldschmidt BM, et al. 1969. Carcinogenicity of halo-ethers. *J Nat Cancer Inst* 43:481-486.
- *Van Duuren BL, Goldschmidt BM, Katz C, et al. 1972. Carcinogenicity of halo-ethers. 11. Structure-activity relationships of analogs of bis(chloromethyl) ether. *J Nat Cancer Inst* 48:1431-1439.
- *Weast RC, ed. 1985. *CRC handbook of chemistry and physics*. 66th ed. Boca Raton, FL: CRC Press.
- *Weiss W. 1976. Chloromethyl ethers, cigarettes, cough and cancer. *J Occup Med* 18:194-199.
- Weiss W. 1980. The cigarette factor in lung cancer due to chloromethyl ethers. *J Occup Med* 22:527-529.
- *Weiss W. 1982. Epidemic curve of respiratory cancer due to chloromethyl ethers. *J Nat Cancer Inst* 69:1265-1270.
- *Weiss W, Boucot KR. 1975. The respiratory effects of chloromethyl methyl ether. *J Am Med Assoc* 234:1139-1142.
- *Weiss W. 1989. Lung cancer due to chloromethyl ethers: Bias in cohort definition. *J Occup Med* 31:102-105.
- *Weiss W, Moser RL, Auerbach O. 1979. Lung cancer in chloromethyl ether workers. *Am Rev Respir Disease* 120:1031-1037.
- *Weiss W, Nash D. 1997. An epidemic of lung cancer due to chloromethyl ethers. 30 Years of observation. *J Occup Environ Med* 39(10):1003-1009.
- *WHO. 2010. Guidelines for indoor air quality: Selected pollutants. Geneva, Switzerland: World Health Organization. http://www.euro.who.int/__data/assets/pdf_file/0009/128169/e94535.pdf. January 08, 2014.
- *WHO. 2017. Guidelines for drinking-water quality. Fourth edition incorporating the first addendum. Geneva, Switzerland: World Health Organization. <http://apps.who.int/iris/bitstream/10665/254637/1/9789241549950-eng.pdf?ua=1>. February 28, 2017.
- Wilbourn J. 1978. Evaluation of the carcinogenicity of chemicals: A review of the monograph program of the International Agency for Research on Cancer (1971-1977). *Cancer Res* 38:877-885.
- Wilbourn J, Haroun L, Haseltine E, et al. 1986. Response of experimental animals to human carcinogens: An analysis based upon the IARC Monographs Programme. *Carcinogenesis* 7:1853-1863.
- Windholz M, ed. 1983. *The Merck index*. Tenth edition. Rahway, NJ: Merck & Co., Inc.

8. REFERENCES

- *Zajdela F, Croisy A, Barbin A, et al. 1980. Carcinogenicity of chloroethylene oxide, an ultimate reactive metabolite of vinyl chloride, and bis(chloromethyl) ether after subcutaneous administration and in initiation-promotion experiments in mice. *Cancer Res* 40:352-356.