

CHAPTER 8. REFERENCES

- Abbott BD, Birnbaum LS, Pratt RM. 1987. TCDD-induced hyperplasia of the ureteral epithelium produces hydronephrosis in murine fetuses. *Teratology* 35(3):329-334. <http://doi.org/10.1002/tera.1420350307>.
- Abraham K, Hille A, Ende M, et al. 1994. Intake and fecal excretion of PCDDs, PCDFs, HCB and PCBs (138, 153, 180) in a breast-fed and a formula-fed infant. *Chemosphere* 29(9-11):2279-2286. [http://doi.org/10.1016/0045-6535\(94\)90395-6](http://doi.org/10.1016/0045-6535(94)90395-6).
- Abraham K, Knoll A, Ende M, et al. 1996. Intake, fecal excretion, and body burden of polychlorinated dibenzo-p-dioxins and dibenzofurans in breast-fed and formula-fed infants. *Pediatr Res* 40(5):671-679. <http://doi.org/10.1203/00006450-199611000-00005>.
- Abraham K, Paepke O, Gross A, et al. 1998. Time course of PCDD/PCB concentrations in breast-feeding mothers and their infants. *Chemosphere* 37(9-12):1731-1741.
- Ahlborg UG, Wærn F, Manzoor E, et al. 1989. Effects of combinations of PCDDs/PCDFs given to Sprague-Dawley rats. *Chemosphere* 18(1-6):283-289. [http://doi.org/10.1016/0045-6535\(89\)90132-x](http://doi.org/10.1016/0045-6535(89)90132-x).
- Akahane M, Matsumoto S, Kanagawa Y, et al. 2018. Long-term health effects of PCBs and related compounds: A comparative analysis of patients suffering from Yusho and the general population. *Arch Environ Contam Toxicol* 74(2):203-217. <http://doi.org/10.1007/s00244-017-0486-6>.
- Almemark M, Frostell B, Finnveden G. 1991. Treatment technologies for organochlorine-containing sludges and concentrates from external treatment of pulp and paper wastewaters. *Water Sci Technol* 24(3-4):319-329. <http://doi.org/10.2166/wst.1991.0488>.
- Amendola G, Barna D, Blosser R, et al. 1989. The occurrence and fate of PCDDs and PCDFs in five bleached kraft pulp and paper mills. *Chemosphere* 18(1-6):1181-1188. [http://doi.org/10.1016/0045-6535\(89\)90253-1](http://doi.org/10.1016/0045-6535(89)90253-1).
- Ankley GT, Niemi GJ, Lodge KB, et al. 1993. Uptake of planar polychlorinated biphenyls and 2,3,7,8-substituted polychlorinated dibenzofurans and dibenzo-p-dioxins by birds nesting in the lower Fox River and Green Bay, Wisconsin, USA. *Arch Environ Contam Toxicol* 24(3):332-344. <http://doi.org/10.1007/bf01128731>.
- Aozasa O, Ohta S, Mase Y, et al. 1995. Comparative studies on bioaccumulation of PCDDs and PCDFs in C57BL/6 and DBA/2 mice treated with a mixture by oral administration. *Chemosphere* 30(9):1819-1828. [http://doi.org/10.1016/0045-6535\(95\)00050-i](http://doi.org/10.1016/0045-6535(95)00050-i).
- Aries E, Anderson DR, Fisher R. 2008. Exposure assessment of workers to airborne PCDD/Fs, PCBs and PAHs at an electric arc furnace steelmaking plant in the UK. *Ann Occup Hyg* 52(4):213-225. <http://doi.org/10.1093/annhyg/men011>.
- Arnot JA, Meylan W, Tunkel J, et al. 2009. A quantitative structure-activity relationship for predicting metabolic biotransformation rates for organic chemicals in fish. *Environ Toxicol Chem* 28(6):1168-1177. <http://doi.org/10.1897/08-289.1>.
- Atkinson R. 1991. Atmospheric lifetimes of dibenzo-p-dioxins and dibenzofurans. *Sci Total Environ* 104(1-2):17-33. [http://doi.org/10.1016/0048-9697\(91\)90005-y](http://doi.org/10.1016/0048-9697(91)90005-y).
- ATSDR. 1989. Decision guide for identifying substance-specific data needs related to toxicological profiles; Notice. Agency for Toxic Substances and Disease Registry. *Fed Regist* 54(174):37618-37634.
- ATSDR. 1998. Toxicological profile for chlorinated dibenzo-p-dioxins (CDDs). Agency for Toxic Substances and Disease Registry. <https://www.atsdr.cdc.gov/ToxProfiles/tp104.pdf>. April 23, 2020.
- ATSDR. 2000. Toxicological profile for polychlorinated biphenyls (PCBs). Agency for Toxic Substances and Disease Registry. <https://www.atsdr.cdc.gov/toxprofiles/tp17.pdf>. April 23, 2020.
- ATSDR. 2012. Addendum to the toxicological profile for chlorinated dibenzo-p-dioxins (CDDs). Atlanta, GA: Agency for Toxic Substances and Disease Registry. https://www.atsdr.cdc.gov/toxprofiles/cdds_addendum.pdf. April 23, 2020.

8. REFERENCES

- ATSDR. 2017. Chlorodibenzofurans. Full SPL data. Substance priority list (SPL) resource page. Agency for Toxic Substances and Disease Registry. https://www.atsdr.cdc.gov/spl/resources/ATSDR_2017_SPL_Toxicity_Values.xlsx. July 30, 2020.
- ATSDR. 2019. Chlorodibenzofurans. Full SPL data. Substance priority list (SPL) resource page. Agency for Toxic Substances and Disease Registry.
- Bajanowski T, Furst P, Wilmers K, et al. 2002. Dioxin in infants - an environmental hazard? *Int J Legal Med* 116(1):27-32. <http://doi.org/10.1007/s004140100257>.
- Ballschmiter K, Buchert H, Niemczyk R, et al. 1986. Automobile exhausts versus municipal-waste incineration as sources of the polychloro-dibenzodioxins (PCDD) and -furans (PCDF) found in the environment. *Chemosphere* 15(7):901-915. [http://doi.org/10.1016/0045-6535\(86\)90055-x](http://doi.org/10.1016/0045-6535(86)90055-x).
- Bandiera S, Farrell K, Mason G, et al. 1984a. Comparative toxicities of the polychlorinated dibenzofuran (PCDF) and biphenyl (PCB) mixtures which persist in Yusho victims. *Chemosphere* 13(4):507-512. [http://doi.org/10.1016/0045-6535\(84\)90145-0](http://doi.org/10.1016/0045-6535(84)90145-0).
- Bandiera S, Sawyer T, Romkes M, et al. 1984b. Polychlorinated dibenzofurans (PCDFs): effects of structure on binding to the 2,3,7,8-TCDD cytosolic receptor protein, AHH induction and toxicity. *Toxicology* 32(2):131-144. [http://doi.org/10.1016/0300-483x\(84\)90132-x](http://doi.org/10.1016/0300-483x(84)90132-x).
- Banks YB, Brewster DW, Birnbaum LS. 1990. Age-related changes in dermal absorption of 2,3,7,8-tetrachlorodibenzo-p-dioxin and 2,3,4,7,8-pentachlorodibenzofuran. *Fundam Appl Toxicol* 15(1):163-173. [http://doi.org/10.1016/0272-0590\(90\)90173-h](http://doi.org/10.1016/0272-0590(90)90173-h).
- Bannister R, Safe S. 1987. The effects of receptor antagonists on the AHH induction activity of 2,3,7,8-TCDD in C57BL/6 and DBA/2 mice: 1,3,6,8-tetrachlorodibenzofuran. *Chemosphere* 16(8-9):1739-1742. [http://doi.org/10.1016/0045-6535\(87\)90160-3](http://doi.org/10.1016/0045-6535(87)90160-3).
- Barnes DG, Dourson M. 1988. Reference dose (RfD): description and use in health risk assessments. *Regul Toxicol Pharmacol* 8(4):471-486. [http://doi.org/10.1016/0273-2300\(88\)90047-5](http://doi.org/10.1016/0273-2300(88)90047-5).
- Bartalini A, Munoz-Arnanz J, Marsili L, et al. 2019. Evaluation of PCDD/Fs, dioxin-like PCBs and PBDEs in sperm whales from the Mediterranean Sea. *Sci Total Environ* 653:1417-1425. <http://doi.org/10.1016/j.scitotenv.2018.10.436>.
- Barton DA, McKeown JJ, Brunck R. 1990. A screening study of the treatability of dioxins and furans in bleach plant filtrates and mill wastewaters. *Chemosphere* 20(10-12):1747-1754. [http://doi.org/10.1016/0045-6535\(90\)90338-t](http://doi.org/10.1016/0045-6535(90)90338-t).
- Beck H, Droß A, Mathar W. 1992. Dependence of PCDD and PCDF levels in human milk on various parameters in Germany II. *Chemosphere* 25(7-10):1015-1020. [http://doi.org/10.1016/0045-6535\(92\)90101-v](http://doi.org/10.1016/0045-6535(92)90101-v).
- Beck H, Dross A, Mathar W. 1994. PCDD and PCDF exposure and levels in humans in Germany. *Environ Health Perspect* 102(Suppl 1):173-185. <http://doi.org/10.1289/ehp.94102s1173>.
- Beck H, Eckart K, Mathar W, et al. 1989. Levels of PCDDs and PCDFs in adipose tissue of occupationally exposed workers. *Chemosphere* 18(1-6):507-516. [http://doi.org/10.1016/0045-6535\(89\)90161-6](http://doi.org/10.1016/0045-6535(89)90161-6).
- Beck H, Droß A, Kleemann WJ, et al. 1990a. PCDD and PCDF concentrations in different organs from infants. *Chemosphere* 20(7-9):903-910. [http://doi.org/10.1016/0045-6535\(90\)90199-4](http://doi.org/10.1016/0045-6535(90)90199-4).
- Beck H, Droß A, Mathar W, et al. 1990b. Influence of different regional emissions and cardboard containers on levels of PCDD, PCDF and related compounds in cow milk. *Chemosphere* 21(6):789-798. [http://doi.org/10.1016/0045-6535\(90\)90266-V](http://doi.org/10.1016/0045-6535(90)90266-V).
- Birmingham B, Thorpe B, Frank R, et al. 1989a. Dietary intake of PCDD and PCDF from food in Ontario, Canada. *Chemosphere* 19(1-6):507-512. [http://doi.org/10.1016/0045-6535\(89\)90360-3](http://doi.org/10.1016/0045-6535(89)90360-3).
- Birmingham B, Gilman A, Grant D, et al. 1989b. PCDD/PCDF multimedia exposure analysis for the Canadian population: Detailed exposure estimation. *Chemosphere* 19(1-6):637-642. [http://doi.org/10.1016/0045-6535\(89\)90383-4](http://doi.org/10.1016/0045-6535(89)90383-4).
- Birnbaum LS. 1985. The role of structure in the disposition of halogenated aromatic xenobiotics. *Environ Health Perspect* 61:11-20. <http://doi.org/10.1289/ehp.856111>.

8. REFERENCES

- Birnbaum LS, Decad GM, Matthews HB. 1980. Disposition and excretion of 2,3,7,8-tetrachlorodibenzofuran in the rat. *Toxicol Appl Pharmacol* 55(2):342-352. [http://doi.org/10.1016/0041-008x\(80\)90096-4](http://doi.org/10.1016/0041-008x(80)90096-4).
- Birnbaum LS, Decad GM, Matthews HB, et al. 1981. Fate of 2,3,7,8-tetrachlorodibenzofuran in the monkey. *Toxicol Appl Pharmacol* 57(2):189-196. [http://doi.org/10.1016/0041-008x\(81\)90279-9](http://doi.org/10.1016/0041-008x(81)90279-9).
- Birnbaum LS, Harris MW, Barnhart ER, et al. 1987a. Teratogenicity of three polychlorinated dibenzofurans in C57BL/6N mice. *Toxicol Appl Pharmacol* 90(2):206-216. [http://doi.org/10.1016/0041-008x\(87\)90328-0](http://doi.org/10.1016/0041-008x(87)90328-0).
- Birnbaum LS, Harris MW, Crawford DD, et al. 1987b. Teratogenic effects of polychlorinated dibenzofurans in combination in C57BL/6N mice. *Toxicol Appl Pharmacol* 91(2):246-255. [http://doi.org/10.1016/0041-008x\(87\)90105-0](http://doi.org/10.1016/0041-008x(87)90105-0).
- Bloom M, Vena J, Olson J, et al. 2006. Chronic exposure to dioxin-like compounds and thyroid function among New York anglers. *Environ Toxicol Pharmacol* 21(3):260-267. <http://doi.org/10.1016/j.etap.2005.09.001>.
- Bobet E, Berard MF, Dann T. 1990. The measurement of PCDD and PCDF in ambient air in southwestern Ontario. *Chemosphere* 20(10-12):1439-1445. [http://doi.org/10.1016/0045-6535\(90\)90295-5](http://doi.org/10.1016/0045-6535(90)90295-5).
- Boda H, Nghi TN, Nishijo M, et al. 2018. Prenatal dioxin exposure estimated from dioxins in breast milk and sex hormone levels in umbilical cord blood in Vietnamese newborn infants. *Sci Total Environ* 615:1312-1318. <http://doi.org/10.1016/j.scitotenv.2017.09.214>.
- Boenke A, Ballschmiter K. 1989. Formation of polychlorinated dibenzofurans in the pyrolysis of chlorinated fluorenones and 9,10-anthraquinones at 400°C. *Fresenius Z Anal Chem* 333(7):723-724. <http://doi.org/10.1007/BF00476581>.
- Bonfanti L, Cioni M, Rossi C, et al. 1990. Evaluation of PCDD/PCDF emission from the combined combustion of RDF with coal. *Chemosphere* 20(10-12):1891-1897. [http://doi.org/10.1016/0045-6535\(90\)90357-y](http://doi.org/10.1016/0045-6535(90)90357-y).
- Bopp RF, Gross ML, Tong H, et al. 1991. A major incident of dioxin contamination: sediments of New Jersey estuaries. *Environ Sci Technol* 25(5):951-956. <http://doi.org/10.1021/es00017a019>.
- Borwitzky H, Schramm KW. 1991. Reduction of dioxin concentrations on contaminated surfaces. *Chemosphere* 22(5-6):485-493. [http://doi.org/10.1016/0045-6535\(91\)90060-q](http://doi.org/10.1016/0045-6535(91)90060-q).
- Boutros PC, Bielefeld KA, Pohjanvirta R, et al. 2009. Dioxin-dependent and dioxin-independent gene batteries: comparison of liver and kidney in AHR-null mice. *Toxicol Sci* 112(1):245-256. <http://doi.org/10.1093/toxsci/kfp191>.
- Boverhof DR, Burgoon LD, Tashiro C, et al. 2006. Comparative toxicogenomic analysis of the hepatotoxic effects of TCDD in Sprague Dawley rats and C57BL/6 mice. *Toxicol Sci* 94(2):398-416. <http://doi.org/10.1093/toxsci/kfl100>.
- Bowes CW, Mulvihill MJ, Simoneit BR, et al. 1975a. Identification of chlorinated dibenzofurans in American polychlorinated biphenyls. *Nature* 256(5515):305-307. <http://doi.org/10.1038/256305b0>.
- Bowes GW, Mulvihill MJ, DeCamp MR, et al. 1975b. Gas chromatographic characteristics of authentic chlorinated dibenzofurans; identification of two isomers in American and Japanese polychlorinated biphenyls. *J Agric Food Chem* 23(6):1222-1223. <http://doi.org/10.1021/jf60202a045>.
- Brewster DW, Birnbaum LS. 1987. Disposition and excretion of 2,3,4,7,8-pentachlorodibenzofuran in the rat. *Toxicol Appl Pharmacol* 90(2):243-252. [http://doi.org/10.1016/0041-008x\(87\)90332-2](http://doi.org/10.1016/0041-008x(87)90332-2).
- Brewster DW, Birnbaum LS. 1988. Disposition of 1,2,3,7,8-pentachlorodibenzofuran in the rat. *Toxicol Appl Pharmacol* 95(3):490-498. [http://doi.org/10.1016/0041-008x\(88\)90367-5](http://doi.org/10.1016/0041-008x(88)90367-5).
- Brewster DW, Uraih LC, Birnbaum LS. 1988. The acute toxicity of 2,3,4,7,8-pentachlorodibenzofuran (4PeCDF) in the male Fischer rat. *Fundam Appl Toxicol* 11(2):236-249. [http://doi.org/10.1016/0272-0590\(88\)90148-0](http://doi.org/10.1016/0272-0590(88)90148-0).
- Brewster DW, Banks YB, Clark AM, et al. 1989. Comparative dermal absorption of 2,3,7,8-tetrachlorodibenzo-p-dioxin and three polychlorinated dibenzofurans. *Toxicol Appl Pharmacol* 97(1):156-166. [http://doi.org/10.1016/0041-008x\(89\)90064-1](http://doi.org/10.1016/0041-008x(89)90064-1).

8. REFERENCES

- Brna TG, Kilgroe JD. 1990. Control of PCDD/PCDF emissions from municipal waste combustion systems. *Chemosphere* 20(10-12):1875-1882. [http://doi.org/10.1016/0045-6535\(90\)90355-w](http://doi.org/10.1016/0045-6535(90)90355-w).
- Broman D, Naf C, Rolff C, et al. 1992. Using ratios of stable nitrogen isotopes to estimate bioaccumulation and flux of polychlorinated dibenzo-p-dioxins (PCDDs) and dibenzofurans (PCDFs) in two food chains from the northern Baltic. *Environ Toxicol Chem* 11(3):331-345. <http://doi.org/10.1002/etc.5620110307>.
- Brown JF, Lawton RW. 1984. Polychlorinated biphenyl (PCB) partitioning between adipose tissue and serum. *Bull Environ Contam Toxicol* 33(3):277-280. <http://doi.org/10.1007/bf01625543>.
- Budinsky RA, Rowlands JC, Casteel S, et al. 2008. A pilot study of oral bioavailability of dioxins and furans from contaminated soils: Impact of differential hepatic enzyme activity and species differences. *Chemosphere* 70(10):1774-1786. <http://doi.org/10.1016/j.chemosphere.2007.08.035>.
- Bumb RR, Crummett WB, Cutie SS, et al. 1980. Trace chemistries of fire: A source of chlorinated dioxins. *Science* 210(4468):385-390. <http://doi.org/10.1126/science.6159682>.
- Burkhard LP, Kuehl DW. 1986. N-octanol/water partition coefficients by reverse phase liquid chromatography/mass spectrometry for eight tetrachlorinated planar molecules. *Chemosphere* 15(2):163-167. [http://doi.org/10.1016/0045-6535\(86\)90568-0](http://doi.org/10.1016/0045-6535(86)90568-0).
- Buser HR. 1976. Preparation of qualitative standard mixtures of polychlorinated dibenzo-p-dioxins and dibenzofurans by ultraviolet and γ -irradiation of the octachloro compounds. *J Chromatogr* 129:303-307. [http://doi.org/10.1016/s0021-9673\(00\)87788-2](http://doi.org/10.1016/s0021-9673(00)87788-2).
- Buser HR. 1979. Formation of polychlorinated dibenzofurans (PCDFs) from the pyrolysis of individual PCB isomers. *Chemosphere* 8(3):157-174. [http://doi.org/10.1016/0045-6535\(79\)90065-1](http://doi.org/10.1016/0045-6535(79)90065-1).
- Cai LY, Izumi S, Suzuki T, et al. 2011. Dioxins in ascites and serum of women with endometriosis: a pilot study. *Hum Reprod* 26(1):117-126. <http://doi.org/10.1093/humrep/deq312>.
- Campbell CA, Teschke K, Bert J, et al. 1996. Pharmacokinetic model of dioxin and furan levels in adipose tissue from sawmill work involving chlorophenolate fungicides. *Chemosphere* 33(12):2373-2381. [http://doi.org/10.1016/s0045-6535\(96\)00341-4](http://doi.org/10.1016/s0045-6535(96)00341-4).
- Campin DN, Buckland SJ, Hannah DJ, et al. 1991. The identification of dioxin sources in an integrated wood processing facility. *Water Sci Technol* 24(3-4):65-74. <http://doi.org/10.2166/wst.1991.0463>.
- Cao X, Ji L, Lin X, et al. 2018. Comprehensive diagnosis of PCDD/F emission from three hazardous waste incinerators. *R Soc Open Sci* 5(7):172056. <http://doi.org/10.1098/rsos.172056>.
- CARB. 1989. Determination of body burdens for polychlorinated dibenzo-p-dioxins (PCDDs) and polychlorinated dibenzofurans (PCDFs) in California residents. Sacramento: California Air Resources Board. PB90148289. <https://ntrl.ntis.gov/NTRL/dashboard/searchResults/titleDetail/PB90148289.xhtml>. June 17, 2020.
- CARB. 1990. Ambient concentrations of PCDDs/PCDFs (polychlorinated dibenzodioxins/dibenzofurans) in the South Coast air basin. Sacramento, CA: California Air Resources Board. PB90169970. <https://ntrl.ntis.gov/NTRL/dashboard/searchResults/titleDetail/PB90169970.xhtml>. June 19, 2020.
- Carrier G, Brunet RC, Brodeur J. 1995a. Modeling of the toxicokinetics of polychlorinated dibenzo-p-dioxins and dibenzofurans in mammals, including humans. I. Nonlinear distribution of PCDD/PCDF body burden between liver and adipose tissues. *Toxicol Appl Pharmacol* 131(2):253-266. <http://doi.org/10.1006/taap.1995.1068>.
- Carrier G, Brunet RC, Brodeur J. 1995b. Modeling of the toxicokinetics of polychlorinated dibenzo-p-dioxins and dibenzofurans in mammals, including humans. II. Kinetics of absorption and disposition of PCDDs/PCDFs. *Toxicol Appl Pharmacol* 131(2):267-276. <http://doi.org/10.1006/taap.1995.1069>.
- Carsch S, Thoma H, Hutzinger O. 1986. Leaching of polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans from municipal waste incinerator fly ash by water and organic solvents. *Chemosphere* 15(9-12):1927-1930. [http://doi.org/10.1016/0045-6535\(86\)90485-6](http://doi.org/10.1016/0045-6535(86)90485-6).

8. REFERENCES

- CDC. 2021. Fourth report on human exposure to environmental chemicals, updated tables, (March 2021). Atlanta, GA: Centers for Disease Control and Prevention. <https://www.cdc.gov/exposurereport/>. July 10, 2021.
- Chang KJ, Lu FJ, Tung TC, et al. 1980. Studies on patients with polychlorinated biphenyl poisoning. 2. Determination of urinary coproporphyrin, uroporphyrin, delta-aminolevulinic acid and porphobilinogen. *Res Commun Chem Pathol Pharmacol* 30(3):547-554.
- Chang KJ, Hsieh KH, Lee TP, et al. 1982a. Immunologic evaluation of patients with polychlorinated biphenyl poisoning: determination of phagocyte Fc and complement receptors. *Environ Res* 28(2):329-334. [http://doi.org/10.1016/0013-9351\(82\)90131-1](http://doi.org/10.1016/0013-9351(82)90131-1).
- Chang KJ, Hsieh KH, Tang SY, et al. 1982b. Immunologic evaluation of patients with polychlorinated biphenyl poisoning: evaluation of delayed-type skin hypersensitive response and its relation to clinical studies. *J Toxicol Environ Health* 9(2):217-223. <http://doi.org/10.1080/15287398209530156>.
- Chang JW, Chen HL, Su HJ, et al. 2016. Abdominal obesity and insulin resistance in people exposed to moderate-to-high levels of dioxin. *PLoS ONE* 11(1):e0145818. <http://doi.org/10.1371/journal.pone.0145818>.
- Chang KJ, Hsieh KH, Lee TP, et al. 1981. Immunologic evaluation of patients with polychlorinated biphenyl poisoning: determination of lymphocyte subpopulations. *Toxicol Appl Pharmacol* 61(1):58-63. [http://doi.org/10.1016/0041-008x\(81\)90007-7](http://doi.org/10.1016/0041-008x(81)90007-7).
- Chang RR, Jarman WM, King CC, et al. 1990. Bioaccumulation of PCDDs and PCDFs in food animals III: A rapid cleanup of biological materials using reverse phase adsorbent columns. *Chemosphere* 20(7-9):881-886. [http://doi.org/10.1016/0045-6535\(90\)90196-z](http://doi.org/10.1016/0045-6535(90)90196-z).
- Chang JW, Chen HL, Su HJ, et al. 2010. Dioxin exposure and insulin resistance in Taiwanese living near a highly contaminated area. *Epidemiology* 21(1):56-61. <http://doi.org/10.1097/EDE.0b013e3181c2fc6e>.
- Chao WY, Hsu CC, Guo YL. 1997. Middle-ear disease in children exposed prenatally to polychlorinated biphenyls and polychlorinated dibenzofurans. *Arch Environ Health* 52(4):257-262. <http://doi.org/10.1080/00039899709602195>.
- Chen PH, Hsu ST. 1986. PCB poisoning from toxic rice-bran oil in Taiwan. In: Waid JS, ed. *PCBs and the environment*. Vol. 3. Boca Raton, FL: CRC Press, 27-38.
- Chen PH, Wong CK, Rappe C, et al. 1985a. Polychlorinated biphenyls, dibenzofurans and quaterphenyls in toxic rice-bran oil and in the blood and tissues of patients with PCB poisoning (Yu-Cheng) in Taiwan. *Environ Health Perspect* 59:59-65. <http://doi.org/10.1289/ehp.59-1568079>.
- Chen RC, Tang SY, Miyata H, et al. 1985b. Polychlorinated biphenyl poisoning: correlation of sensory and motor nerve conduction, neurologic symptoms, and blood levels of polychlorinated biphenyls, quaterphenyls, and dibenzofurans. *Environ Res* 37(2):340-348. [http://doi.org/10.1016/0013-9351\(85\)90114-8](http://doi.org/10.1016/0013-9351(85)90114-8).
- Chen YC, Guo YL, Hsu CC, et al. 1992. Cognitive development of Yu-Cheng ('oil disease') children prenatally exposed to heat-degraded PCBs. *JAMA* 268(22):3213-3218.
- Chia LG, Chu FL. 1984. Neurological studies on polychlorinated biphenyl (PCB)-poisoned patients. *Am J Ind Med* 5(1-2):117-126.
- Chia LG, Chu FL. 1985. A clinical and electrophysiological study of patients with polychlorinated biphenyl poisoning. *J Neurol Neurosurg Psychiatry* 48(9):894-901. <http://doi.org/10.1136/jnnp.48.9.894>.
- Choudhry GG, Hutzinger O. 1982. Mechanistic aspects of the thermal formation of halogenated organic compounds including polychlorinated dibenzo-p-dioxins. Part II. Thermochemical generation and destruction of dibenzofurans and dibenzo-p-dioxins. *Toxicological & Environmental Chemistry* 5(1):67-93. <http://doi.org/10.1080/02772248209356970>.
- Choudhry GG, Sundstrom G, Ruzo LO, et al. 1977. Photochemistry of chlorinated diphenyl ethers. *J Agric Food Chem* 25(6):1371-1376. <http://doi.org/10.1021/jf60214a004>.

8. REFERENCES

- Christmann W, Kasiske D, Klöppel KD, et al. 1989a. Combustion of polyvinylchloride - an important source for the formation of PCDD/PCDF. *Chemosphere* 19(1-6):387-392. [http://doi.org/10.1016/0045-6535\(89\)90340-8](http://doi.org/10.1016/0045-6535(89)90340-8).
- Christmann W, Klöppel KD, Partsch H, et al. 1989b. PCDD/PCDF and chlorinated phenols in wood preserving formulations for household use. *Chemosphere* 18(1-6):861-865. [http://doi.org/10.1016/0045-6535\(89\)90210-5](http://doi.org/10.1016/0045-6535(89)90210-5).
- Clark KE, Mackay D. 1991. Dietary uptake and biomagnification of four chlorinated hydrocarbons by guppies. *Environ Toxicol Chem* 10(9):1205-1217. <http://doi.org/10.1002/etc.5620100912>.
- Clement RE, Tosine HM, Ali B. 1985. Levels of polychlorinated dibenzo-p-dioxins and dibenzofurans in wood burning stoves and fireplaces. *Chemosphere* 14(6-7):815-819. [http://doi.org/10.1016/0045-6535\(85\)90194-8](http://doi.org/10.1016/0045-6535(85)90194-8).
- Clement RE, Suter SA, Tosine HM. 1989a. Analysis of large volume water samples near chemical dump sites using the aqueous phase liquid extractor (APLE). *Chemosphere* 18(1-6):133-140. [http://doi.org/10.1016/0045-6535\(89\)90113-6](http://doi.org/10.1016/0045-6535(89)90113-6).
- Clement RE, Tosine HM, Osborne J, et al. 1987a. Emissions of chlorinated organics from a municipal sewage sludge burning incinerator. *Chemosphere* 16(8-9):1895-1900. [http://doi.org/10.1016/0045-6535\(87\)90184-6](http://doi.org/10.1016/0045-6535(87)90184-6).
- Clement RE, Tosine HM, Taguchi V, et al. 1987b. Investigation of American lobster, *Homarus americanus*, for the presence of chlorinated dibenzo-p-dioxins and dibenzofurans. *Bull Environ Contam Toxicol* 39(6):1069-1075. <http://doi.org/10.1007/bf01689600>.
- Clement RE, Tashiro C, Suter S, et al. 1989b. Chlorinated dibenzo-p-dioxins (CDDs) and dibenzofurans (CDFs) in effluents and sludges from pulp and paper mills. *Chemosphere* 18(1-6):1189-1197. [http://doi.org/10.1016/0045-6535\(89\)90254-3](http://doi.org/10.1016/0045-6535(89)90254-3).
- Clement RE, Suter SA, Reiner E, et al. 1989c. Concentrations of chlorinated dibenzo-p-dioxins and dibenzofurans in effluents and centrifuged particulates from Ontario pulp and paper mills. *Chemosphere* 19(1-6):649-654. [http://doi.org/10.1016/0045-6535\(89\)90385-8](http://doi.org/10.1016/0045-6535(89)90385-8).
- Couture LA, Harris MW, Birnbaum LS. 1989. Developmental toxicity of 2,3,4,7,8-pentachlorodibenzofuran in the Fischer 344 rat. *Fundam Appl Toxicol* 12(2):358-366. [http://doi.org/10.1016/0272-0590\(89\)90052-3](http://doi.org/10.1016/0272-0590(89)90052-3).
- Cramer G, Bolger M, Henry S, et al. 1991. USFDA assessment of exposure to 2,3,7,8-TCDD and 2,3,7,8-TCDF from foods contacting bleached paper products. *Chemosphere* 23(8-10):1537-1550. [http://doi.org/10.1016/0045-6535\(91\)90179-h](http://doi.org/10.1016/0045-6535(91)90179-h).
- Crandall MS, Kinnes GM, Hartle RW. 1992. Levels of chlorinated dioxins and furans in three occupational environments. *Chemosphere* 25(1-2):255-258. [http://doi.org/10.1016/0045-6535\(92\)90527-x](http://doi.org/10.1016/0045-6535(92)90527-x).
- Crofton KM, Craft ES, Hedge JM, et al. 2005. Thyroid-hormone-disrupting chemicals: evidence for dose-dependent additivity or synergism. *Environ Health Perspect* 113(11):1549-1554. <http://doi.org/10.1289/ehp.8195>.
- Czuczwa JM, Hites RA. 1986a. Sources and fate of PCDD and PCDF. *Chemosphere* 15(9-12):1417-1420. [http://doi.org/10.1016/0045-6535\(86\)90419-4](http://doi.org/10.1016/0045-6535(86)90419-4).
- Czuczwa JM, Hites RA. 1986b. Airborne dioxins and dibenzofurans: sources and fates. *Environ Sci Technol* 20(2):195-200. <http://doi.org/10.1021/es00144a015>.
- Czuczwa JM, Niessen F, Hites RA. 1985. Historical record of polychlorinated dibenzo-p-dioxins and dibenzofurans in Swiss lake sediments. *Chemosphere* 14(9):1175-1179. [http://doi.org/10.1016/0045-6535\(85\)90139-0](http://doi.org/10.1016/0045-6535(85)90139-0).
- Czuczwa J, Katona V, Pitts G, et al. 1989. Analysis of fog samples for PCDD and PCDF. *Chemosphere* 18(1-6):847-850. [http://doi.org/10.1016/0045-6535\(89\)90207-5](http://doi.org/10.1016/0045-6535(89)90207-5).
- Dahl P, Lindstrom G, Wiberg K, et al. 1995. Absorption of polychlorinated biphenyls, dibenzo-p-dioxins and dibenzofurans by breast-fed infants. *Chemosphere* 30(12):2297-2306. [http://doi.org/10.1016/0045-6535\(95\)00102-e](http://doi.org/10.1016/0045-6535(95)00102-e).

8. REFERENCES

- Dahmer SC, Tetreault GR, Hall RI, et al. 2015. Historical decline and altered congener patterns of polychlorinated dibenzo-p-dioxins and dibenzofurans in fish and sediment in response to process changes at a pulp mill discharging into Jackfish Bay, Lake Superior. *Environ Toxicol Chem* 34(11):2489-2502. <http://doi.org/10.1002/etc.3126>.
- Davis D, Safe S. 1988. Immunosuppressive activities of polychlorinated dibenzofuran congeners: quantitative structure-activity relationships and interactive effects. *Toxicol Appl Pharmacol* 94(1):141-149. [http://doi.org/10.1016/0041-008x\(88\)90344-4](http://doi.org/10.1016/0041-008x(88)90344-4).
- De Jongh J, Belfroid A, Sinnige T, et al. 1992. Disposition, elimination and enzyme induction of 1,2,3,7,8-PnCDD, 1,2,3,6,7,8-HxCDD and 2,3,4,7,8-PnCDF in the liver of the mouse after a single oral dose. *Chemosphere* 25(12):1851-1859.
- De Jongh J, Nieboer R, Schrodgers I, et al. 1993. Toxicokinetic mixture interactions between chlorinated aromatic hydrocarbons in the liver of the C57BL/6J mouse: 2. Polychlorinated dibenzo-p-dioxins (PCDDs), dibenzofurans (PCDFs) and biphenyls (PCBs). *Arch Toxicol* 67(9):598-604. <http://doi.org/10.1007/bf01974066>.
- De Vault D, Dunn W, Bergqvist P-A, et al. 1989. Polychlorinated dibenzofurans and polychlorinated dibenzo-p-dioxins in great lakes fish: A baseline and interlake comparison. *Environ Toxicol Chem* 8(11):1013-1022. <http://doi.org/10.1002/etc.5620081107>.
- Decad GM, Birnbaum LS, Matthews HB. 1981a. 2,3,7,8-Tetrachlorodibenzofuran tissue distribution and excretion in guinea pig. *Toxicol Appl Pharmacol* 57(2):231-240. [http://doi.org/10.1016/0041-008x\(81\)90284-2](http://doi.org/10.1016/0041-008x(81)90284-2).
- Decad GM, Birnbaum LS, Matthews HB. 1981b. Distribution and excretion of 2,3,7,8-tetrachlorodibenzofuran in C57BL/6J and DBA/2J mice. *Toxicol Appl Pharmacol* 59(3):564-573. [http://doi.org/10.1016/0041-008x\(81\)90311-2](http://doi.org/10.1016/0041-008x(81)90311-2).
- Denison MS, Soshilov AA, He G, et al. 2011. Exactly the same but different: promiscuity and diversity in the molecular mechanisms of action of the aryl hydrocarbon (dioxin) receptor. *Toxicol Sci* 124(1):1-22. <http://doi.org/10.1093/toxsci/kfr218>.
- des Rosiers PE. 1983. Remedial measures for wastes containing polychlorinated dibenzo-p-dioxins (PCDDs) and dibenzofurans (PCDFs): destruction, containment or process modification. *Ann Occup Hyg* 27(1):57-72. <http://doi.org/10.1093/annhyg/27.1.57>.
- des Rosiers PE. 1986. Methodologies for materials contaminated with PCDDs and related compounds. *Chemosphere* 15(9-12):1513-1528. [http://doi.org/10.1016/0045-6535\(86\)90432-7](http://doi.org/10.1016/0045-6535(86)90432-7).
- des Rosiers PE. 1987. National dioxin study. *ACS Symp Ser* 338:34-53. <http://doi.org/10.1021/bk-1987-0338.ch003>.
- des Rosiers PE, Lee A. 1986. PCBs fires: Correlation of chlorobenzene isomer and PCB homolog contents of PCB fluids with PCDD and PCDF contents of soot. *Chemosphere* 15(9-12):1313-1132. [http://doi.org/10.1016/0045-6535\(86\)90406-6](http://doi.org/10.1016/0045-6535(86)90406-6).
- DeVito MJ, Ross DR, Birnbaum LS. 1995. Disposition of PCDD/PCDF in mice. *Organohalogen Compounds* 25:11-15. <http://dioxin20xx.org/wp-content/uploads/pdfs/1995/95-68.pdf>. October 7, 2019.
- DeVito MJ, Maier WE, Diliberto JJ, et al. 1993. Comparative ability of various PCBs, PCDFs, and TCDD to induce cytochrome P450 1A1 and 1A2 activity following 4 weeks of treatment. *Fundam Appl Toxicol* 20(1):125-130.
- DeVito MJ, Diliberto JJ, Ross DG, et al. 1997. Dose-response relationships for polyhalogenated dioxins and dibenzofurans following subchronic treatment in mice. I. CYP1A1 and CYP1A2 enzyme activity in liver, lung, and skin. *Toxicol Appl Pharmacol* 147(2):267-280. <http://doi.org/10.1006/taap.1997.8261>.
- DeVito MJ, Ross DG, Dupuy AE, et al. 1998. Dose-response relationships for disposition and hepatic sequestration of polyhalogenated dibenzo-p-dioxins, dibenzofurans, and biphenyls following subchronic treatment in mice. *Toxicol Sci* 46(2):223-234. <http://doi.org/10.1006/toxs.1998.2530>.

8. REFERENCES

- Dewailly E, Weber JP, Gingras S, et al. 1991. Coplanar PCBs in human milk in the province of Quebec, Canada: are they more toxic than dioxin for breast fed infants? *Bull Environ Contam Toxicol* 47(4):491-498. <http://doi.org/10.1007/bf01700935>.
- Dickerson R, Howie L, Davis D, et al. 1990. The structure-dependent effects of heptachlorodibenzofuran isomers in male C57BL/6 mice: immunotoxicity and monooxygenase enzyme induction. *Fundam Appl Toxicol* 15(2):298-307. [http://doi.org/10.1016/0272-0590\(90\)90056-p](http://doi.org/10.1016/0272-0590(90)90056-p).
- Diliberto JJ, Burgin D, Birnbaum LS. 1997. Role of CYP1A2 in hepatic sequestration of dioxin: studies using CYP1A2 knock-out mice. *Biochem Biophys Res Commun* 236(2):431-433. <http://doi.org/10.1006/bbrc.1997.6973>.
- Diliberto JJ, Burgin DE, Birnbaum LS. 1999. Effects of CYP1A2 on disposition of 2,3,7, 8-tetrachlorodibenzo-p-dioxin, 2,3,4,7,8-pentachlorodibenzofuran, and 2,2',4,4',5,5'-hexachlorobiphenyl in CYP1A2 knockout and parental (C57BL/6N and 129/Sv) strains of mice. *Toxicol Appl Pharmacol* 159(1):52-64. <http://doi.org/10.1006/taap.1999.8720>.
- Diliberto JJ, Akubue PI, Luebke RW, et al. 1995. Dose-response relationships of tissue distribution and induction of Cyp1A1 and Cyp1A2 enzymatic activities following acute exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in mice. *Toxicol Appl Pharmacol* 130(2):197-208. <http://doi.org/10.1006/taap.1995.1025>.
- DOE. 2018a. Table 3: Protective Action Criteria (PAC) Rev. 29a based on applicable 60-minute AEGLs, ERPGs, or TEELs. The chemicals are listed by CASRN. June 2018. Oak Ridge, TN: U.S. Department of Energy. https://edms.energy.gov/pac/docs/Revision_29A_Table3.pdf. April 12, 2020.
- DOE. 2018b. Protective Action Criteria (PAC) with AEGLs, ERPGs, & TEELs: Rev. 29A, June 2018. Oak Ridge, TN: U.S. Department of Energy. <https://edms.energy.gov/pac/>. April 12, 2020.
- Doyle EA, Fries GF. 1986. Induction of aryl hydrocarbon hydroxylase by chlorinated dibenzofurans in rats. *Chemosphere* 15(9-12):1745-1748. [http://doi.org/10.1016/0045-6535\(86\)90462-5](http://doi.org/10.1016/0045-6535(86)90462-5).
- Draper WM, Park H, Stephens RD. 1991. Determination of OCDD and OCDF in soils and biological samples by GC-ECD. In: Friedman D, ed. *Waste testing and quality assurance*. Vol. 3. West Conshohocken, PA: ASTM International, 87-105.
- Drechsler WD. 1986. Destruction of PCDD/PCDF by non-thermal methods. *Chemosphere* 15(9-12):1529-1534. [http://doi.org/10.1016/0045-6535\(86\)90433-9](http://doi.org/10.1016/0045-6535(86)90433-9).
- Eadon G, Kaminsky L, Silkworth J, et al. 1986. Calculation of 2,3,7,8-TCDD equivalent concentrations of complex environmental contaminant mixtures. *Environ Health Perspect* 70:221-227. <http://doi.org/10.1289/ehp.8670221>.
- Edgerton SA, Czuczwa JM, Rench JD, et al. 1989. Ambient air concentrations of polychlorinated dibenzo-p-dioxins and dibenzofurans in Ohio: Sources and health risk assessment. *Chemosphere* 18(9-10):1713-1730. [http://doi.org/10.1016/0045-6535\(89\)90455-4](http://doi.org/10.1016/0045-6535(89)90455-4).
- Eitzer BD, Hites RA. 1988. Vapor pressures of chlorinated dioxins and dibenzofurans. *Environ Sci Technol* 22(11):1362-1364. <http://doi.org/10.1021/es00176a018>.
- Eitzer BD, Hites RA. 1989a. Atmospheric transport and deposition of polychlorinated dibenzo-p-dioxins and dibenzofurans. *Environ Sci Technol* 23(11):1396-1401. <http://doi.org/10.1021/es00069a011>.
- Eitzer BD, Hites RA. 1989b. Polychlorinated dibenzo-p-dioxins and dibenzofurans in the ambient atmosphere of Bloomington, Indiana. *Environ Sci Technol* 23(11):1389-1395. <http://doi.org/10.1021/es00069a010>.
- Elliott JE, Butler RW, Norstrom RJ, et al. 1989. Environmental contaminants and reproductive success of great blue herons *Ardea herodias* in British Columbia, 1986-1987. *Environ Pollut* 59(2):91-114. [http://doi.org/10.1016/0269-7491\(89\)90099-7](http://doi.org/10.1016/0269-7491(89)90099-7).
- EPA. 1985. Health assessment document for polychlorinated dibenzo-p-dioxins. U.S. Environmental Protection Agency. EPA600884014F. <https://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=30001CBN.txt>. April 24, 2020.

8. REFERENCES

- EPA. 1989a. NHATS broad scan analysis: population estimates from fiscal year 1982 specimens. Washington, DC: U.S. Environmental Protection Agency. EPA56059000. <https://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=91012GAH.txt>. June 17, 2020.
- EPA. 1989b. Interim procedures for estimating risks associated with exposure to mixtures of chlorinated dibenzo-p-dioxins and dibenzofurans (CDDs and CDFs) and 1989 update. Washington, DC: U.S. Environmental Protection Agency. EPA625389016. <https://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=300047JJ.txt>. June 17, 2020.
- EPA. 1990a. Waste analysis and identification and listing of hazardous waste. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 264-341 and 40 CFR 261, App VII.
- EPA. 1990b. Interim methods for development of inhalation reference concentrations. Washington, DC: U.S. Environmental Protection Agency. EPA600890966A. <https://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=30001JXM.txt>. June 19, 2020.
- EPA. 1991. Proposed regulation of land application of sludge from pulp and paper mills using chlorine and chlorine derivative bleaching processes. U.S. Environmental Protection Agency. Fed Regist 56(91):21802-21803. <https://www.loc.gov/item/fr056091/>. June 17, 2020.
- EPA. 1994. Method 1613: Tetra- through octa-chlorinated dioxins and furans by isotope dilution HRGC/HRMS. Washington, DC: U.S. Environmental Protection Agency. https://www.epa.gov/sites/production/files/2015-08/documents/method_1613b_1994.pdf. April 22, 2020.
- EPA. 2005. Toxic chemical release inventory reporting forms and instructions: Revised 2004 version. Section 313 of the Emergency Planning and Community Right-to-Know Act (Title III of the Superfund Amendments and Reauthorization Act of 1986). U.S. Environmental Protection Agency. EPA260B05001. <https://19january2017snapshot.epa.gov/sites/production/files/documents/ry2004rfi.pdf>. June 19, 2020.
- EPA. 2007a. 2001 National sewage sludge survey report. Washington, DC: U.S. Environmental Protection Agency. EPA822R07006. <https://www.epa.gov/biosolids/sewage-sludge-surveys>. April 23, 2020.
- EPA. 2007b. Method 8280B: Polychlorinated dibenzo-p-dioxins (pcdds) and polychlorinated dibenzofurans (PCDFs) by high-resolution gas chromatography/low-resolution mass spectrometry (HRGC/LRMS). U.S. Environmental Protection Agency. https://www.epa.gov/sites/production/files/2015-12/documents/8280b_0.pdf. April 23, 2020.
- EPA. 2009. National primary drinking water regulations. Washington, DC: U.S. Environmental Protection Agency. EPA816F090004. https://www.epa.gov/sites/production/files/2016-06/documents/npwdr_complete_table.pdf. September 7, 2017.
- EPA. 2013. National Dioxin Air Monitoring Network (NDAMN): Report of the results of atmospheric measurements of polychlorinated dibenzo-p-dioxins (PCDDs), polychlorinated dibenzofurans (PCDFs), and dioxin-like polychlorinated biphenyls (PCBs) in rural and remote areas of the United States from June 1998 through November 2004. Washington, DC: U.S. Environmental Protection Agency. EPA600R13183F. PB2014104547. http://ofmpub.epa.gov/eims/eimscomm.getfile?p_download_id=515193. April 20, 2020.
- EPA. 2018a. 2018 Edition of the drinking water standards and health advisories. Washington, DC: U.S. Environmental Protection Agency. EPA822S12001. <https://www.epa.gov/sites/production/files/2018-03/documents/dwtable2018.pdf>. July 25, 2018.
- EPA. 2018b. Acute Exposure Guideline Levels (AEGs) values. U.S. Environmental Protection Agency. https://www.epa.gov/sites/production/files/2018-08/documents/compiled_aegls_update_27jul2018.pdf. April 12, 2020.
- Erickson MD. 1989. PCDFs and related compounds produced from PCB fires - A review. Chemosphere 19(1-6):161-165. [http://doi.org/10.1016/0045-6535\(89\)90305-6](http://doi.org/10.1016/0045-6535(89)90305-6).

8. REFERENCES

- Everett CJ, Thompson OM. 2014. Dioxins, furans and dioxin-like PCBs in human blood: causes or consequences of diabetic nephropathy? *Environ Res* 132:126-131. <http://doi.org/10.1016/j.envres.2014.03.043>.
- Everett CJ, Thompson OM. 2016. Association of dioxins, furans and dioxin-like PCBs in human blood with nephropathy among US teens and young adults. *Rev Environ Health* 31(2):195-201. <http://doi.org/10.1515/reveh-2015-0031>.
- Fahrig R, Nilsson C, Rappe C. 1978. Genetic activity of chlorophenols and chlorophenol impurities. *Environmental Science and Research* 12:325-338. http://doi.org/10.1007/978-1-4615-8948-8_28.
- Farrell K, Safe L, Safe S. 1987. Synthesis and aryl hydrocarbon receptor binding properties of radiolabeled polychlorinated dibenzofuran congeners. *Arch Biochem Biophys* 259(1):185-195. [http://doi.org/10.1016/0003-9861\(87\)90485-1](http://doi.org/10.1016/0003-9861(87)90485-1).
- FDA. 2007. Dioxin analysis results/exposure estimates. U.S. Food and Drug Administration. <https://www.fda.gov/food/chemicals/dioxin-analysis-resultsexposure-estimates>. April 23, 2020.
- FDA. 2020. Substances added to food. Washington, DC: U.S. Food and Drug Administration. <https://www.accessdata.fda.gov/scripts/fdcc/?set=FoodSubstances>. April 12, 2020.
- Finley B, Fehling K, Warmerdam J, et al. 2009. Oral bioavailability of polychlorinated dibenzo-p-dioxins/dibenzofurans in industrial soils. *Human and Ecological Risk Assessment: An International Journal* 15(6):1146-1167. <http://doi.org/10.1080/10807030903304765>.
- Fishbein L. 1992. Exposure from occupational versus other sources. *Scand J Work Environ Health* 18 Suppl 1:5-16.
- Flesch-Janys D, Becher H, Gurn P, et al. 1996. Elimination of polychlorinated dibenzo-p-dioxins and dibenzofurans in occupationally exposed persons. *J Toxicol Environ Health* 47(4):363-378. <http://doi.org/10.1080/009841096161708>.
- Fletcher N, Wahlstrom D, Lundberg R, et al. 2005. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) alters the mRNA expression of critical genes associated with cholesterol metabolism, bile acid biosynthesis, and bile transport in rat liver: a microarray study. *Toxicol Appl Pharmacol* 207(1):1-24. <http://doi.org/10.1016/j.taap.2004.12.003>.
- Frank APCG, Schrap SM. 1990. Bioaccumulation of some polychlorinated dibenzo-p-dioxins and octachlorodibenzofuran in the guppy (*Poecilia reticulata*). *Chemosphere* 20(5):495-512. [http://doi.org/10.1016/0045-6535\(90\)90107-5](http://doi.org/10.1016/0045-6535(90)90107-5).
- Fricker C, Hardy JK. 1990. Characterization of commercially available coffee filter papers. *J Environ Sci Health A* 25(8):927-936. <http://doi.org/10.1080/10934529009375609>.
- Friesen KJ, Vilks J, Muir DCG. 1990. Aqueous solubilities of selected 2,3,7,8-substituted polychlorinated dibenzofurans (PCDFs). *Chemosphere* 20(1-2):27-32. [http://doi.org/10.1016/0045-6535\(90\)90084-7](http://doi.org/10.1016/0045-6535(90)90084-7).
- Frommberger R. 1991. Cork products: a potential source of polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans. *Chemosphere* 23(2):133-139. [http://doi.org/10.1016/0045-6535\(91\)90102-j](http://doi.org/10.1016/0045-6535(91)90102-j).
- Fu YA. 1984. Ocular manifestation of polychlorinated biphenyls intoxication. *Am J Ind Med* 5(1-2):127-132.
- Funatsu I, Yamashita F, Yoshikane T, et al. 1971. [A chlorobiphenyl induced fetopathy]. *Fukuoka Igaku Zasshi* 62:139-149. (Japanese)
- Fürst P, Fürst C, Groebel W. 1990. Levels of PCDDs and PCDFs in food-stuffs from the Federal Republic of Germany. *Chemosphere* 20(7-9):787-792. [http://doi.org/10.1016/0045-6535\(90\)90183-t](http://doi.org/10.1016/0045-6535(90)90183-t).
- Fürst P, Fürst C, Wilmers K. 1992. PCDDs and PCDFs in human milk-statistical evaluation of a 6-years survey. *Chemosphere* 25(7-10):1029-1038. [http://doi.org/10.1016/0045-6535\(92\)90103-x](http://doi.org/10.1016/0045-6535(92)90103-x).
- Fürst P, Krüger C, Meemken H, et al. 1989. PCDD and PCDF levels in human milk - dependence on the period of lactation. *Chemosphere* 18:439-444.

8. REFERENCES

- Garå A, Andersson K, Nilsson C, et al. 1981. Synthesis of halogenated diphenyl ethers and dibenzofurans - A discussion of specific isomers available. *Chemosphere* 10(4):365-390. [http://doi.org/10.1016/0045-6535\(81\)90233-2](http://doi.org/10.1016/0045-6535(81)90233-2).
- Gardner AM, White KD. 1990. Polychlorinated dibenzofurans in the edible portion of selected fish. *Chemosphere* 21(1-2):215-222. [http://doi.org/10.1016/0045-6535\(90\)90393-8](http://doi.org/10.1016/0045-6535(90)90393-8).
- Garner AJ, Pagano JJ. 2019. Trends of polychlorinated dioxins, polychlorinated furans, and dioxin-like polychlorinated biphenyls in Chinook and Coho salmonid eggs from a Great Lakes tributary. *Environ Pollut* 247:1039-1045. <http://doi.org/10.1016/j.envpol.2019.01.117>.
- Geyer HJ, Scheunert I, Korte F. 1987. Correlation between the bioconcentration potential of organic environmental chemicals in humans and their n-octanol/water partition coefficients. *Chemosphere* 16(1):239-252. [http://doi.org/10.1016/0045-6535\(87\)90128-7](http://doi.org/10.1016/0045-6535(87)90128-7).
- Gibbs J, Hansen M, Ferrario J. 2003. Ambient air sampling for dioxins, furans and coplanar PCBs in an urban industrialized corridor in Calcasieu Parish, Louisiana. *Organohalogen Compounds* 61:514-517. <http://dioxin20xx.org/wp-content/uploads/pdfs/2003/03-269.pdf>. April 23, 2020.
- Gladen BC, Rogan WJ, Ragan NB, et al. 1988. Urinary porphyrins in children exposed transplacentally to polyhalogenated aromatics in Taiwan. *Arch Environ Health* 43(1):54-58. <http://doi.org/10.1080/00039896.1988.9934374>.
- Gladen BC, Taylor JS, Wu YC, et al. 1990. Dermatological findings in children exposed transplacentally to heat-degraded polychlorinated biphenyls in Taiwan. *Br J Dermatol* 122(6):799-808. <http://doi.org/10.1111/j.1365-2133.1990.tb06269.x>.
- Glasser H, Chang DP, Hickman DC. 1991. An analysis of biomedical waste incineration. *J Air Waste Manage Assoc* 41(9):1180-1188. <http://doi.org/10.1080/10473289.1991.10466913>.
- Glidden RM, Brown PJ, Sittig RA, et al. 1990. Determination of 2,3,7,8-tetrachlorodibenzo-p-dioxin and 2,3,7,8-tetrachlorodibenzofuran in cow's milk. *Chemosphere* 20(10-12):1619-1624. [http://doi.org/10.1016/0045-6535\(90\)90321-j](http://doi.org/10.1016/0045-6535(90)90321-j).
- Goldfarb TD, Harrad SJ. 1991. Consideration of the environmental impact of the volatilization of PCDDs and PCDFs. *Chemosphere* 23(11-12):1669-1674. [http://doi.org/10.1016/0045-6535\(91\)90015-6](http://doi.org/10.1016/0045-6535(91)90015-6).
- Goldstein JA, Safe S. 1989. Mechanism of action and structure-activity relationships for the chlorinated dibenzo-p-dioxins and related compounds. In: Kimbrough RD, Jensen AA, eds. *Halogenated biphenyls, naphthalenes, dibenzodioxins and related compounds*. 2nd ed. New York, NY: Elsevier, 239-293.
- Goldstein JA, Hass JR, Linko P, et al. 1978. 2,3,7,8-Tetrachlorodibenzofuran in a commercially available 99% pure polychlorinated biphenyl isomer identified as the inducer of hepatic cytochrome P-448 and aryl hydrocarbon hydroxylase in the rat. *Drug Metab Dispos* 6(3):258-264.
- Golor G, Yamashita K, McLachlan MS, et al. 1993. Comparison of the kinetics of chlorinated and brominated dioxins and furans in the rat. *Organohalogen Compounds* 13:203-206. <http://dioxin20xx.org/wp-content/uploads/pdfs/1993/93-114.pdf>. October 4, 2019.
- Gorski T. 1981. Presence of polychlorinated dibenzo-p-dioxins in latex nipples. *Bull Environ Contam Toxicol* 27(1):68-71. <http://doi.org/10.1007/bf01610988>.
- Graham M, Hileman FD, Orth RG, et al. 1986. Chlorocarbons in adipose tissue from a Missouri population. *Chemosphere* 15(9-12):1595-1600. [http://doi.org/10.1016/0045-6535\(86\)90442-x](http://doi.org/10.1016/0045-6535(86)90442-x).
- Gray AP, Dipinto VM, Solomon IJ. 1976. Synthesis of specific polychlorinated dibenzofurans. *J Org Chem* 41(14):2428-2434. <http://doi.org/10.1021/jo00876a016>.
- Groce DF, Alley CC, Patterson DG. 1989. Synthesis of tetrachloro-, pentachloro-, and hexachlorodibenzofurans from the pyrolysis of polychlorinated biphenyls. *Chemosphere* 19(1-6):225-232. [http://doi.org/10.1016/0045-6535\(89\)90316-0](http://doi.org/10.1016/0045-6535(89)90316-0).
- Guo YL, Lin CJ, Yao WJ, et al. 1994a. Musculoskeletal changes in children prenatally exposed to polychlorinated biphenyls and related compounds (Yu-cheng children). *J Toxicol Environ Health* 41(1):83-93. <http://doi.org/10.1080/15287399409531828>.

8. REFERENCES

- Guo YL, Chen YC, Yu ML, et al. 1994b. Early development of Yu-Cheng children born seven to twelve years after the Taiwan PCB outbreak. *Chemosphere* 29(9-11):2395-2404.
- Guo YL, Lai TJ, Chen SJ, et al. 1995. Gender-related decrease in Raven's progressive matrices scores in children prenatally exposed to polychlorinated biphenyls and related contaminants. *Bull Environ Contam Toxicol* 55(1):8-13. <http://doi.org/10.1007/bf00212382>.
- Guo YL, Yu ML, Hsu CC, et al. 1999. Chloracne, goiter, arthritis, and anemia after polychlorinated biphenyl poisoning: 14-year follow-Up of the Taiwan Yucheng cohort. *Environ Health Perspect* 107(9):715-719. <http://doi.org/10.1289/ehp.99107715>.
- Guo YL, Hsu PC, Hsu CC, et al. 2000. Semen quality after prenatal exposure to polychlorinated biphenyls and dibenzofurans. *Lancet* 356(9237):1240-1241. [http://doi.org/10.1016/S0140-6736\(00\)02792-6](http://doi.org/10.1016/S0140-6736(00)02792-6).
- Hagenmaier H, Berchtold A. 1986. Analysis of waste from production of Na-pentachlorophenolate for polychlorinated dibenzodioxins (PCDD) and dibenzofurans (PCDF). *Chemosphere* 15(9-12):1991-1994. [http://doi.org/10.1016/0045-6535\(86\)90499-6](http://doi.org/10.1016/0045-6535(86)90499-6).
- Hagenmaier H, Brunner H. 1987. Isomerspecific analysis of pentachlorophenol and sodium pentachlorophenate for 2,3,7,8-substituted PCDD and PCDF at sub-ppb levels. *Chemosphere* 16(8-9):1759-1764. [http://doi.org/10.1016/0045-6535\(87\)90164-0](http://doi.org/10.1016/0045-6535(87)90164-0).
- Hagenmaier H, She J, Lindig C. 1992. Persistence of polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans in contaminated soil at Maulach and Rastatt in Southwest Germany. *Chemosphere* 25(7-10):1449-1456. [http://doi.org/10.1016/0045-6535\(92\)90168-q](http://doi.org/10.1016/0045-6535(92)90168-q).
- Hagenmaier H, Brunner H, Haag R, et al. 1987. Copper-catalyzed dechlorination/hydrogenation of polychlorinated dibenzo-p-dioxins, polychlorinated dibenzofurans, and other chlorinated aromatic compounds. *Environ Sci Technol* 21(11):1085-1088. <http://doi.org/10.1021/es00164a007>.
- Hagenmaier H, Wiesmuller T, Golor G, et al. 1990. Transfer of various polychlorinated dibenzo-p-dioxins and dibenzofurans (PCDDs and PCDFs) via placenta and through milk in a marmoset monkey. *Arch Toxicol* 64(8):601-615. <http://doi.org/10.1007/BF01974688>.
- Hagenmaier H, Horch K, Fahlenkamp H, et al. 1991. Destruction of PCDD and PCDF in refuse incineration plants by primary and secondary measures. *Chemosphere* 23(8-10):1429-1437. [http://doi.org/10.1016/0045-6535\(91\)90167-c](http://doi.org/10.1016/0045-6535(91)90167-c).
- Hahn JL, Von Dem Fange HP, Westerman G. 1989. A comparison of ambient and workplace dioxin levels from testing in and around modern resource recovery facilities with predicted ground level concentrations of dioxins from stack emission testing with corresponding workplace health risks. *Chemosphere* 19(1-6):629-636. [http://doi.org/10.1016/0045-6535\(89\)90382-2](http://doi.org/10.1016/0045-6535(89)90382-2).
- Hardell L, Fredrikson M, Eriksson M, et al. 1995. Adipose tissue concentrations of dioxins and dibenzofurans in patients with malignant lymphoproliferative diseases and in patients without a malignant disease. *Eur J Cancer Prev* 4(3):225-229.
- Hardell L, Lindstrom G, Liljegren G, et al. 1996. Increased concentrations of octachlorodibenzo-p-dioxin in cases with breast cancer - results from a case-control study. *Eur J Cancer Prev* 5(5):351-357. <http://doi.org/10.1097/00008469-199610000-00007>.
- Hardell L, Lindstrom G, van Bavel B, et al. 2001. Adipose tissue concentrations of dioxins and dibenzofurans, titers of antibodies to Epstein-Barr virus early antigen and the risk for non-Hodgkin lymphoma. *Environ Res* 87(2):99-107. <http://doi.org/10.1006/enrs.2001.4295>.
- Harper N, Connor K, Safe S. 1993. Immunotoxic potencies of polychlorinated biphenyl (PCB), dibenzofuran (PCDF) and dibenzo-p-dioxin (PCDD) congeners in C57BL/6 and DBA/2 mice. *Toxicology* 80(2-3):217-227. [http://doi.org/10.1016/0300-483x\(93\)90183-s](http://doi.org/10.1016/0300-483x(93)90183-s).
- Harrad SJ, Fernandes AR, Creaser CS, et al. 1991a. Domestic coal combustion as a source of PCDDs and PCDFs in the British environment. *Chemosphere* 23(3):255-261. [http://doi.org/10.1016/0045-6535\(91\)90181-c](http://doi.org/10.1016/0045-6535(91)90181-c).
- Harrad SJ, Malloy TA, Khan MA, et al. 1991b. Levels and sources of PCDDs, PCDFs, chlorophenols (CPs) and chlorobenzenes (CBzs) in composts from a municipal yard waste composting facility. *Chemosphere* 23(2):181-191. [http://doi.org/10.1016/0045-6535\(91\)90105-m](http://doi.org/10.1016/0045-6535(91)90105-m).

8. REFERENCES

- Hassoun EA. 1987. In vivo and in vitro interactions of TCDD and other ligands of the Ah-receptor: effect on embryonic and fetal tissues. *Arch Toxicol* 61(2):145-149. <http://doi.org/10.1007/bf00661373>.
- Hassoun E, d'Argy R, Dencker L, et al. 1984. Teratogenicity of 2,3,7,8-tetrachlorodibenzofuran in BXD recombinant inbred strains. *Toxicol Lett* 23(1):37-42. [http://doi.org/10.1016/0378-4274\(84\)90007-9](http://doi.org/10.1016/0378-4274(84)90007-9).
- Hassoun EA, Li F, Abushaban A, et al. 2000. The relative abilities of TCDD and its congeners to induce oxidative stress in the hepatic and brain tissues of rats after subchronic exposure. *Toxicology* 145(2-3):103-113. [http://doi.org/10.1016/s0300-483x\(99\)00221-8](http://doi.org/10.1016/s0300-483x(99)00221-8).
- Hassoun EA, Wang H, Abushaban A, et al. 2002. Induction of oxidative stress in the tissues of rats after chronic exposure to TCDD, 2,3,4,7,8-pentachlorodibenzofuran, and 3,3',4,4',5-pentachlorobiphenyl. *J Toxicol Environ Health* 65(12):825-842. <http://doi.org/10.1080/00984100290071054>.
- Hattemer-Frey HA, Travis CC. 1989. Comparison of human exposure to dioxin from municipal waste incineration and background environmental contamination. *Chemosphere* 18(1-6):643-649. [http://doi.org/10.1016/0045-6535\(89\)90177-x](http://doi.org/10.1016/0045-6535(89)90177-x).
- Haws LC, Su SH, Harris M, et al. 2006. Development of a refined database of mammalian relative potency estimates for dioxin-like compounds. *Toxicol Sci* 89(1):4-30. <http://doi.org/10.1093/toxsci/kfi294>.
- Hayabuchi H, Yoshimura T, Kuratsune M. 1979. Consumption of toxic rice oil by 'Yusho' patients and its relation to the clinical response and latent period. *Food Cosmet Toxicol* 17(5):455-461. [http://doi.org/10.1016/0015-6264\(79\)90004-x](http://doi.org/10.1016/0015-6264(79)90004-x).
- Hayward DG, Charles JM, Voss de Bettancourt C, et al. 1989. PCDD and PCDF in breast milk as correlated with fish consumption in Southern California. *Chemosphere* 18(1-6):455-468. [http://doi.org/10.1016/0045-6535\(89\)90155-0](http://doi.org/10.1016/0045-6535(89)90155-0).
- Hayward DG, Petreas MX, Goldman LR, et al. 1991. Assessing the risks from 2,3,7,8 TCDD and TCDF in milk packaged in paper. *Chemosphere* 23(11-12):1551-1559. [http://doi.org/10.1016/0045-6535\(91\)90002-u](http://doi.org/10.1016/0045-6535(91)90002-u).
- Hebert CD, Harris MW, Elwell MR, et al. 1990. Relative toxicity and tumor-promoting ability of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD), 2,3,4,7,8-pentachlorodibenzofuran (PCDF), and 1,2,3,4,7,8-hexachlorodibenzofuran (HCDF) in hairless mice. *Toxicol Appl Pharmacol* 102(2):362-377. [http://doi.org/10.1016/0041-008x\(90\)90033-q](http://doi.org/10.1016/0041-008x(90)90033-q).
- Heindl A, Hutzinger O. 1989. Search for industrial sources of PCDD/PCDFs: IV. Phthalocyanine dyes. *Chemosphere* 18(1-6):1207-1211. [http://doi.org/10.1016/0045-6535\(89\)90256-7](http://doi.org/10.1016/0045-6535(89)90256-7).
- Hensley AR, Scott A, Rosenfeld PE, et al. 2007. Attic dust and human blood samples collected near a former wood treatment facility. *Environ Res* 105(2):194-199. <http://doi.org/10.1016/j.envres.2007.03.011>.
- Hodson PV, McWhirter M, Ralph K, et al. 1992. Effects of bleached kraft mill effluent on fish in the St. Maurice River, Quebec. *Environ Toxicol Chem* 11(11):1635-1651.
- Holcomb M, Yao C, Safe S. 1988. Biologic and toxic effects of polychlorinated dibenzo-p-dioxin and dibenzofuran congeners in the guinea pig. Quantitative structure-activity relationships. *Biochem Pharmacol* 37(8):1535-1539. [http://doi.org/10.1016/0006-2952\(88\)90015-9](http://doi.org/10.1016/0006-2952(88)90015-9).
- Hood A, Klaassen CD. 2000. Effects of microsomal enzyme inducers on outer-ring deiodinase activity toward thyroid hormones in various rat tissues. *Toxicol Appl Pharmacol* 163(3):240-248. <http://doi.org/10.1006/taap.1999.8883>.
- Hood A, Allen ML, Liu Y, et al. 2003. Induction of T4 UDP-GT activity, serum thyroid stimulating hormone, and thyroid follicular cell proliferation in mice treated with microsomal enzyme inducers. *Toxicol Appl Pharmacol* 188(1):6-13. [http://doi.org/10.1016/s0041-008x\(02\)00071-6](http://doi.org/10.1016/s0041-008x(02)00071-6).
- Hryhorczuk DO, Orris P, Kominsky JR, et al. 1986. PCB, PCDF, and PCDD exposure following a transformer fire: Chicago. *Chemosphere* 15(9-12):1297-1303. [http://doi.org/10.1016/0045-6535\(86\)90404-2](http://doi.org/10.1016/0045-6535(86)90404-2).

8. REFERENCES

- Hsu MML, Chang JC, Hsu CC. 1993. Nail changes in PCB poisoning. *Organohalogen Compounds* 13:251-252. <http://dioxin20xx.org/wp-content/uploads/pdfs/1993/93-154.pdf>. April 23, 2020.
- Hsu ST, Ma CI, Hsu SK, et al. 1984. Discovery and epidemiology of PCB poisoning in Taiwan. *Am J Ind Med* 5(1-2):71-79.
- Hsu ST, Ma CI, Hsu SK, et al. 1985. Discovery and epidemiology of PCB poisoning in Taiwan: a four-year followup. *Environ Health Perspect* 59:5-10. <http://doi.org/10.1289/ehp.59-1568088>.
- Hsu PC, Lai TJ, Guo NW, et al. 2005. Serum hormones in boys prenatally exposed to polychlorinated biphenyls and dibenzofurans. *J Toxicol Environ Health A* 68(17-18):1447-1456. <http://doi.org/10.1080/15287390590967360>.
- Hsu PC, Li MC, Lee YC, et al. 2016. Polychlorinated biphenyls and dibenzofurans increased abnormal sperm morphology without alterations in aneuploidy: The Yucheng study. *Chemosphere* 165:294-297. <http://doi.org/10.1016/j.chemosphere.2016.09.035>.
- Huisman M, Eerenstein SE, Koopman-Esseboom C, et al. 1995. Perinatal exposure to polychlorinated biphenyls and dioxins through dietary intake. *Chemosphere* 31(10):4273-4287.
- Humppi T. 1986. Synthesis of polychlorinated phenoxyphenols (PCPP), phenoxyanisoles (PCPA), dibenzo-p-dioxins (PCDD), dibenzofurans (PCDF) and diphenyl ethers (PCDE). *Chemosphere* 15(9-12):2003-2006. [http://doi.org/10.1016/0045-6535\(86\)90502-3](http://doi.org/10.1016/0045-6535(86)90502-3).
- Hunt GT, Maisel BE. 1990. Atmospheric PCDDs/PCDFs in wintertime in a northeastern U.S. urban coastal environment. *Chemosphere* 20(10-12):1455-1462. [http://doi.org/10.1016/0045-6535\(90\)90297-7](http://doi.org/10.1016/0045-6535(90)90297-7).
- Hutzinger O, Fiedler H. 1989. Sources and emissions of PCDD/PCDF. *Chemosphere* 18(1-6):23-32. [http://doi.org/10.1016/0045-6535\(89\)90102-1](http://doi.org/10.1016/0045-6535(89)90102-1).
- Hutzinger O, Safe S, Wentzell BR, et al. 1973. Photochemical degradation of di- and octachlorodibenzofuran. *Environ Health Perspect* 5:267-271. <http://doi.org/10.1289/ehp.7305267>.
- Hutzinger O, Choudhry GG, Chittim BG, et al. 1985a. Formation of polychlorinated dibenzofurans and dioxins during combustion, electrical equipment fires and PCB incineration. *Environ Health Perspect* 60:3-9. <http://doi.org/10.1289/ehp.85603>.
- Hutzinger O, Blumich MJ, Berg MDV, et al. 1985b. Sources and fate of PCDDs and PCDFs: An overview. *Chemosphere* 14:581-560.
- IARC. 1997. IARC Monographs on the evaluation of carcinogenic risks to humans. Lyon, France: International Agency for Research on Cancer. Vol. 69. <https://publications.iarc.fr/87>. June 17, 2020.
- IARC. 2012. 2,3,7,8-Tetrachlorodibenzopara-dioxin, 2,3,4,7,8-pentachlorodibenzofuran, and 3,3',4,4',5-pentachlorobiphenyl. IARC monographs on the evaluation of carcinogenic risks to humans. Volume 110F. Chemical agents and related occupations. Lyon, France: International Agency for Research on Cancer. <http://publications.iarc.fr/123>. April 20, 2020.
- Iida T, Hirakawa H, Matsueda T, et al. 1999a. Polychlorinated dibenzo-p-dioxins and related compounds: correlations of levels in human tissues and in blood. *Chemosphere* 38(12):2767-2774. [http://doi.org/10.1016/s0045-6535\(98\)00478-0](http://doi.org/10.1016/s0045-6535(98)00478-0).
- Iida T, Hirakawa H, Matsueda T, et al. 1999b. Recent trend of polychlorinated dibenzo-p-dioxins and their related compounds in the blood and sebum of Yusho and Yu-Cheng patients. *Chemosphere* 38(5):981-993.
- Iida T, Todaka T, Hirakawa H, et al. 2007. Concentration and distribution of dioxins and related compounds in human tissues. *Chemosphere* 67(9):S263-271. <http://doi.org/10.1016/j.chemosphere.2006.05.107>.
- Imamura T, Kanagawa Y, Matsumoto S, et al. 2007. Relationship between clinical features and blood levels of pentachlorodibenzofuran in patients with Yusho. *Environ Toxicol* 22(2):124-131. <http://doi.org/10.1002/tox.20251>.
- Imamura T, Matsumoto S, Akahane M, et al. 2009. Cutaneous symptoms such as acneform eruption and pigmentation are closely associated with blood levels of 2,3,4,7,8-penta-chlorodibenzofurans in

8. REFERENCES

- Yusho patients, using data mining analysis. *BMC Res Notes* 2:27. <http://doi.org/10.1186/1756-0500-2-27>.
- Ioannou YM, Birnbaum LS, Matthews HB. 1983. Toxicity and distribution of 2,3,7,8-tetrachlorodibenzofuran in male guinea pigs. *J Toxicol Environ Health* 12(4-6):541-553. <http://doi.org/10.1080/15287398309530448>.
- IRIS. 2020. Chlorodibenzofurans. Integrated risk information system. Washington, DC: U.S. Environmental Protection Agency. https://cfpub.epa.gov/ncea/iris_drafts/atoz.cfm. April 20, 2020.
- Iyer R, Aggarwal J, Iken B. 2016. A review of the Texas, USA San Jacinto Superfund site and the deposition of polychlorinated dibenzo-p-dioxins and dibenzofurans in the San Jacinto River and Houston Ship Channel. *Environ Sci Pollut Res Int* 23(23):23321-23338. <http://doi.org/10.1007/s11356-016-7501-8>.
- Jackson JA, Diliberto JJ, Birnbaum LS. 1993. Estimation of octanol-water partition coefficients and correlation with dermal absorption for several polyhalogenated aromatic hydrocarbons. *Fundam Appl Toxicol* 21(3):334-344.
- Jarman WM, Burns SA, Chang RR, et al. 1993. Determination of PCDDS, PCDFs, and PCBs in California peregrine falcons (*Falco peregrinus*) and their eggs. *Environ Toxicol Chem* 12(1):105-114. <http://doi.org/10.1002/etc.5620120112>.
- Jay K, Stieglitz L. 1991. On the mechanism of formation of polychlorinated aromatic compounds with copper(II) chloride. *Chemosphere* 22(11):987-996. [http://doi.org/10.1016/0045-6535\(91\)90299-s](http://doi.org/10.1016/0045-6535(91)90299-s).
- Jensen AA. 1987. Polychlorobiphenyls (PCBs), polychlorodibenzo-p-dioxins (PCDDs) and polychlorodibenzofurans (PCDFs) in human milk, blood and adipose tissue. *Sci Total Environ* 64(3):259-293. [http://doi.org/10.1016/0048-9697\(87\)90250-6](http://doi.org/10.1016/0048-9697(87)90250-6).
- Jödicke B, Enge M, Helge H, et al. 1992. Fecal excretion of PCDDs/PCDFs in a 3-month-old breast-fed infant. *Chemosphere* 25(7-10):1061-1065.
- Johnson KL, Cummings AM, Birnbaum LS. 1997. Promotion of endometriosis in mice by polychlorinated dibenzo-p-dioxins, dibenzofurans, and biphenyls. *Environ Health Perspect* 105(7):750-755. <http://doi.org/10.1289/ehp.97105750>.
- Johnson CW, Williams WC, Copeland CB, et al. 2000. Sensitivity of the SRBC PFC assay versus ELISA for detection of immunosuppression by TCDD and TCDD-like congeners. *Toxicology* 156(1):1-11. [http://doi.org/10.1016/s0300-483x\(00\)00330-9](http://doi.org/10.1016/s0300-483x(00)00330-9).
- Jones PD, Giesy JP, Newsted JL, et al. 1993. 2,3,7,8-Tetrachlorodibenzo-p-dioxin equivalents in tissues of birds at Green Bay, Wisconsin, USA. *Arch Environ Contam Toxicol* 24(3):345-354. <http://doi.org/10.1007/bf01128732>.
- Jordan RJ. 1987. The feasibility of wet scrubbing for treating waste-to-energy flue gas. *J Air Pollut Control Assoc* 37(4):422-430. <http://doi.org/10.1080/08940630.1987.10466237>.
- Kanagawa Y, Matsumoto S, Koike S, et al. 2008. Association of clinical findings in Yusho patients with serum concentrations of polychlorinated biphenyls, polychlorinated quarterphenyls and 2,3,4,7,8-pentachlorodibenzofuran more than 30 years after the poisoning event. *Environ Health* 7:47. <http://doi.org/10.1186/1476-069X-7-47>.
- Kang HK, Watanabe KK, Breen J, et al. 1991. Dioxins and dibenzofurans in adipose tissue of US Vietnam veterans and controls. *Am J Public Health* 81(3):344-349. <http://doi.org/10.2105/ajph.81.3.344>.
- Kashimoto T, Miyata H. 1986. Differences between Yusho and other kinds of poisoning involving only PCBs. In: Waid JS, ed. *PCBs and the environment*. Vol. 3. Boca Raton, FL: CRC Press, 1-26.
- Kawano S, Hiraga K. 1978. Polychlorinated dibenzofurans - potent inducers of rat hepatic drug-metabolizing enzymes. *Jpn J Pharmacol* 28(2):305-315. <http://doi.org/10.1254/jjp.28.305>.
- Keenan RE, Sullivan MJ. 1989. Assessing potential health risks of dioxin in paper products. *Environ Sci Technol* 23(6):643-644. <http://doi.org/10.1021/es00064a002>.
- Kelada FS. 1990. Occupational intake by dermal exposure to polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans in pulp mill industry. *Am Ind Hyg Assoc J* 51(9):519-521. <http://doi.org/10.1080/15298669091370022>.

8. REFERENCES

- Kerger BD, Leung HW, Scott PK, et al. 2007a. An adaptable internal dose model for risk assessment of dietary and soil dioxin exposures in young children. *Toxicol Sci* 100(1):224-237. <http://doi.org/10.1093/toxsci/kfm199>.
- Kerger BD, Leung HW, Scott PK, et al. 2007b. Refinements on the age-dependent half-life model for estimating child body burdens of polychlorodibenzodioxins and dibenzofurans. *Chemosphere* 67(9):S272-278. <http://doi.org/10.1016/j.chemosphere.2006.05.108>.
- Kerkvliet NI, Brauner JA, Matlock JP. 1985. Humoral immunotoxicity of polychlorinated diphenyl ethers, phenoxyphenols, dioxins and furans present as contaminants of technical grade pentachlorophenol. *Toxicology* 36(4):307-324. [http://doi.org/10.1016/0300-483x\(85\)90033-2](http://doi.org/10.1016/0300-483x(85)90033-2).
- Keys B, Piskorska-Pliszczynska J, Safe S. 1986. Polychlorinated dibenzofurans as 2,3,7,8-TCDD antagonists: in vitro inhibition of monooxygenase enzyme induction. *Toxicol Lett* 31(2):151-158. [http://doi.org/10.1016/0378-4274\(86\)90009-3](http://doi.org/10.1016/0378-4274(86)90009-3).
- Khairy MA, Noonan GO, Lohmann R. 2019. Uptake of hydrophobic organic compounds, including organochlorine pesticides, polybrominated diphenyl ethers, and perfluoroalkyl acids in fish and blue crabs of the lower Passaic River, New Jersey, USA. *Environ Toxicol Chem* 38(4):872-882. <http://doi.org/10.1002/etc.4354>.
- Khera KS. 1992. Extraembryonic tissue changes induced by 2,3,7,8-tetrachlorodibenzo-p-dioxin and 2,3,4,7,8-pentachlorodibenzofuran with a note on direction of maternal blood flow in the labyrinth of C57BL/6N mice. *Teratology* 45(6):611-627. <http://doi.org/10.1002/tera.1420450606>.
- Kitunen VH, Salkinoja-Salonen MS. 1989. Occurrence of PCDDs and PCDFs in pulp and board products. *Chemosphere* 19(1-6):721-726. [http://doi.org/10.1016/0045-6535\(89\)90397-4](http://doi.org/10.1016/0045-6535(89)90397-4).
- Kitunen VH, Valo RJ, Salkinoja-Salonen MS. 1987. Contamination of soil around wood-preserving facilities by polychlorinated aromatic compounds. *Environ Sci Technol* 21(1):96-101. <http://doi.org/10.1021/es00155a012>.
- Kjeller LO, Jones KC, Johnston AE, et al. 1991. Increases in the polychlorinated dibenzo-p-dioxin and -furan content of soils and vegetation since the 1840s. *Environ Sci Technol* 25(9):1619-1627. <http://doi.org/10.1021/es00021a014>.
- Koester CJ, Hites RA. 1992. Photodegradation of polychlorinated dioxins and dibenzofurans adsorbed to fly ash. *Environ Sci Technol* 26(3):502-507. <http://doi.org/10.1021/es00027a008>.
- Kondo H, Tanio K, Nagaura Y, et al. 2018. Sleep disorders among Yusho patients highly intoxicated with dioxin-related compounds: A 140-case series. *Environ Res* 166:261-268. <http://doi.org/10.1016/j.envres.2018.05.033>.
- Konishi K, Sasaki S, Kato S, et al. 2009. Prenatal exposure to PCDDs/PCDFs and dioxin-like PCBs in relation to birth weight. *Environ Res* 109(7):906-913. <http://doi.org/10.1016/j.envres.2009.07.010>.
- Körner W, Dawidowsky N, Hagenmaier H. 1993. Fecal excretion rates of PCDDs and PCDFs in two breast-fed infants. *Chemosphere* 27(1-3):157-162.
- Koshioka M, Iizuka H, Kanazawa J, et al. 2014. Photodegradation of octachlorodi-benzofuran in 1,4-dioxane under xenon lamp irradiation. *Agric Biol Chem* 51(3):949-952. <http://doi.org/10.1080/00021369.1987.10868130>.
- Krowke R. 1986. Studies on distribution and embryotoxicity of different PCDD and PCDF in mice and marmosets. *Chemosphere* 15(9-12):2011-2022. [http://doi.org/10.1016/0045-6535\(86\)90504-7](http://doi.org/10.1016/0045-6535(86)90504-7).
- Kuehl DW, Haebler R, Potter C. 1991. Chemical residues in dolphins from the U.S. Atlantic coast including Atlantic bottlenose obtained during the 1987/88 mass mortality. *Chemosphere* 22(11):1071-1084. [http://doi.org/10.1016/0045-6535\(91\)90308-z](http://doi.org/10.1016/0045-6535(91)90308-z).
- Kuehl DW, Cook PM, Batterman AR, et al. 1987. Bioavailability of polychlorinated dibenzo-p-dioxins and dibenzofurans from contaminated Wisconsin River sediment to carp. *Chemosphere* 16(4):667-679. [http://doi.org/10.1016/0045-6535\(87\)90003-8](http://doi.org/10.1016/0045-6535(87)90003-8).
- Kunita N, Kashimoto T, Miyata H, et al. 1984. Causal agents of Yusho. *Am J Ind Med* 5(1-2):45-58.
- Kurachi M, Hashimoto S, Obata A, et al. 2002. Identification of 2,3,7,8-tetrachlorodibenzo-p-dioxin-responsive genes in mouse liver by serial analysis of gene expression. *Biochem Biophys Res Commun* 292(2):368-377. <http://doi.org/10.1006/bbrc.2002.6669>.

8. REFERENCES

- Kuratsune M. 1989. Yusho, with reference to Yu-cheng. In: Kimbrough RD, Jensen AA, eds. Halogenated biphenyls, terphenyls, naphthalene, dibenzodioxins and related products. Vol. 4. 2nd ed. Amsterdam: Elsevier Science Publishers, 381-400.
- Kuratsune M, Nakamura Y, Ikeda M, et al. 1987. Analysis of deaths seen among patients with Yusho. *Chemosphere* 16(8-9):2085-2088. [http://doi.org/10.1016/0045-6535\(87\)90212-8](http://doi.org/10.1016/0045-6535(87)90212-8).
- Kuroiwa Y, Murai YST. 1969. [Neurological and nerve conduction velocity studies on 23 patients with chlorobiphenyls poisoning]. *Fukuoka Igaku Zasshi* 60:462-463. (Japanese)
- Kuroki H, Haraguchi K, Masuda Y. 1984. Synthesis of polychlorinated dibenzofuran isomers and their gas chromatographic profiles. *Chemosphere* 13(4):561-573. [http://doi.org/10.1016/0045-6535\(84\)90151-6](http://doi.org/10.1016/0045-6535(84)90151-6).
- Kusuda M. 1971. [A study on the sexual functions of women suffering from rice-bran oil poisoning]. *Sanka to Fujinka* 38:1063-1072. (Japanese)
- Kuwatsuka Y, Shimizu K, Akiyama Y, et al. 2014. Yusho patients show increased serum IL-17, IL-23, IL-1beta, and TNFalpha levels more than 40 years after accidental polychlorinated biphenyl poisoning. *J Immunotoxicol* 11(3):246-249. <http://doi.org/10.3109/1547691X.2013.835890>.
- LaKind JS, Hays SM, Aylward LL, et al. 2009. Perspective on serum dioxin levels in the United States: an evaluation of the NHANES data. *J Expo Sci Environ Epidemiol* 19(4):435-441. <http://doi.org/10.1038/jes.2008.63>.
- Lan SJ, Yen YY, Yang CH, et al. 1987. [A study on the birth weight of transplacental Yu-cheng babies]. *Gaoxiong Yi Xue Ke Xue Za Zhi* 3(4):273-282. (Chinese)
- Lan SJ, Yen YY, Lan JL, et al. 1990. Immunity of PCB transplacental Yu-cheng children in Taiwan. *Bull Environ Contam Toxicol* 44(2):224-229. <http://doi.org/10.1007/bf01700140>.
- LeBel GL, Williams DT, Benoit FM. 1992. Chlorinated dibenzodioxins and dibenzofurans in consumer paper products. *Chemosphere* 25(11):1683-1690. [http://doi.org/10.1016/0045-6535\(92\)90315-i](http://doi.org/10.1016/0045-6535(92)90315-i).
- LeBel GL, Williams DT, Benoit FM, et al. 1990. Polychlorinated dibenzodioxins and dibenzofurans in human adipose tissue samples from five Ontario municipalities. *Chemosphere* 21(12):1465-1475. [http://doi.org/10.1016/0045-6535\(90\)90052-u](http://doi.org/10.1016/0045-6535(90)90052-u).
- Leung HW, Kerger BD, Paustenbach DJ. 2006. Elimination half-lives of selected polychlorinated dibenzodioxins and dibenzofurans in breast-fed human infants. *J Toxicol Environ Health* 69(6):437-443. <http://doi.org/10.1080/15287390500246886>.
- Leung HW, Kerger BD, Paustenbach D, et al. 2005. Age- and concentration-dependent elimination half lives of chlorinated dibenzofurans in Yusho and Yucheng patients. *Organohalogen Compounds* 67:1589-1592.
- Leung HW, Kerger BD, Paustenbach DJ, et al. 2007. Concentration and age-dependent elimination kinetics of polychlorinated dibenzofurans in Yucheng and Yusho patients. *Toxicol Ind Health* 23(8):493-501. <http://doi.org/10.1177/0748233708089024>.
- Li MC, Tsai PC, Chen PC, et al. 2013. Mortality after exposure to polychlorinated biphenyls and dibenzofurans: 30 years after the "Yucheng accident". *Environ Res* 120:71-75. <http://doi.org/10.1016/j.envres.2012.09.003>.
- Li MC, Chen PC, Tsai PC, et al. 2015a. Mortality after exposure to polychlorinated biphenyls and polychlorinated dibenzofurans: a meta-analysis of two highly exposed cohorts. *Int J Cancer* 137(6):1427-1432. <http://doi.org/10.1002/ijc.29504>.
- Li MC, Wu HP, Yang CY, et al. 2015b. Gestational exposure to polychlorinated biphenyls and dibenzofurans induced asymmetric hearing loss: Yucheng children study. *Environ Res* 137:65-71. <http://doi.org/10.1016/j.envres.2014.12.002>.
- Ligon WV, Dorn SB, May RJ, et al. 1989. Chlorodibenzofuran and chlorodibenzo-p-dioxin levels in Chilean mummies dated to about 2800 years before the present. *Environ Sci Technol* 23(10):1286-1290. <http://doi.org/10.1021/es00068a016>.
- Lin KC, Guo NW, Tsai PC, et al. 2008. Neurocognitive changes among elderly exposed to PCBs/PCDFs in Taiwan. *Environ Health Perspect* 116(2):184-189. <http://doi.org/10.1289/ehp.10134>.

8. REFERENCES

- Lin KC, Huang PC, Yeh PS, et al. 2010. Comparing mini-mental state examination and attention and digit span in elderly exposed to polychlorinated biphenyls and polychlorinated dibenzofurans. *Psychogeriatrics* 10(4):191-197. <http://doi.org/10.1111/j.1479-8301.2010.00341.x>.
- Lindner G, Jenkins AC, McCormack J, et al. 1990. Dioxins and furans in emissions from medical waste incinerators. *Chemosphere* 20(10-12):1793-1800. [http://doi.org/10.1016/0045-6535\(90\)90344-s](http://doi.org/10.1016/0045-6535(90)90344-s).
- Lofroth G, Zebuhr Y. 1992. Polychlorinated dibenzo-p-dioxins (PCDDs) and dibenzofurans (PCDFs) in mainstream and sidestream cigarette smoke. *Bull Environ Contam Toxicol* 48(6):789-794. <http://doi.org/10.1007/bf00201136>.
- Lu YC, Wu YC. 1985. Clinical findings and immunological abnormalities in Yu-Cheng patients. *Environ Health Perspect* 59:17-29. <http://doi.org/10.1289/ehp.59-1568085>.
- Lu FJ, Chang KJ, Lin SC, et al. 1980. [Studies on patients with polychlorinated biphenyls poisoning. Determination of urinary coproporphyrin, uroporphyrin, delta-aminolevulinic acid, and porphobilinogen (author's transl)]. *Taiwan Yi Xue Hui Za Zhi* 79(11):990-995. (Chinese)
- Lucier GW, Sunahara GI, Wong TK. 1990. Placental markers of human exposure to polychlorinated dibenzofurans and polychlorinated biphenyls: implications for risk assessment. *IARC Sci Publ* 104(104):55-62.
- Lucier GW, Nelson KG, Everson RB, et al. 1987. Placental markers of human exposure to polychlorinated biphenyls and polychlorinated dibenzofurans. *Environ Health Perspect* 76:79-87. <http://doi.org/10.1289/ehp.877679>.
- Lundgren K, Collman GW, Wang-Wuu S, et al. 1988. Cytogenetic and chemical detection of human exposure to polyhalogenated aromatic hydrocarbons. *Environ Mol Mutagen* 11(1):1-11. <http://doi.org/10.1002/em.2850110103>.
- Luster MI, Faith RE, Clark G. 1979a. Laboratory studies on the immune effects of halogenated aromatics. *Ann N Y Acad Sci* 320:473-486. <http://doi.org/10.1111/j.1749-6632.1979.tb56628.x>.
- Luster MI, Faith RE, Lawson LD. 1979b. Effects of 2,3,7,8-tetrachlorodibenzofuran (TCDF) on the immune system in guinea pigs. *Drug Chem Toxicol* 2(1-2):49-60. <http://doi.org/10.3109/01480547908993181>.
- Madsen C, Larsen JC. 1989. Relative toxicity of chlorinated dibenzo-p-dioxins, and dibenzofurans measured by thymus weight and liver enzyme induction in perinatally dosed rats, 2,3,7,8-TCDD, 2,3,4,7,8-PeCDF, 1,2,3,7,8-PeCDD. *Chemosphere* 18(1-6):955-966. [http://doi.org/10.1016/0045-6535\(89\)90223-3](http://doi.org/10.1016/0045-6535(89)90223-3).
- Maisel BE, Hunt GT. 1990. Background concentrations of PCDDs/PCDFs in ambient air - A comparison of toxic equivalency factor (TEF) models. *Chemosphere* 20(7-9):771-778. [http://doi.org/10.1016/0045-6535\(90\)90181-r](http://doi.org/10.1016/0045-6535(90)90181-r).
- Mäntykoski K, Paasivirta J, Mannila E. 1989. Combustion products of biosludge from pulp mill. *Chemosphere* 19(1-6):413-416. [http://doi.org/10.1016/0045-6535\(89\)90344-5](http://doi.org/10.1016/0045-6535(89)90344-5).
- Marklund S, Rappe C, Tysklind M, et al. 1987. Identification of polychlorinated dibenzofurans and dioxins in exhausts from cars run on leaded gasoline. *Chemosphere* 16(1):29-36. [http://doi.org/10.1016/0045-6535\(87\)90105-6](http://doi.org/10.1016/0045-6535(87)90105-6).
- Marshall NB, Kerkvliet NI. 2010. Dioxin and immune regulation: emerging role of aryl hydrocarbon receptor in the generation of regulatory T cells. *Ann N Y Acad Sci* 1183:25-37. <http://doi.org/10.1111/j.1749-6632.2009.05125.x>.
- Martinez-Zamora MA, Mattioli L, Parera J, et al. 2015. Increased levels of dioxin-like substances in adipose tissue in patients with deep infiltrating endometriosis. *Hum Reprod* 30(5):1059-1068. <http://doi.org/10.1093/humrep/dev026>.
- Maruyama W, Yoshida K, Tanaka T, et al. 2002. Possible range of dioxin concentration in human tissues: simulation with a physiologically based model. *J Toxicol Environ Health A* 65(24):2053-2073. <http://doi.org/10.1080/00984100290071829>.
- Maruyama W, Yoshida K, Tanaka T, et al. 2003. Simulation of dioxin accumulation in human tissues and analysis of reproductive risk. *Chemosphere* 53(4):301-313. [http://doi.org/10.1016/S0045-6535\(03\)00015-8](http://doi.org/10.1016/S0045-6535(03)00015-8).

8. REFERENCES

- Mason G, Sawyer T, Keys B, et al. 1985. Polychlorinated dibenzofurans (PCDFs): Correlation between in vivo and in vitro structure-activity relationships. *Toxicology* 37(1-2):1-12. [http://doi.org/10.1016/0300-483x\(85\)90108-8](http://doi.org/10.1016/0300-483x(85)90108-8).
- Masuda Y. 2001. Fate of PCDF/PCB congeners and change of clinical symptoms in patients with Yusho PCB poisoning for 30 years. *Chemosphere* 43(4-7):925-930. [http://doi.org/10.1016/s0045-6535\(00\)00452-5](http://doi.org/10.1016/s0045-6535(00)00452-5).
- Masuda Y, Yoshimura H. 1984. Polychlorinated biphenyls and dibenzofurans in patients with Yusho and their toxicological significance: a review. *Am J Ind Med* 5(1-2):31-44.
- Masuda Y, Kuroki H, Haraguchi K, et al. 1985. PCB and PCDF congeners in the blood and tissues of Yusho and Yu-cheng patients. *Environ Health Perspect* 59:53-58. <http://doi.org/10.1289/ehp.59-1568081>.
- Matsumoto S, Akahane M, Kanagawa Y, et al. 2009. Variation in half-life of penta-chlorodibenzofuran (PeCDF) blood level among Yusho patients. *Chemosphere* 77(5):658-662. <http://doi.org/10.1016/j.chemosphere.2009.08.009>.
- Matsumoto S, Kanagawa Y, Koike S, et al. 2010. Twenty-year changes of penta-chlorodibenzofuran (PeCDF) level and symptoms in Yusho patients, using association analysis. *BMC Res Notes* 3:129. <http://doi.org/10.1186/1756-0500-3-129>.
- Matsumoto S, Akahane M, Kanagawa Y, et al. 2016. Change in decay rates of dioxin-like compounds in Yusho patients. *Environ Health* 15(1):95. <http://doi.org/10.1186/s12940-016-0178-0>.
- Matsumura F. 2009. The significance of the nongenomic pathway in mediating inflammatory signaling of the dioxin-activated Ah receptor to cause toxic effects. *Biochem Pharmacol* 77(4):608-626. <http://doi.org/10.1016/j.bcp.2008.10.013>.
- McConnell EE. 1989. Acute and chronic toxicity and carcinogenesis in animals. In: Kimbrough RD, Jensen AA, eds. *Halogenated biphenyls, terphenyls, naphthalenes, dibenzodioxins and related products*. Vol. 4. 2nd ed. Amsterdam: Elsevier Science Publishers, 161-193.
- McKee P, Burt A, McCurvin D, et al. 1990. Levels of dioxins, furans and other organic contaminants in harbour sediments near a wood preserving plant using pentachlorophenol and creosote. *Chemosphere* 20(10-12):1679-1685. [http://doi.org/10.1016/0045-6535\(90\)90329-r](http://doi.org/10.1016/0045-6535(90)90329-r).
- McLachlan MS. 1993. Digestive tract absorption of polychlorinated dibenzo-p-dioxins, dibenzofurans, and biphenyls in a nursing infant. *Toxicol Appl Pharmacol* 123(1):68-72. <http://doi.org/10.1006/taap.1993.1222>.
- McLaughlin DL, Pearson RG, Clement RE. 1989. Concentrations of chlorinated dibenzo-p-dioxins (CDD) and dibenzofurans (CDF) in soil from the vicinity of a large refuse incinerator in Hamilton Ontario. *Chemosphere* 18(1-6):851-854. [http://doi.org/10.1016/0045-6535\(89\)90208-7](http://doi.org/10.1016/0045-6535(89)90208-7).
- McNulty WP, Pomerantz I, Farrell T. 1981. Chronic toxicity of 2,3,7,8-tetrachlorodibenzofuran for rhesus macaques. *Food Cosmet Toxicol* 19(1):57-65. [http://doi.org/10.1016/0015-6264\(81\)90304-7](http://doi.org/10.1016/0015-6264(81)90304-7).
- Meyer C, O'Keefe P, Hilker D, et al. 1989. A survey of twenty community water systems in New York State for PCDDs and PCDFs. *Chemosphere* 19(1-6):21-26. [http://doi.org/10.1016/0045-6535\(89\)90284-1](http://doi.org/10.1016/0045-6535(89)90284-1).
- Milbrath MO, Wenger Y, Chang CW, et al. 2009. Apparent half-lives of dioxins, furans, and polychlorinated biphenyls as a function of age, body fat, smoking status, and breast-feeding. *Environ Health Perspect* 117(3):417-425. <http://doi.org/10.1289/ehp.11781>.
- Mitoma C, Mine Y, Utani A, et al. 2015. Current skin symptoms of Yusho patients exposed to high levels of 2,3,4,7,8-pentachlorinated dibenzofuran and polychlorinated biphenyls in 1968. *Chemosphere* 137:45-51. <http://doi.org/10.1016/j.chemosphere.2015.03.070>.
- Miyata H, Takayama K, Ogaki J, et al. 1989. Levels of PCDDs, coplanar PCBs and PCDFs in patients with Yusho disease and in the Yusho oil. *Chemosphere* 18(1-6):407-416. [http://doi.org/10.1016/0045-6535\(89\)90149-5](http://doi.org/10.1016/0045-6535(89)90149-5).

8. REFERENCES

- Moore JA, Gupta BN, Vos JG. 1976. Toxicity of 2,3,7,8-tetrachlorodibenzofuran-preliminary results. In: National conference on polychlorinated biphenyls. Washington, DC: U.S. Environmental Protection Agency, 77-80.
- Moore JA, McConnell EE, Dalgard DW, et al. 1979. Comparative toxicity of three halogenated dibenzofurans in guinea pigs, mice, and rhesus monkeys. *Ann N Y Acad Sci* 320:151-163. <http://doi.org/10.1111/j.1749-6632.1979.tb56598.x>.
- Morales L, Dachs J, Gonzalez-Gaya B, et al. 2014. Background concentrations of polychlorinated dibenzo-p-dioxins, dibenzofurans, and biphenyls in the global oceanic atmosphere. *Environ Sci Technol* 48(17):10198-10207. <http://doi.org/10.1021/es5023619>.
- Morrissey RE, Schwetz BA. 1989. Reproductive and developmental toxicity in animals. In: Kimbrough RD, Jensen AA, eds. Halogenated biphenyls, terphenyls, naphthalene, dibenzodioxins and related products. 2nd ed. Amsterdam, Netherlands: Elsevier Science Publishers, 195-225.
- Moser GA, McLachlan MS. 2001. The influence of dietary concentration on the absorption and excretion of persistent lipophilic organic pollutants in the human intestinal tract. *Chemosphere* 45(2):201-211. [http://doi.org/10.1016/s0045-6535\(00\)00551-8](http://doi.org/10.1016/s0045-6535(00)00551-8).
- Muir DCG, Lawrence S, Holoka M, et al. 1992. Partitioning of polychlorinated dioxins and furans between water, sediments and biota in lake mesocosms. *Chemosphere* 25(1-2):119-124. [http://doi.org/10.1016/0045-6535\(92\)90494-c](http://doi.org/10.1016/0045-6535(92)90494-c).
- Mukerjee D, Papke O, Karmaus W. 1989. Indoor air contamination with polychlorinated dibenzo-p-dioxins and dibenzofurans. *Toxicol Ind Health* 5(5):731-745. <http://doi.org/10.1177/074823378900500511>.
- Mundy KJ, Brown RS, Pettit K, et al. 1989. Environmental assessment at and around a chemical waste treatment facility I - Measurements on PCDFs and PCDDs. *Chemosphere* 19(1-6):381-386. [http://doi.org/10.1016/0045-6535\(89\)90339-1](http://doi.org/10.1016/0045-6535(89)90339-1).
- Muto H, Takizawa Y. 1991. The liquid-phase photolyses of tetra- and octa-CDDs and their CDFs in hexane solution. *Chem Lett (Jpn)* 20(2):273-276. <http://doi.org/10.1246/cl.1991.273>.
- Muto H, Saito K, Shinada M, et al. 1991. Concentrations of polychlorinated dibenzo-p-dioxins and dibenzofurans from chemical manufacturers and waste disposal facilities. *Environ Res* 54(2):170-182. [http://doi.org/10.1016/s0013-9351\(05\)80099-4](http://doi.org/10.1016/s0013-9351(05)80099-4).
- Nadal M, Mari M, Schuhmacher M, et al. 2019. Monitoring dioxins and furans in plasma of individuals living near a hazardous waste incinerator: Temporal trend after 20 years. *Environ Res* 173:207-211. <http://doi.org/10.1016/j.envres.2019.03.051>.
- Naf C, Broman D, Pettersen H, et al. 1992. Flux estimates and pattern recognition of particulate polycyclic aromatic hydrocarbons, polychlorinated dibenzo-p-dioxins, and dibenzofurans in the waters outside various emission sources on the Swedish Baltic coast. *Environ Sci Technol* 26(7):1444-1457. <http://doi.org/10.1021/es00031a024>.
- Nagai J, Furukawa M, Tojo A, et al. 1971. [Coulometric and gas-chromatographic determinations of urinary 17-ketosteroids: Survey of chlorobiphenyls poisoning patients by these methods]. *Fukuoka Igaku Zasshi* 62:51-65. (Japanese)
- Nagao T, Golor G, Hagenmaier H, et al. 1993. Teratogenic potency of 2,3,4,7,8-pentachlorodibenzofuran and of three mixtures of polychlorinated dibenzo-p-dioxins and dibenzofurans in mice. Problems with risk assessment using TCDD toxic-equivalency factors. *Arch Toxicol* 67(9):591-597.
- Nagayama J, Masuda Y, Kuratsune M. 1977. Determination of polychlorinated dibenzofurans in tissues of patients with 'Yusho'. *Food Cosmet Toxicol* 15(3):195-198. [http://doi.org/10.1016/s0015-6264\(77\)80389-1](http://doi.org/10.1016/s0015-6264(77)80389-1).
- Nagayama J, Tsuji H, Iida T, et al. 2001. Effects of contamination level of dioxins and related chemicals on thyroid hormone and immune response systems in patients with "Yusho". *Chemosphere* 43(4-7):1005-1010.

8. REFERENCES

- Nagayama J, Kohno H, Kunisue T, et al. 2007. Concentrations of organochlorine pollutants in mothers who gave birth to neonates with congenital hypothyroidism. *Chemosphere* 68(5):972-976. <http://doi.org/10.1016/j.chemosphere.2007.01.010>.
- Nakajima S, Saijo Y, Kato S, et al. 2006. Effects of prenatal exposure to polychlorinated biphenyls and dioxins on mental and motor development in Japanese children at 6 months of age. *Environ Health Perspect* 114(5):773-778. <http://doi.org/10.1289/ehp.8614>.
- Nakajima S, Saijo Y, Miyashita C, et al. 2017. Sex-specific differences in effect of prenatal exposure to dioxin-like compounds on neurodevelopment in Japanese children: Sapporo cohort study. *Environ Res* 159:222-231. <http://doi.org/10.1016/j.envres.2017.08.006>.
- Nakanishi Y, Shigematsu N, Kurita Y, et al. 1985. Respiratory involvement and immune status in Yusho patients. *Environ Health Perspect* 59:31-36. <http://doi.org/10.1289/ehp.59-1568074>.
- Nakano T, Tsuji M, Okuno T. 1990. Distribution of PCDDs, PCDFs and PCBs in the atmosphere. *Atmos Environ* 24(6):1361-1368. [http://doi.org/10.1016/0960-1686\(90\)90042-1](http://doi.org/10.1016/0960-1686(90)90042-1).
- Nakano S, Noguchi T, Takekoshi H, et al. 2005. Maternal-fetal distribution and transfer of dioxins in pregnant women in Japan, and attempts to reduce maternal transfer with *Chlorella* (*Chlorella pyrenoidosa*) supplements. *Chemosphere* 61(9):1244-1255. <http://doi.org/10.1016/j.chemosphere.2005.03.080>.
- Nakao T, Aozasa O, Ohta S, et al. 2002. Assessment of human exposure to PCDDs, PCDFs and Co-PCBs using hair as a human pollution indicator sample I: Development of analytical method for human hair and evaluation for exposure assessment. *Chemosphere* 48(8):885-896. [http://doi.org/10.1016/s0045-6535\(02\)00156-x](http://doi.org/10.1016/s0045-6535(02)00156-x).
- Narang AS, Swami K, Narang RS, et al. 1991. Pyrolysis and combustion of liquids and solids containing pentachlorophenol. *Chemosphere* 22(11):1029-1043. [http://doi.org/10.1016/0045-6535\(91\)90304-v](http://doi.org/10.1016/0045-6535(91)90304-v).
- NAS/NRC. 1989. Report of the oversight committee. In: *Biologic markers in reproductive toxicology*. Washington, DC: National Academy of Sciences, National Research Council, National Academy Press, 15-35.
- Neamah WH, Busbee PB, Alghetaa H, et al. 2020. AhR Activation leads to alterations in the gut microbiome with consequent effect on induction of myeloid derived suppressor cells in a CXCR2-dependent manner. *International Journal of Molecular Sciences* 21(24) <http://doi.org/10.3390/ijms21249613>.
- Nebert DW, Robinson JR, Niwa A, et al. 1975. Genetic expression of aryl hydrocarbon hydroxylase activity in the mouse. *Arch Biochem Biophys* 85(2 Pt 2 Suppl 1):393-414. <http://doi.org/10.1002/jcp.1040850407>.
- Needham LL, Patterson DG, Alley CC, et al. 1987. Polychlorinated dibenzo-p-dioxins and dibenzofurans levels in persons with high and normal levels of 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Chemosphere* 16(8-9):2027-2031. [http://doi.org/10.1016/0045-6535\(87\)90203-7](http://doi.org/10.1016/0045-6535(87)90203-7).
- Neubert D, Wiesmuller T, Abraham K, et al. 1990. Persistence of various polychlorinated dibenzo-p-dioxins and dibenzofurans (PCDDs and PCDFs) in hepatic and adipose tissue of marmoset monkeys. *Arch Toxicol* 64(6):431-442. <http://doi.org/10.1007/BF01977624>.
- Nguyen ATN, Nishijo M, Pham TT, et al. 2018. Sex-specific effects of perinatal dioxin exposure on eating behavior in 3-year-old Vietnamese children. *BMC Pediatr* 18(1):213. <http://doi.org/10.1186/s12887-018-1171-2>.
- NIOSH. 2018. NIOSH pocket guide to chemical hazards. Index of CAS numbers. Atlanta, GA: National Institute for Occupational Safety and Health. <https://www.cdc.gov/niosh/npg/npgdcas.html>. April 20, 2020.
- Nishijo M, Tai PT, Nui NM, et al. 2012. Effects of dioxins exposure on social emotional behavior of children living in a hot spot area, Vietnam. *Organohalogen Compounds* 74:1332-1335. <http://dioxin20xx.org/wp-content/uploads/pdfs/2012/1341.pdf>. October 7, 2019.

8. REFERENCES

- Norstrom RJ, Simon M, Muir DCG. 1990. Polychlorinated dibenzo-p-dioxins and dibenzofurans in marine mammals in the Canadian North. *Environ Pollut* 66(1):1-19. [http://doi.org/10.1016/0269-7491\(90\)90195-i](http://doi.org/10.1016/0269-7491(90)90195-i).
- Norström Å, Chaudhary S, Albro PW, et al. 1979. Synthesis of chlorinated dibenzofurans and chlorinated amino-dibenzofurans from the corresponding diphenyl ethers and nitro-diphenylethers. *Chemosphere* 8(6):331-343. [http://doi.org/10.1016/0045-6535\(79\)90121-8](http://doi.org/10.1016/0045-6535(79)90121-8).
- Norwood CB, Hackett M, Pruell RJ, et al. 1989. Polychlorinated dibenzo-p-dioxins and dibenzofurans in selected estuarine sediments. *Chemosphere* 18(1-6):553-560. [http://doi.org/10.1016/0045-6535\(89\)90166-5](http://doi.org/10.1016/0045-6535(89)90166-5).
- NTP. 2006. Toxicology and carcinogenesis studies of 2,3,4,7,8-pentachlorodibenzofuran (PeCDF) (Cas No. 57117-31-4) in female Harlan Sprague-Dawley rats (gavage studies). National Toxicology Program. TR525. https://ntp.niehs.nih.gov/ntp/htdocs/lt_rpts/tr526.pdf. June 19, 2020.
- NTP. 2016. CASRN index. In: Report on carcinogens. 14th ed. Research Triangle Park, NC: National Toxicology Program, <https://ntp.niehs.nih.gov/pubhealth/roc/index-1.html#P>. March 1, 2017.
- Nygren M, Hansson M, Sjöström M, et al. 1988. Development and validation of a method for determination of PCDDs and PCDFs in human blood plasma. A multivariate comparison of blood and adipose tissue levels between Viet Nam veterans and matched controls. *Chemosphere* 17(9):1663-1692. [http://doi.org/10.1016/0045-6535\(88\)90095-1](http://doi.org/10.1016/0045-6535(88)90095-1).
- Oanh NTP, Kido T, Honma S, et al. 2018. Androgen disruption by dioxin exposure in 5-year-old Vietnamese children: Decrease in serum testosterone level. *Sci Total Environ* 640-641:466-474. <http://doi.org/10.1016/j.scitotenv.2018.05.257>.
- Öberg T, Bergström JGT. 1987. Emission and chlorination pattern of PCDD/PCDF predicted from indicator parameters. *Chemosphere* 16(6):1221-1230. [http://doi.org/10.1016/0045-6535\(87\)90059-2](http://doi.org/10.1016/0045-6535(87)90059-2).
- Öberg LG, Rappe C. 1992. Biochemical formation of PCDD/Fs from chlorophenols. *Chemosphere* 25:49-45.
- Oehme M. 1991. Dispersion and transport paths of toxic persistent organochlorines to the Arctic - levels and consequences. *Sci Total Environ* 106(1-2):43-53. [http://doi.org/10.1016/0048-9697\(91\)90019-b](http://doi.org/10.1016/0048-9697(91)90019-b).
- Oehme M, Manø S, Bjerke B. 1989. Formation of polychlorinated dibenzofurans and dibenzo-p-dioxins by production processes for magnesium and refined nickel. *Chemosphere* 18(7-8):1379-1389. [http://doi.org/10.1016/0045-6535\(89\)90029-5](http://doi.org/10.1016/0045-6535(89)90029-5).
- Oehme M, Bartonova A, Knutzen J. 1990. Estimation of polychlorinated dibenzofuran and dibenzo-p-dioxin contamination of a coastal region using isomer profiles in crabs. *Environ Sci Technol* 24(12):1836-1841. <http://doi.org/10.1021/es00082a008>.
- Ogura I. 2004. Half-life of each dioxin and PCB congener in the human body. *Organohalogen Compounds* 66:3329-3337. <http://dioxin20xx.org/wp-content/uploads/pdfs/2004/04-286.pdf>. October 7, 2019.
- Ogura I, Masunaga S, Nakanishi J. 2001. Congener-specific characterization of PCDDs/PCDFs in atmospheric deposition: comparison of profiles among deposition, source, and environmental sink. *Chemosphere* 45(2):173-183. [http://doi.org/10.1016/s0045-6535\(00\)00584-1](http://doi.org/10.1016/s0045-6535(00)00584-1).
- Oishi S, Hiraga K. 1978. Is a mixture of polychlorinated dibenzofurans an inducer of hepatic porphyria? *Food Cosmet Toxicol* 16(1):47-48. [http://doi.org/10.1016/s0015-6264\(78\)80326-5](http://doi.org/10.1016/s0015-6264(78)80326-5).
- Oishi S, Hiraga K. 1980. Effect of polychlorinated biphenyl, dibenzofuran and dibenzo-p-dioxin on the susceptibility of male mice to endotoxin. *J Environ Sci Health B* 15(1):77-85. <http://doi.org/10.1080/03601238009372165>.
- Oishi S, Morita M, Fukuda H. 1978. Comparative toxicity of polychlorinated biphenyls and dibenzofurans in rats. *Toxicol Appl Pharmacol* 43(1):13-22. [http://doi.org/10.1016/s0041-008x\(78\)80028-3](http://doi.org/10.1016/s0041-008x(78)80028-3).

8. REFERENCES

- O'Keefe P, Hilker D, Meyer C, et al. 1984. Tetrachlorodibenzo-p-dioxins and tetrachlorodibenzofurans in Atlantic coast striped bass and in selected Hudson River fish, waterfowl and sediments. *Chemosphere* 13(8):849-860. [http://doi.org/10.1016/0045-6535\(84\)90158-9](http://doi.org/10.1016/0045-6535(84)90158-9).
- Okumura M. 1984. Past and current medical states of Yusho patients. *Am J Ind Med* 5(1-2):13-18.
- Okumura M, Yamanaka M, Nakamuta S. 1979. [Ten year follow-up study on serum triglyceride levels in 24 patients with PCB poisoning (author's transl)]. *Fukuoka Igaku Zasshi* 70(4):208-210. (Japanese)
- Olson JR, McGarrigle BP, Gigliotti PJ, et al. 1994. Hepatic uptake and metabolism of 2,3,7,8-tetrachlorodibenzo-p-dioxin and 2,3,7,8-tetrachlorodibenzofuran. *Fundam Appl Toxicol* 22(4):631-640. <http://doi.org/10.1006/faat.1994.1069>.
- Onozuka D, Yoshimura T, Kaneko S, et al. 2009. Mortality after exposure to polychlorinated biphenyls and polychlorinated dibenzofurans: a 40-year follow-up study of Yusho patients. *Am J Epidemiol* 169(1):86-95. <http://doi.org/10.1093/aje/kwn295>.
- Opperhuizen A, Sijm DTHM. 1990. Bioaccumulation and biotransformation of polychlorinated dibenzo-p-dioxins and dibenzofurans in fish. *Environ Toxicol Chem* 9(2):175-186. <http://doi.org/10.1002/etc.5620090207>.
- OSHA. 2019a. Occupational safety and health standards. Subpart Z - Toxic and hazardous substances. Air contaminants. Table Z-1: Limits for air contaminants. Code of Federal Regulations. 29 CFR 1910.1000. Occupational Safety and Health Administration. <https://www.osha.gov/laws-regs/regulations/standardnumber/1910/1910.1000TABLEZ1>. October 25, 2019.
- OSHA. 2019b. Occupational safety and health standards for shipyard employment. Subpart Z - Toxic and hazardous substances. Air contaminants. Code of Federal Regulations. 29 CFR 1915.1000. Occupational Safety and Health Administration. <https://www.osha.gov/laws-regs/regulations/standardnumber/1915/1915.1000>. October 25, 2019.
- OSHA. 2019c. Safety and health regulations for construction. Subpart D - Occupational health and environment controls. Gases, vapors, fumes, dusts, and mists. Code of Federal Regulations. 29 CFR 1926.55 Appendix A. Occupational Safety and Health Administration. <https://www.osha.gov/laws-regs/regulations/standardnumber/1926/1926.55AppA>. October 25, 2019.
- Pagano JJ, Garner AJ, McGoldrick DJ, et al. 2018. Age-corrected trends and toxic equivalence of PCDD/F and CP-PCBs in lake trout and walleye from the Great Lakes: 2004-2014. *Environ Sci Technol* 52(2):712-721. <http://doi.org/10.1021/acs.est.7b05568>.
- Palauschek N, Scholz B. 1987. Destruction of polychlorinated dibenzo-p-dioxins and dibenzofurans in contaminated water samples using ozone. *Chemosphere* 16(8-9):1857-1863. [http://doi.org/10.1016/0045-6535\(87\)90177-9](http://doi.org/10.1016/0045-6535(87)90177-9).
- Päpke O, Ball M, Lis ZA, et al. 1989. PCDD/PCDF in whole blood samples of unexposed persons. *Chemosphere* 19(1-6):941-948. [http://doi.org/10.1016/0045-6535\(89\)90437-2](http://doi.org/10.1016/0045-6535(89)90437-2).
- Patterson DG, Holler JS, Smith SJ, et al. 1986. Human adipose data for 2,3,7,8-tetrachlorodibenzo-p-dioxin in certain U.S. samples. *Chemosphere* 15(9-12):2055-2060. [http://doi.org/10.1016/0045-6535\(86\)90512-6](http://doi.org/10.1016/0045-6535(86)90512-6).
- Patterson DG, Hampton L, Lapeza CR, et al. 1987. High-resolution gas chromatographic/high-resolution mass spectrometric analysis of human serum on a whole-weight and lipid basis for 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Anal Chem* 59(15):2000-2005. <http://doi.org/10.1021/ac00142a023>.
- Patterson DG, Fürst P, Henderson LO, et al. 1989. Partitioning of in vivo bound PCDDs/PCDFs among various compartments in whole blood. *Chemosphere* 19(1-6):135-142. [http://doi.org/10.1016/0045-6535\(89\)90301-9](http://doi.org/10.1016/0045-6535(89)90301-9).
- Petreas MX, Goldman LR, Hayward DG, et al. 1991. Biotransfer and bioaccumulation of PCDD/PCDFs from soil: Controlled exposure studies of chickens. *Chemosphere* 23(11-12):1731-1741. [http://doi.org/10.1016/0045-6535\(91\)90021-5](http://doi.org/10.1016/0045-6535(91)90021-5).
- Petty JD, Stalling DL, Smith LM, et al. 1983. Occurrence and potential impact of PCDFs and PCDDs in aquatic ecosystems. *Trace Subst Environ Health* 17:96-102.

8. REFERENCES

- Piacitelli LA, Sweeney MH, Fingerhut MA, et al. 1992. Serum levels of PCDDs and PCDFs among workers exposed to 2,3,7,8-TCDD contaminated chemicals. *Chemosphere* 25(1-2):251-254. [http://doi.org/10.1016/0045-6535\(92\)90526-w](http://doi.org/10.1016/0045-6535(92)90526-w).
- Pluess N, Poiger H, Hohbach C, et al. 1988a. Subchronic toxicity of some chlorinated dibenzofurans (PCDFs) and a mixture of PCDFs and chlorinated dibenzodioxins (PCDDs) in rats. *Chemosphere* 17(5):973-984. [http://doi.org/10.1016/0045-6535\(88\)90068-9](http://doi.org/10.1016/0045-6535(88)90068-9).
- Pluess N, Poiger H, Hohbach C, et al. 1988b. Subchronic toxicity of 2,3,4,7,8-pentachlorodibenzofuran (PeCDF) in rats. *Chemosphere* 17(6):1099-1110. [http://doi.org/10.1016/0045-6535\(88\)90176-2](http://doi.org/10.1016/0045-6535(88)90176-2).
- Pluim HJ, Wever J, Koppe JG, et al. 1993. Intake and faecal excretion of chlorinated dioxins and dibenzofurans in breast-fed infants at different ages. *Chemosphere* 26(11):1947-1952.
- Pluim HJ, Van Der Slikke JW, Olie K, et al. 1994. Dioxins and vitamin K status of the newborn. *J Environ Sci Health A* 29(4):793-802.
- Poiger H, Pluess N, Buser HR. 1989. The metabolism of selected PCDFs in the rat. *Chemosphere* 18(1-6):259-264. [http://doi.org/10.1016/0045-6535\(89\)90129-x](http://doi.org/10.1016/0045-6535(89)90129-x).
- Poland A, Knutson JC. 1982. 2,3,7,8-Tetrachlorodibenzo-p-dioxin and related halogenated aromatic hydrocarbons: examination of the mechanism of toxicity. *Annu Rev Pharmacol Toxicol* 22:517-554. <http://doi.org/10.1146/annurev.pa.22.040182.002505>.
- Poland A, Glover E, Kende AS. 1976. Stereospecific, high affinity binding of 2,3,7,8-tetrachlorodibenzo-p-dioxin by hepatic cytosol. Evidence that the binding species is receptor for induction of aryl hydrocarbon hydroxylase. *J Biol Chem* 251(16):4936-4946.
- Poland A, Palen D, Glover E. 1982. Tumour promotion by TCDD in skin of HRS/J hairless mice. *Nature* 300(5889):271-273. <http://doi.org/10.1038/300271a0>.
- Portier CJ, Edler L, Jung D, et al. 1999. Half-lives and body burdens for dioxin and dioxin-like compounds in humans estimated from an occupational cohort in Germany. *Organohalogen Compounds* 42:129-138. <http://dioxin20xx.org/wp-content/uploads/pdfs/1999/99-453.pdf>. October 7, 2019.
- Rappe C. 1987. Global distribution of polychlorinated dioxins and dibenzofurans. *ACS Symp Ser* 338:20-33. <http://doi.org/10.1021/bk-1987-0338.ch002>.
- Rappe C. 1989. Analytical methods and exposure assessment. *Chemosphere* 18:17-12.
- Rappe C. 1991. Sources of and human exposure to PCDDs and PCDFs. In: *Banbury Report 35: Biological basis for risk assessment of dioxins and related compounds* Cold Spring Harbor, NY: Cold Spring Harbor Laboratory, 121-131.
- Rappe C. 1992. Dietary exposure and human levels of PCDs and PCDFs. *Chemosphere* 25(1-2):231-234.
- Rappe C, Buser HR. 1981. Occupational exposure to polychlorinated dioxins and dibenzofurans. *ACS Symp Ser* 149:319-342. <http://doi.org/10.1021/bk-1981-0149.ch020>.
- Rappe C, Buser HR, Bosshardt HP. 1979. Dioxins, dibenzofurans and other polyhalogenated aromatics: production, use, formation, and destruction. *Ann NY Acad Sci* 320(1 Health Effect):1-18. <http://doi.org/10.1111/j.1749-6632.1979.tb56588.x>.
- Rappe C, Nygren M, Lindstrom G, et al. 1987. Polychlorinated dibenzofurans and dibenzo-p-dioxins and other chlorinated contaminants in cow milk from various locations in Switzerland. *Environ Sci Technol* 21(10):964-970. <http://doi.org/10.1021/es50001a009>.
- Rappe C, Marklund S, Kjeller LO, et al. 1989. Long-range transport of PCDDs and PCDFs on airborne particles. *Chemosphere* 18:1283-1129.
- Rappe C, Glas B, Kjeller LO, et al. 1990a. Levels of PCDDs and PCDFs in products and effluent from the Swedish pulp and paper industry and chloralkali process. *Chemosphere* 20(10-12):1701-1706. [http://doi.org/10.1016/0045-6535\(90\)90332-n](http://doi.org/10.1016/0045-6535(90)90332-n).
- Rappe C, Andersson R, Lundström K, et al. 1990b. Levels of polychlorinated dioxins and dibenzofurans in commercial detergents and related products. *Chemosphere* 21(1-2):43-50. [http://doi.org/10.1016/0045-6535\(90\)90376-5](http://doi.org/10.1016/0045-6535(90)90376-5).

8. REFERENCES

- Rappe C, Lundstroem G, Glas B, et al. 1990c. Levels of PCDDs in milk cartons in commercial milk. *Chemosphere* 20:1649-1165.
- Rappe C, Kjeller L, Kulp S, et al. 1991a. Levels, profile and pattern of PCDDs and PCDFs in samples related to the production and use of chlorine. *Chemosphere* 23(11-12):1629-1636. [http://doi.org/10.1016/0045-6535\(91\)90010-B](http://doi.org/10.1016/0045-6535(91)90010-B).
- Rappe C, Bergqvist P, Kjeller L, et al. 1991b. Levels and patterns of PCDD and PCDF contamination in fish, crabs, and lobsters from Newark Bay and the New York Bight. *Chemosphere* 22(3-4):239-266. [http://doi.org/10.1016/0045-6535\(91\)90314-4](http://doi.org/10.1016/0045-6535(91)90314-4).
- Rayne S, Ikonomou MG, Butt CM, et al. 2005. Polychlorinated dioxins and furans from the World Trade Center attacks in exterior window films from lower Manhattan in New York City. *Environ Sci Technol* 39(7):1995-2003. <http://doi.org/10.1021/es049211k>.
- Reed LW, Hunt GT, Maisel BE, et al. 1990. Baseline assessment of PCDDs/PCDFs in the vicinity of the Elk River, Minnesota generating station. *Chemosphere* 21(1-2):159-171. [http://doi.org/10.1016/0045-6535\(90\)90388-a](http://doi.org/10.1016/0045-6535(90)90388-a).
- Reid NW, Orr DB, Shackleton MN, et al. 1990. Monitoring dioxins and dibenzofurans in precipitation in Ontario. *Chemosphere* 20(10-12):1467-1472. [http://doi.org/10.1016/0045-6535\(90\)90299-9](http://doi.org/10.1016/0045-6535(90)90299-9).
- Reischl A, Reissinger M, Thoma H, et al. 1989. Accumulation of organic air constituents by plant surfaces: Part IV. *Chemosphere* 18(1-6):561-568. [http://doi.org/10.1016/0045-6535\(89\)90167-7](http://doi.org/10.1016/0045-6535(89)90167-7).
- Remmers J, Dupuy A, McDaniel D, et al. 1992. Polychlorinated dibenzo-p-dioxin and dibenzofuran contamination in chloranil and carbazole violet. *Chemosphere* 25(7-10):1505-1508. [http://doi.org/10.1016/0045-6535\(92\)90177-S](http://doi.org/10.1016/0045-6535(92)90177-S).
- RePORTER. 2020. Chlorinated dibenzofurans. National Institutes of Health, Research Portfolio Online Reporting Tools. <https://projectreporter.nih.gov/reporter.cfm>. April 16, 2020.
- Reynolds P, Hurley SE, Petreas M, et al. 2005. Adipose levels of dioxins and risk of breast cancer. *Cancer Causes Control* 16(5):525-535. <http://doi.org/10.1007/s10552-004-7840-5>.
- Richardson TA, Klaassen CD. 2010. Role of UDP-glucuronosyltransferase (UGT) 2B2 in metabolism of triiodothyronine: effect of microsomal enzyme inducers in Sprague Dawley and UGT2B2-deficient Fischer 344 rats. *Toxicol Sci* 116(2):413-421. <http://doi.org/10.1093/toxsci/kfq125>.
- Roberts SM, Lowney YW, Stuchal LD. 2019. Bioaccessibility of polychlorinated dioxins and furans in soil from a Superfund site. *Chemosphere* 214:418-423. <http://doi.org/10.1016/j.chemosphere.2018.09.044>.
- Rogan WJ. 1989. Yu-Cheng. In: Kimbrough RD, Jensen AA, eds. Halogenated biphenyls, terphenyls, naphthalenes, dibenzodioxins and related products. 2nd ed. Amsterdam: Elsevier Science Publishers, 401-415.
- Rogan WJ, Gladen BC, Hung KL, et al. 1988. Congenital poisoning by polychlorinated biphenyls and their contaminants in Taiwan. *Science* 241(4863):334-336.
- Rohde S, Moser GA, Papke O, et al. 1999. Clearance of PCDD/Fs via the gastrointestinal tract in occupationally exposed persons. *Chemosphere* 38(14):3397-3410. [http://doi.org/10.1016/s0045-6535\(98\)00551-7](http://doi.org/10.1016/s0045-6535(98)00551-7).
- Rordorf BF. 1989. Prediction of vapor pressures, boiling points and enthalpies of fusion for twenty-nine halogenated dibenzo-p-dioxins and fifty-five dibenzofurans by a vapor pressure correlation method. *Chemosphere* 18(1-6):783-788. [http://doi.org/10.1016/0045-6535\(89\)90196-3](http://doi.org/10.1016/0045-6535(89)90196-3).
- Ross DG, Crofton KM, DeVito MJ. 2000. Changes in thyroxine glucuronidation and total serum thyroxine in rats treated with polychlorinated biphenyls and dibenzofurans. *Organohalogen Compounds* 49:243-246. <http://dioxin20xx.org/wp-content/uploads/pdfs/2000/00-436.pdf>. October 4, 2019.
- Ryan JJ. 1986. Variation of dioxins and furans in human tissues. *Chemosphere* 15(9-12):1585-1593. [http://doi.org/10.1016/0045-6535\(86\)90441-8](http://doi.org/10.1016/0045-6535(86)90441-8).
- Ryan JJ, Lizotte R, Lewis D. 1987b. Human tissue levels of PCDDs and PCDFs from a fatal pentachlorophenol poisoning. *Chemosphere* 16(8-9):1989-1996. [http://doi.org/10.1016/0045-6535\(87\)90199-8](http://doi.org/10.1016/0045-6535(87)90199-8).

8. REFERENCES

- Ryan JJ, Gasiewicz TA, Brown JF. 1990. Human body burden of polychlorinated dibenzofurans associated with toxicity based on the Yusho and Yucheng incidents. *Fundam Appl Toxicol* 15(4):722-731. [http://doi.org/10.1016/0272-0590\(90\)90188-p](http://doi.org/10.1016/0272-0590(90)90188-p).
- Ryan JJ, Guo Y, Masuda Y. 2001. Long-term human elimination and half lives of PCDFs in Yu-Cheng individuals. *Organohalogen Compounds* 54:304-307. <http://dioxin20xx.org/wp-content/uploads/pdfs/2001/01-466.pdf>. October 7, 2019.
- Ryan JJ, Schechter A, Lizotte R, et al. 1985a. Tissue distribution of dioxins and furans in humans from the general population. *Chemosphere* 14(6-7):929-932. [http://doi.org/10.1016/0045-6535\(85\)90218-8](http://doi.org/10.1016/0045-6535(85)90218-8).
- Ryan JJ, Lizotte R, Sakuma T, et al. 1985b. Chlorinated dibenzo-p-dioxins, chlorinated dibenzofurans, and pentachlorophenol in Canadian chicken and pork samples. *J Agric Food Chem* 33(6):1021-1026. <http://doi.org/10.1021/jf00066a003>.
- Ryan JJ, Schechter A, Sun NF, et al. 1986. Distribution of chlorinated dibenzo-p-dioxins and chlorinated dibenzofurans in human tissues from the general population. In: Rappe C, Choudhury G, Keith LH, eds. *Chlorinated dioxins and dibenzofurans in Perspective*. Chelsea, MI: Lewis Publishers, Inc., 3-16.
- Ryan JJ, Schechter A, Masuda Y, et al. 1987a. Comparison of PCDDs and PCDFs in the tissues of Yusho patients with those from the general population in Japan and China. *Chemosphere* 16(8-9):2017-2025. [http://doi.org/10.1016/0045-6535\(87\)90202-5](http://doi.org/10.1016/0045-6535(87)90202-5).
- Ryan JJ, Conacher HBS, Panopio LG, et al. 1991. Gas-chromatographic separations of all 136 tetra- to octa-polychlorinated dibenzo-para-dioxins and polychlorinated dibenzofurans on 9 different stationary phases. *J Chromatogr* 541:131-118.
- Ryan JJ, Shewchuk C, Lau BPY, et al. 1992. Polychlorinated dibenzo-p-dioxins and polychlorinated dibenzofurans in Canadian bleached paperboard milk containers (1988-1989) and their transfer to fluid milk. *J Agric Food Chem* 40(5):919-923.
- Ryan JJ, Levesque D, Panopio LG, et al. 1993. Elimination of polychlorinated dibenzofurans (PCDFs) and polychlorinated biphenyls (PCBs) from human blood in the Yusho and Yu-Cheng rice oil poisonings. *Arch Environ Contam Toxicol* 24(4):504-512. <http://doi.org/10.1007/bf01146170>.
- Safe S. 1990a. Polychlorinated dibenzofurans: Environmental impact, toxicology and risk assessment. In: Saxena J, ed. *Hazard assessment of chemicals - Current developments*. Vol. 7. Washington, DC: Hemisphere Publishing Company, 283-327.
- Safe S. 1990b. Polychlorinated biphenyls (PCBs), dibenzo-p-dioxins (PCDDs), dibenzofurans (PCDFs), and related compounds: environmental and mechanistic considerations which support the development of toxic equivalency factors (TEFs). *Crit Rev Toxicol* 21(1):51-88. <http://doi.org/10.3109/10408449009089873>.
- Safe S. 1991. Polychlorinated dibenzofurans: Environmental impact, toxicology, and risk assessment. *Toxic Subst J* 11:177-228.
- Safe S. 2001. Molecular biology of the Ah receptor and its role in carcinogenesis. *Toxicol Lett* 120(1-3):1-7. [http://doi.org/10.1016/s0378-4274\(01\)00301-0](http://doi.org/10.1016/s0378-4274(01)00301-0).
- Safe SH, Safe LM. 1984. Synthesis and characterization of twenty-two purified polychlorinated dibenzofuran congeners. *J Agric Food Chem* 32(1):68-67. <http://doi.org/10.1021/jf00121a018>.
- Safe S, Bunce NJ, Chittim B, et al. 1977. Photodecomposition of halogenated aromatic compounds. In: Keith LH, ed. *Identification and analysis of organic pollutants in water*. Ann Arbor, MI: Ann Arbor Science, 35-47.
- Safe S, Mason G, Keys B, et al. 1986. Polychlorinated dibenzo-p-dioxins and dibenzofurans: Correlation between in vitro and in vivo structure-activity relationships (SARs). *Chemosphere* 15(9-12):1725-1731. [http://doi.org/10.1016/0045-6535\(86\)90460-1](http://doi.org/10.1016/0045-6535(86)90460-1).
- Safe S, Brown KW, Donnelly KC, et al. 1992. Polychlorinated dibenzo-p-dioxins and dibenzofurans associated with wood-preserving chemical sites: Biomonitoring with pine needles. *Environ Sci Technol* 26(2):394-396. <http://doi.org/10.1021/es00026a023>.

8. REFERENCES

- Salisbury TB, Marcinkiewicz JL. 2002. In utero and lactational exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin and 2,3,4,7,8-pentachlorodibenzofuran reduces growth and disrupts reproductive parameters in female rats. *Biol Reprod* 66(6):1621-1626. <http://doi.org/10.1095/biolreprod66.6.1621>.
- Schechter A. 1991. Dioxins and related chemicals in humans and in the environment. In: Banbury Report 35: Biological basis for risk assessment of dioxins and related compounds. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory, 169-214.
- Schechter A, Gasiewicz TA. 1987a. Human breast milk levels of dioxins and dibenzofurans: Significance with respect to current risk assessments. *ACS Symp Ser* 338:162-173. <http://doi.org/10.1021/bk-1987-0338.ch012>.
- Schechter A, Gasiewicz TA. 1987b. Health hazard assessment of chlorinated dioxins and dibenzofurans contained in human milk. *Chemosphere* 16(8-9):2147-2154. [http://doi.org/10.1016/0045-6535\(87\)90223-2](http://doi.org/10.1016/0045-6535(87)90223-2).
- Schechter A, Ryan JJ. 1989. Blood and adipose tissue levels of PCDDs/PCDFs over three years in a patient after exposure to polychlorinated dioxins and dibenzofurans. *Chemosphere* 18(1-6):635-642. [http://doi.org/10.1016/0045-6535\(89\)90176-8](http://doi.org/10.1016/0045-6535(89)90176-8).
- Schechter A, Ryan JJ, Gitlitz G. 1986. Chlorinated dioxin and dibenzofuran levels in human adipose tissues from exposed and control populations In: Rappe C, Choudhury G, Keith LH, eds. Chlorinated dioxins and dibenzofurans in perspective. Chelsea, MI: Lewis Publishers, Inc., 51-65.
- Schechter A, Ryan JJ, Constable JD. 1987. Polychlorinated dibenzo-p-dioxin and polychlorinated dibenzofuran levels in human breast milk from Vietnam compared with cow's milk and human breast milk from the North American continent. *Chemosphere* 16(8-9):2003-2016. [http://doi.org/10.1016/0045-6535\(87\)90201-3](http://doi.org/10.1016/0045-6535(87)90201-3).
- Schechter A, Mes J, Davies D. 1989a. Polychlorinated biphenyl (PCB), DDT, DDE and hexachlorobenzene (HCB) and PCDD/F isomer levels in various organs in autopsy tissue from North American patients. *Chemosphere* 18(1-6):811-818. [http://doi.org/10.1016/0045-6535\(89\)90201-4](http://doi.org/10.1016/0045-6535(89)90201-4).
- Schechter A, Ryan JJ, Constable JD. 1989c. Chlorinated dioxins and dibenzofurans in human milk from Japan, India, and the United States of America. *Chemosphere* 18(1-6):975-980. [http://doi.org/10.1016/0045-6535\(89\)90225-7](http://doi.org/10.1016/0045-6535(89)90225-7).
- Schechter A, Pöpke O, Ball M. 1990a. Evidence for transplacental transfer of dioxins from mother to fetus: Chlorinated dioxin and dibenzofuran levels in the livers of stillborn infants. *Chemosphere* 21(8):1017-1022. [http://doi.org/10.1016/0045-6535\(90\)90124-C](http://doi.org/10.1016/0045-6535(90)90124-C).
- Schechter A, Ryan JJ, Kostyniak PJ. 1990b. Decrease over a six year period of dioxin and dibenzofuran tissue levels in a single patient following exposure. *Chemosphere* 20(7-9):911-917. [http://doi.org/10.1016/0045-6535\(90\)90200-D](http://doi.org/10.1016/0045-6535(90)90200-D).
- Schechter A, Tiernan T, Schaffner F, et al. 1985a. Patient fat biopsies for chemical analysis and liver biopsies for ultrastructural characterization after exposure to polychlorinated dioxins, furans and PCBs. *Environ Health Perspect* 60:241-254. <http://doi.org/10.1289/ehp.8560241>.
- Schechter A, Ryan JJ, Lizotte R, et al. 1985b. Chlorinated dibenzodioxins and dibenzofurans in human adipose tissue from exposed and control New York State patients. *Chemosphere* 14:933-993.
- Schechter A, Fürst P, Fürst C, et al. 1989b. Levels of polychlorinated dibenzodioxins and dibenzofurans in cow's milk and in soy bean derived infant formulas sold in the United States and other countries. *Chemosphere* 19(1-6):913-918. [http://doi.org/10.1016/0045-6535\(89\)90431-1](http://doi.org/10.1016/0045-6535(89)90431-1).
- Schechter A, Fürst P, Ryan JJ, et al. 1989d. Polychlorinated dioxin and dibenzofuran levels from human milk from several locations in the United States, Germany and Vietnam. *Chemosphere* 19:979-984.
- Schechter A, Fürst P, Fürst C, et al. 1990c. Levels of dioxins, dibenzofurans and other chlorinated xenobiotics in human milk from the Soviet Union. *Chemosphere* 20:927-993.
- Schechter AJ, Malkin R, Papke O, et al. 1991a. Dioxin levels in blood of municipal incinerator workers. *Med Sci Res* 19:331-332.

8. REFERENCES

- Schechter A, Fürst P, Fürst C, et al. 1991b. Dioxins, dibenzofurans and selected chlorinated organic compounds in human milk and blood from Cambodia, Germany, Thailand, the USA, the USSR, and Vietnam. *Chemosphere* 23(11-12):1903-1912. [http://doi.org/10.1016/0045-6535\(91\)90037-E](http://doi.org/10.1016/0045-6535(91)90037-E).
- Schechter A, Pöpke O, Ball M, et al. 1991c. Partitioning of dioxins and dibenzofurans: Whole blood, blood plasma and adipose tissue. *Chemosphere* 23(11-12):1913-1919. [http://doi.org/10.1016/0045-6535\(91\)90038-F](http://doi.org/10.1016/0045-6535(91)90038-F).
- Schechter A, Startin J, Wright C, et al. 1993. Dioxin levels in food from the United States with estimated daily intake. *Organohalogen Compounds* 13:97-100. <http://dioxin20xx.org/wp-content/uploads/pdfs/1993/93-302.pdf>. April 24, 2020.
- Schlummer M, Moser GA, McLachlan MS. 1998. Digestive tract absorption of PCDD/Fs, PCBs, and HCB in humans: mass balances and mechanistic considerations. *Toxicol Appl Pharmacol* 152(1):128-137. <http://doi.org/10.1006/taap.1998.8487>.
- Schoeny R. 1982. Mutagenicity testing of chlorinated biphenyls and chlorinated dibenzofurans. *Mutat Res* 101(1):45-56. [http://doi.org/10.1016/0165-1218\(82\)90164-1](http://doi.org/10.1016/0165-1218(82)90164-1).
- Schramm KW, Kuettner T, Weber S, et al. 1992. Dioxin hair analysis as monitoring pool. *Chemosphere* 24(3):351-358. [http://doi.org/10.1016/0045-6535\(92\)90303-9](http://doi.org/10.1016/0045-6535(92)90303-9).
- Schrey P, Wittsiepe J, Mackrodt P, et al. 1998. Human fecal PCDD/F-excretion exceeds the dietary intake. *Chemosphere* 37(9-12):1825-1831. [http://doi.org/10.1016/s0045-6535\(98\)00248-3](http://doi.org/10.1016/s0045-6535(98)00248-3).
- Sedman RM, Esparza JR. 1991. Evaluation of the public health risks associated with semivolatile metal and dioxin emissions from hazardous waste incinerators. *Environ Health Perspect* 94:181-187. <http://doi.org/10.1289/ehp.94-1567951>.
- Sherman RK, Clement RE, Tashiro C. 1990. The distribution of polychlorinated dibenzo-p-dioxins and dibenzofurans in Jackfish Bay, Lake Superior, in relation to a kraft pulp mill effluent. *Chemosphere* 20(10-12):1641-1648. [http://doi.org/10.1016/0045-6535\(90\)90324-m](http://doi.org/10.1016/0045-6535(90)90324-m).
- Shigematsu N, Norimatsu Y, Ishibashi T, et al. 1971. [Clinical and experimental studies on respiratory involvement in chlorobiphenyls poisoning]. *Fukuoka Igaku Zasshi* 62:150-156. (Japanese)
- Shigematsu N, Ishimaru S, Ikeda T, et al. 1977. [Further studies on respiratory disorders in polychlorinated biphenyls (PCB) poisoning. Relationship between respiratory disorders and PCB concentrations in blood and sputum (author's transl)]. *Fukuoka Igaku Zasshi* 68(3):133-138. (Japanese)
- Siebert PC, Alston DR, Walsh JF, et al. 1987. Statistical properties of available worldwide MSW combustion dioxin/furan emissions. *Proc APCA Annu Meet* 80:1-20.
- Sijm DTHM, Wever H, de Vries PJ, et al. 1989. Octan-1-ol/water partition coefficients of polychlorinated dibenzo-p-dioxins and dibenzofurans: Experimental values determined with a stirring method. *Chemosphere* 19(1-6):263-266. [http://doi.org/10.1016/0045-6535\(89\)90321-4](http://doi.org/10.1016/0045-6535(89)90321-4).
- Smith RM, O'Keefe PW, Hilker DR, et al. 1986. Determination of picogram per cubic meter concentrations of tetra- and pentachlorinated dibenzofurans and dibenzo-p-dioxins in indoor air by high-resolution gas chromatography/high-resolution mass spectrometry. *Anal Chem* 58(12):2414-2420. <http://doi.org/10.1021/ac00125a014>.
- Smith RM, O'Keefe PW, Aldous KM, et al. 1990. Chlorinated dibenzofurans and dioxins in atmospheric samples from cities in New York. *Environ Sci Technol* 24:1502-1150.
- Sola-Gutierrez C, Schroder S, San Roman MF, et al. 2019. PCDD/Fs traceability during triclosan electrochemical oxidation. *J Hazard Mater* 369:584-592. <http://doi.org/10.1016/j.jhazmat.2019.02.066>.
- Someshwar AV, Jain AK, Whittemore RC, et al. 1990. The effects of sludge burning on the PCDD/PCDF content of ashes from pulp and paper mill hog fuel boilers. *Chemosphere* 20(10-12):1715-1722. [http://doi.org/10.1016/0045-6535\(90\)90334-p](http://doi.org/10.1016/0045-6535(90)90334-p).
- Stalling DL, Norstrom RJ, Smith LM, et al. 1985. Patterns of PCDD, PCDF, and PCB contamination in Great Lakes fish and birds and their characterization by principal components analysis. *Chemosphere* 14(6-7):627-643. [http://doi.org/10.1016/0045-6535\(85\)90171-7](http://doi.org/10.1016/0045-6535(85)90171-7).

8. REFERENCES

- Stanley JS, Boggess KE, Onstot J, et al. 1986. PCDDs and PCDFs in human adipose tissue from the EPA FY82 NHATS repository. *Chemosphere* 15(9-12):1605-1612. [http://doi.org/10.1016/0045-6535\(86\)90444-3](http://doi.org/10.1016/0045-6535(86)90444-3).
- Steer P, Tashiro C, Clement R, et al. 1990. Ambient air sampling of polychlorinated dibenzo-p-dioxins and dibenzofurans in Ontario: preliminary results. *Chemosphere* 20(10-12):1431-1437. [http://doi.org/10.1016/0045-6535\(90\)90294-4](http://doi.org/10.1016/0045-6535(90)90294-4).
- Stephens RD. 1986. Transformer fire Spear Street Tower One Market Plaza San Francisco, California. *Chemosphere* 15(9-12):1281-1289. [http://doi.org/10.1016/0045-6535\(86\)90402-9](http://doi.org/10.1016/0045-6535(86)90402-9).
- Stieglitz L, Zwick G, Beck J, et al. 1989. Carbonaceous particles in fly ash -a source for the de-novo synthesis of organochloro compounds. *Chemosphere* 19(1-6):283-290. [http://doi.org/10.1016/0045-6535\(89\)90325-1](http://doi.org/10.1016/0045-6535(89)90325-1).
- Sun XL, Kido T, Honma S, et al. 2016. Influence of dioxin exposure upon levels of prostate-specific antigen and steroid hormones in Vietnamese men. *Environ Sci Pollut Res Int* 23(8):7807-7813. <http://doi.org/10.1007/s11356-015-5931-3>.
- Sun XL, Kido T, Honma S, et al. 2017. The relationship between dioxins exposure and risk of prostate cancer with steroid hormone and age in Vietnamese men. *Sci Total Environ* 595:842-848. <http://doi.org/10.1016/j.scitotenv.2017.04.013>.
- Sutter TR, Greenlee WF. 1992. Classification of members of the Ah gene battery. *Chemosphere* 25(1-2):223-226.
- Svenson A, Kjeller L, Rappe C. 1989a. Enzymatic chlorophenol oxidation as a means of chlorinated dioxin and dibenzofuran formation. *Chemosphere* 19:585-558.
- Svenson A, Kjeller L, Rappe C. 1989b. Enzyme-mediated formation of 2,3,7,8-tetra-substituted chlorinated dibenzodioxins and dibenzofurans. *Environ Sci Technol* 23:900-990.
- Svensson BG, Nilsson A, Hansson M, et al. 1991. Exposure to dioxins and dibenzofurans through the consumption of fish. *N Engl J Med* 324(1):8-12. <http://doi.org/10.1056/NEJM199101033240102>.
- Tai H, McReynolds JH, Goldstein JA, et al. 1993. Cytochrome P4501A1 mediates the metabolism of 2,3,7,8-tetrachlorodibenzofuran in the rat and human. *Toxicol Appl Pharmacol* 123(1):34-42. <http://doi.org/10.1006/taap.1993.1218>.
- Tai PT, Nishijo M, Anh NTN, et al. 2013. Perinatal dioxin exposure and neurodevelopment of children aged 3 years - a follow up study in Vietnam. *Organohalogen Compounds* 75:1321-1324. <http://dioxin20xx.org/wp-content/uploads/pdfs/2013/8009.pdf>. October 7, 2019.
- Takayama K, Miyata H, Mimura M, et al. 1991. Evaluation of biological effects of polychlorinated compounds found in contaminated cooking oil responsible for the disease "Yusho". *Chemosphere* 22(5-6):537-546. [http://doi.org/10.1016/0045-6535\(91\)90065-1](http://doi.org/10.1016/0045-6535(91)90065-1).
- Takeshita R, Akimoto Y. 1989. Control of PCDD and PCDF formation in fluidized bed incinerators. *Chemosphere* 19(1-6):345-352. [http://doi.org/10.1016/0045-6535\(89\)90334-2](http://doi.org/10.1016/0045-6535(89)90334-2).
- Taki I, Hisanaga S, Amagase Y. 1969. [Report on Yusho (chlorobiphenyls poisoning) pregnant women and their fetuses]. *Fukuoka Igaku Zasshi* 60:471-474. (Japanese)
- Tanabe S, Kannan N, Wakimoto T, et al. 1989. Isomer-specific determination and toxic evaluation of potentially hazardous coplanar PCBs, dibenzofurans and dioxins in the tissues of "Yusho" PCB poisoning victim and in the causal oil. *Toxicological & Environmental Chemistry* 24(4):215-231. <http://doi.org/10.1080/02772248909357494>.
- Taucher JA, Hannah DJ, Green NJL, et al. 1992. PCDD, PCDF and PCB emissions under variable operating conditions from a waste oil furnace. *Chemosphere* 25(7-10):1429-1433. [http://doi.org/10.1016/0045-6535\(92\)90165-n](http://doi.org/10.1016/0045-6535(92)90165-n).
- Taura J, Takeda T, Fujii M, et al. 2014. 2,3,4,7,8-Pentachlorodibenzofuran is far less potent than 2,3,7,8-tetrachlorodibenzo-p-dioxin in disrupting the pituitary-gonad axis of the rat fetus. *Toxicol Appl Pharmacol* 281(1):48-57. <http://doi.org/10.1016/j.taap.2014.09.001>.
- Tawara K, Nishijo M, Nakagawa H, et al. 2008. Association of having birth defect children with serum dioxin levels in Vietnam. *Organohalogen Compounds* 69:568-571. <http://dioxin20xx.org/wp-content/uploads/pdfs/2007/07-305.pdf>. October 4, 2019.

8. REFERENCES

- Tawara K, Nishijo M, Honda R, et al. 2009. Effects of maternal dioxin exposure on newborn size at birth among Japanese mother-infant pairs. *Environ Health Prev Med* 14(2):88-95. <http://doi.org/10.1007/s12199-008-0061-x>.
- Thoma H, Mücke W, Kauert G. 1990. Comparison of the polychlorinated dibenzo-p-dioxin and dibenzofuran in human tissue and human liver. *Chemosphere* 20(3-4):433-442. [http://doi.org/10.1016/0045-6535\(90\)90074-4](http://doi.org/10.1016/0045-6535(90)90074-4).
- Thomas RG. 1982. Volatilization from water. In: Lyman WH, Riehl WF, Rosenblatt DH, eds. *Handbook of chemical property estimation methods*. New York, NY: McGraw-Hill Book Company, 15-11 to 15-34.
- Thompson HC, Kendall DC, Korfmacher WA, et al. 1986. Assessment of the contamination of a multibuilding facility by polychlorinated biphenyls, polychlorinated dibenzo-p-dioxins, and polychlorinated dibenzofurans. *Environ Sci Technol* 20(6):597-603. <http://doi.org/10.1021/es00148a009>.
- Tiernan TO, Taylor ML, Van Ness GF, et al. 1984. Analyses of human tissues for chlorinated dibenzo-p-dioxins and chlorinated dibenzofurans: The state of the art. In: Lowrance WW, ed. *Public health risks of the dioxins*. New York, NY: Rockefeller University, 31-56.
- Tiernan TO, Taylor ML, Garrett JH, et al. 1985. Sources and fate of polychlorinated dibenzodioxins, dibenzofurans and related compounds in human environments. *Environ Health Perspect* 59:145-158. <http://doi.org/10.1289/ehp.59-1568078>.
- Tiernan TO, Wagel DJ, Garrett JH, et al. 1989. Laboratory and field tests to demonstrate the efficacy of KPEG reagent for detoxification of hazardous wastes containing polychlorinated dibenzo-p-dioxins (PCDD) and dibenzofurans (PCDF) and soils contaminated with such chemical wastes. *Chemosphere* 18(1-6):835-841. [http://doi.org/10.1016/0045-6535\(89\)90205-1](http://doi.org/10.1016/0045-6535(89)90205-1).
- Tirler W, Voto G, Donega M. 2001. PCDD/F, PCB and hexachlorobenzene levels in hair. *Organohalogen Compounds* 52:290-292. <http://dioxin20xx.org/wp-content/uploads/pdfs/2001/01-534.pdf>. October 7, 2019.
- Todaka T, Hirakawa H, Kajiwara J, et al. 2010. Relationship between the concentrations of polychlorinated dibenzo-p-dioxins, polychlorinated dibenzofurans, and polychlorinated biphenyls in maternal blood and those in breast milk. *Chemosphere* 78(2):185-192. <http://doi.org/10.1016/j.chemosphere.2009.09.047>.
- Tondeur Y, Beckert WF, Billets S, et al. 1989. Method 8290: An analytical protocol for the multimedia characterization of polychlorinated dibenzodioxins and dibenzofurans by high-resolution gas chromatography/high-resolution mass spectrometry. *Chemosphere* 18(1-6):119-131. [http://doi.org/10.1016/0045-6535\(89\)90112-4](http://doi.org/10.1016/0045-6535(89)90112-4).
- Tong HY, Karasek FW. 1986. Comparison of PCDD and PCDF in flyash collected from municipal incinerators of different countries. *Chemosphere* 15(9-12):1219-1224. [http://doi.org/10.1016/0045-6535\(86\)90394-2](http://doi.org/10.1016/0045-6535(86)90394-2).
- Travis CC, Hattemer-Frey HA. 1989. A perspective on dioxin emissions from municipal solid waste incinerators. *Risk Anal* 9(1):91-97. <http://doi.org/10.1111/j.1539-6924.1989.tb01223.x>.
- TRI18. 2020. TRI explorer: Providing access to EPA's toxics release inventory data. Washington, DC: Toxics Release Inventory. U.S. Environmental Protection Agency. <http://www.epa.gov/triexplorer/>. February 10, 2020.
- Tsai PC, Ko YC, Huang W, et al. 2007. Increased liver and lupus mortalities in 24-year follow-up of the Taiwanese people highly exposed to polychlorinated biphenyls and dibenzofurans. *Sci Total Environ* 374(2-3):216-222. <http://doi.org/10.1016/j.scitotenv.2006.12.024>.
- Tsukimori K, Tokunaga S, Shibata S, et al. 2008. Long-term effects of polychlorinated biphenyls and dioxins on pregnancy outcomes in women affected by the Yusho incident. *Environ Health Perspect* 116(5):626-630. <http://doi.org/10.1289/ehp.10686>.
- Tsukimori K, Uchi H, Mitoma C, et al. 2011. Comparison of the concentrations of polychlorinated biphenyls and dioxins in mothers affected by the Yusho incident and their children. *Chemosphere* 84(7):928-935. <http://doi.org/10.1016/j.chemosphere.2011.06.009>.

8. REFERENCES

- Tsukimori K, Morokuma S, Hori T, et al. 2013. Characterization of placental transfer of polychlorinated dibenzo-p-dioxins, dibenzofurans and polychlorinated biphenyls in normal pregnancy. *J Obstet Gynaecol* 39(1):83-90. <http://doi.org/10.1111/j.1447-0756.2012.01906.x>.
- Tue NM, Suzuki G, Takahashi S, et al. 2013. Dioxin-related compounds in house dust from New York State: occurrence, in vitro toxic evaluation and implications for indoor exposure. *Environ Pollut* 181:75-80. <http://doi.org/10.1016/j.envpol.2013.06.010>.
- Tuomisto J, Airaksinen R, Kiviranta H, et al. 2016. A pharmacokinetic analysis and dietary information are necessary to confirm or reject the hypothesis on persistent organic pollutants causing type 2 diabetes. *Toxicol Lett* 261:41-48. <http://doi.org/10.1016/j.toxlet.2016.08.024>.
- Tysklind M, Rappe C. 1991. Photolytic transformation of polychlorinated dioxins and dibenzofurans in fly ash. *Chemosphere* 23(8-10):1365-1375. [http://doi.org/10.1016/0045-6535\(91\)90161-6](http://doi.org/10.1016/0045-6535(91)90161-6).
- Uemura H, Arisawa K, Hiyoshi M, et al. 2009. Prevalence of metabolic syndrome associated with body burden levels of dioxin and related compounds among Japan's general population. *Environ Health Perspect* 117(4):568-573. <http://doi.org/10.1289/ehp.0800012>.
- Uzawa H, Ito Y, Notomi A, et al. 1969. [Hyperglyceridemia resulting from intake of rice oil contaminated with chlorinated biphenyls]. *Fukuoka Igaku Zasshi* 60:449-454. (Japanese)
- Vainio H, Hesso A, Jappinen P. 1989. Chlorinated dioxins and dibenzofurans in the environment - a hazard to public health? *Scand J Work Environ Health* 15(6):377-382. <http://doi.org/10.5271/sjweh.1837>.
- Van den Berg M, Olie K, Hutzinger. 1985. Polychlorinated dibenzofurans (PCDFs): Environmental occurrence and physical, chemical and biological properties. *Toxicological & Environmental Chemistry* 9(3):171-217. <http://doi.org/10.1080/02772248509357079>.
- van den Berg M, van der Wielen FWM, Olie K, et al. 1986. The presence of PCDDs and PCDFs in human breast milk from the Netherlands. *Chemosphere* 15(6):693-706. [http://doi.org/10.1016/0045-6535\(86\)90035-4](http://doi.org/10.1016/0045-6535(86)90035-4).
- Van den Berg M, de Jongh J, Eckhart P, et al. 1989. Disposition and elimination of three polychlorinated dibenzofurans in the liver of the rat. *Fundam Appl Toxicol* 12:738-774.
- Van den Berg M, Birnbaum LS, Denison M, et al. 2006. The 2005 World Health Organization reevaluation of human and mammalian toxic equivalency factors for dioxins and dioxin-like compounds. *Toxicol Sci* 93(2):223-241. <http://doi.org/10.1093/toxsci/kfl055>.
- van Ede KI, Aylward LL, Andersson PL, et al. 2013. Tissue distribution of dioxin-like compounds: potential impacts on systemic relative potency estimates. *Toxicol Lett* 220(3):294-302. <http://doi.org/10.1016/j.toxlet.2013.05.001>.
- Van Luong H, Tai PT, Nishijo M, et al. 2018. Association of dioxin exposure and reproductive hormone levels in men living near the Bien Hoa airbase, Vietnam. *Sci Total Environ* 628-629:484-489. <http://doi.org/10.1016/j.scitotenv.2018.02.094>.
- Van Tung D, Kido T, Honma S, et al. 2016. Low birth weight of Vietnamese infants is related to their mother's dioxin and glucocorticoid levels. *Environ Sci Pollut Res Int* 23(11):10922-10929. <http://doi.org/10.1007/s11356-016-6233-0>.
- Vanden Heuvel JP, Clark GC, Tritscher A, et al. 1994. Accumulation of polychlorinated dibenzo-p-dioxins and dibenzofurans in liver of control laboratory rats. *Fundam Appl Toxicol* 23(3):465-469.
- Vartiainen T, Jaakkola JJ, Saarikoski S, et al. 1998. Birth weight and sex of children and the correlation to the body burden of PCDDs/PCDFs and PCBs of the mother. *Environ Health Perspect* 106(2):61-66. <http://doi.org/10.1289/ehp.9810661>.
- Veerkamp W, Wever J, Hutzinger. 1981. The metabolism of some chlorinated dibenzofurans by rats. *Chemosphere* 10:397-340.
- Venier M, Ferrario J, Hites RA. 2009. Polychlorinated dibenzo-p-dioxins and dibenzofurans in the atmosphere around the Great Lakes. *Environ Sci Technol* 43(4):1036-1041. <http://doi.org/10.1021/es802644w>.
- Venkatesan AK, Halden RU. 2014. Contribution of polybrominated dibenzo-p-dioxins and dibenzofurans (PBDD/Fs) to the toxic equivalency of dioxin-like compounds in archived biosolids

8. REFERENCES

- from the U.S. EPA's 2001 national sewage sludge survey. *Environ Sci Technol* 48(18):10843-10849. <http://doi.org/10.1021/es503110j>.
- Vigh E, Colombo A, Benfenati E, et al. 2013. Individual breast milk consumption and exposure to PCBs and PCDD/Fs in Hungarian infants: a time-course analysis of the first three months of lactation. *Sci Total Environ* 449:336-344. <http://doi.org/10.1016/j.scitotenv.2013.01.024>.
- Villanueva EC, Jennings RW, Burse VW, et al. 1974. Evidence of chlorodibenzo-p-dioxin and chlorodibenzofuran in hexachlorobenzene. *J Agric Food Chem* 22(5):916-917. <http://doi.org/10.1021/jf60195a021>.
- Vos JG, Luster MI. 1989. Immune alterations. In: Kimbrough RD JA, ed. *Halogenated biphenyls, terphenyls, naphthalenes, dibenzodioxins and related products*. 2nd ed. Amsterdam: Elsevier Science Publishers, 295-322.
- Waddell D, Chittim B, Clement R, et al. 1990. Database of PCDD/PCDF levels in ambient air and in samples related to the pulp and paper industry. *Chemosphere* 20(10-12):1463-1466. [http://doi.org/10.1016/0045-6535\(90\)90298-8](http://doi.org/10.1016/0045-6535(90)90298-8).
- Wang S, Guo YL, Lin C, et al. 2002. Body burden of polychlorinated dibenzo-p-dioxins, dibenzofurans and biphenyls in pregnant women and their infants - correlation between prenatal and postnatal exposure to infants. *Organohalogen Compounds* 55:255-258. <http://dioxin20xx.org/wp-content/uploads/pdfs/2002/02-61.pdf>. October 4, 2019.
- Wang S, Chen T, Hsu J, et al. 2003. Neonatal and childhood teeth in relation to perinatal exposure to polychlorinated biphenyls and dibenzofurans: observations of the Yucheng children in Taiwan. *Environ Res* 93(2):131-137. [http://doi.org/10.1016/s0013-9351\(03\)00040-9](http://doi.org/10.1016/s0013-9351(03)00040-9).
- Wang S, Lin C, Guo YL, et al. 2004. Infant exposure to polychlorinated dibenzo-p-dioxins, dibenzofurans and biphenyls (PCDD/Fs, PCBs) - correlation between prenatal and postnatal exposure. *Chemosphere* 54(10):1459-1473. <http://doi.org/10.1016/j.chemosphere.2003.08.012>.
- Wang S, Tsai P, Yang C, et al. 2008. Increased risk of diabetes and polychlorinated biphenyls and dioxins: a 24-year follow-up study of the Yucheng cohort. *Diabetes Care* 31(8):1574-1579. <http://doi.org/10.2337/dc07-2449>.
- Wang Z, Hang JG, Feng H, et al. 2019. Effects of perinatal dioxin exposure on development of children: a 3-year follow-up study of China cohort. *Environ Sci Pollut Res Int* 26(20):20780-20786. <http://doi.org/10.1007/s11356-019-05362-0>.
- Watanabe MX, Kunisue T, Ueda N, et al. 2013. Toxicokinetics of dioxins and other organochlorine compounds in Japanese people: association with hepatic CYP1A2 expression levels. *Environ Int* 53:53-61. <http://doi.org/10.1016/j.envint.2012.12.008>.
- Weber H, Birnbaum LS. 1985. 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) and 2,3,7,8-tetrachlorodibenzofuran (TCDF) in pregnant C57BL/6N mice: distribution to the embryo and excretion. *Arch Toxicol* 57(3):159-162. <http://doi.org/10.1007/bf00290880>.
- Weber H, Lamb JC, Harris MW, et al. 1984. Teratogenicity of 2,3,7,8-tetrachlorodibenzofuran (TCDF) in mice. *Toxicol Lett* 20(2):183-188. [http://doi.org/10.1016/0378-4274\(84\)90145-0](http://doi.org/10.1016/0378-4274(84)90145-0).
- Weber H, Harris MW, Haseman JK, et al. 1985. Teratogenic potency of TCDD, TCDF and TCDD-TCDF combinations in C57BL/6N mice. *Toxicol Lett* 26(2-3):159-167. [http://doi.org/10.1016/0378-4274\(85\)90161-4](http://doi.org/10.1016/0378-4274(85)90161-4).
- Weichenthal S, Van Rijswijk D, Kulka R, et al. 2015. The impact of a landfill fire on ambient air quality in the north: A case study in Iqaluit, Canada. *Environ Res* 142:46-50. <http://doi.org/10.1016/j.envres.2015.06.018>.
- Whitlock JP. 1999. Induction of cytochrome P4501A1. *Annu Rev Pharmacol Toxicol* 39:103-125. <http://doi.org/10.1146/annurev.pharmtox.39.1.103>.
- Whittemore RC, LaFleur LE, Gillespie WJ, et al. 1990. USEPA/paper industry cooperative dioxin study: The 104 mill study. *Chemosphere* 20(10-12):1625-1632. [http://doi.org/10.1016/0045-6535\(90\)90322-k](http://doi.org/10.1016/0045-6535(90)90322-k).

8. REFERENCES

- WHO. 2000. Air quality guidelines for Europe. 2nd. Copenhagen, Denmark: World Health Organization. http://www.euro.who.int/__data/assets/pdf_file/0005/74732/E71922.pdf. April 22, 2020.
- WHO. 2010. Guidelines for indoor air quality: Selected pollutants. Geneva, Switzerland: World Health Organization. http://www.euro.who.int/__data/assets/pdf_file/0009/128169/e94535.pdf. April 25, 2012.
- WHO. 2017. Guidelines for drinking-water quality. 4th incorporating the first addendum. Geneva, Switzerland: World Health Organization. <http://apps.who.int/iris/bitstream/10665/254637/1/9789241549950-eng.pdf?ua=1>. February 28, 2017.
- Wiberg K, Lundström K, Glas B, et al. 1989. PCDDs and PCDFs in consumers' paper products. *Chemosphere* 19(1-6):735-740. [http://doi.org/10.1016/0045-6535\(89\)90400-1](http://doi.org/10.1016/0045-6535(89)90400-1).
- Williams DT, LeBel GL, Benoit FM. 1992. Polychlorodibenzodioxins and polychlorodibenzofurans in dioxazine dyes and pigments. *Chemosphere* 24(2):169-180. [http://doi.org/10.1016/0045-6535\(92\)90390-d](http://doi.org/10.1016/0045-6535(92)90390-d).
- Wittsiepe J, Erenkamper B, Welge P, et al. 2007a. Bioavailability of PCDD/F from contaminated soil in young Goettingen minipigs. *Chemosphere* 67(9):S355-364. <http://doi.org/10.1016/j.chemosphere.2006.05.129>.
- Wittsiepe J, Fürst P, Schrey P, et al. 2007b. PCDD/F and dioxin-like PCB in human blood and milk from German mothers. *Chemosphere* 67(9):S286-294. <http://doi.org/10.1016/j.chemosphere.2006.05.118>.
- Yamagishi T, Miyazaki T, Akiyama K, et al. 1981. Polychlorinated dibenzo-p-dioxins and dibenzofurans in commercial diphenyl ether herbicides, and in freshwater fish collected from the application area. *Chemosphere* 10(10):1137-1144. [http://doi.org/10.1016/0045-6535\(81\)90184-3](http://doi.org/10.1016/0045-6535(81)90184-3).
- Yamaguchi A, Yoshimura T, Kuratsune M. 1971. [Investigation concerning babies born from women who consumed oil contaminated with chlorobiphenyl]. *Fukuoka Igaku Zasshi* 62(1):117-122. (Japanese)
- Yanders AF, Orazio CE, Puri RK, et al. 1989. On translocation of 2,3,7,8-tetrachlorodibenzo-p-dioxin: Time dependent analysis at the Times Beach experimental site. *Chemosphere* 19(1-6):429-432. [http://doi.org/10.1016/0045-6535\(89\)90347-0](http://doi.org/10.1016/0045-6535(89)90347-0).
- Yang C, Yu M, Guo H, et al. 2005. The endocrine and reproductive function of the female Yucheng adolescents prenatally exposed to PCBs/PCDFs. *Chemosphere* 61(3):355-360. <http://doi.org/10.1016/j.chemosphere.2005.02.089>.
- Yang C, Wang Y, Chen P, et al. 2008. Exposure to a mixture of polychlorinated biphenyls and polychlorinated dibenzofurans resulted in a prolonged time to pregnancy in women. *Environ Health Perspect* 116(5):599-604. <http://doi.org/10.1289/ehp.10715>.
- Yang C, Huang T, Lin K, et al. 2011. Menstrual effects among women exposed to polychlorinated biphenyls and dibenzofurans. *Environ Res* 111(2):288-294. <http://doi.org/10.1016/j.envres.2010.11.011>.
- Yen Y, Lan S, Yang C, et al. 1994. Follow-up study of intrauterine growth of transplacental Yu-Cheng babies in Taiwan. *Bull Environ Contam Toxicol* 53(5):633-641. <http://doi.org/10.1007/bf00196933>.
- Yoshihara S, Nagata K, Yoshimura H, et al. 1981. Inductive effect on hepatic enzymes and acute toxicity of individual polychlorinated dibenzofuran congeners in rats. *Toxicol Appl Pharmacol* 59(3):580-588. [http://doi.org/10.1016/0041-008x\(81\)90313-6](http://doi.org/10.1016/0041-008x(81)90313-6).
- Yoshimura T. 1974. [Epidemiological study on Yusho babies born to mothers who had consumed oil contaminated by PCB]. *Fukuoka Igaku Zasshi* 65:74-80. (Japanese)
- Yoshimura T, Kaneko S, Hayabuchi H. 2001. Sex ratio in offspring of those affected by dioxin and dioxin-like compounds: the Yusho, Seveso, and Yucheng incidents. *Occup Environ Med* 58(8):540-541. <http://doi.org/10.1136/oem.58.8.540>.

8. REFERENCES

- Young AL. 1984. Analysis of dioxins and furans in human adipose tissue. In: Lowrance WW, ed. Public health risks of the dioxins: Proceedings of a symposium held in New York City on October 19-20, 1983. New York, NY: Rockefeller University, 63-75.
- Yu M, Hsu C, Gladen BC, et al. 1991. In utero PCB/PCDF exposure: relation of developmental delay to dysmorphology and dose. *Neurotoxicol Teratol* 13(2):195-202. [http://doi.org/10.1016/0892-0362\(91\)90011-k](http://doi.org/10.1016/0892-0362(91)90011-k).
- Yu M, Guo YL, Hsu C, et al. 1997. Increased mortality from chronic liver disease and cirrhosis 13 years after the Taiwan "Yucheng" ("oil disease") incident. *Am J Ind Med* 31(2):172-175. [http://doi.org/10.1002/\(sici\)1097-0274\(199702\)31:2<172::aid-ajim6>3.0.co;2-1](http://doi.org/10.1002/(sici)1097-0274(199702)31:2<172::aid-ajim6>3.0.co;2-1).
- Yu M, Hsin J, Hsu C, et al. 1998. The immunologic evaluation of the Yucheng children. *Chemosphere* 37(9-12):1855-1865. [http://doi.org/10.1016/s0045-6535\(98\)00251-3](http://doi.org/10.1016/s0045-6535(98)00251-3).
- Zacharewski T, Safe L, Safe S, et al. 1989. Comparative analysis of polychlorinated dibenzo-p-dioxin and dibenzofuran congeners in Great Lakes fish extracts by gas chromatography-mass spectrometry and in vitro enzyme induction activities. *Environ Sci Technol* 23(6):730-735. <http://doi.org/10.1021/es00064a012>.
- Zeytun A, McKallip RJ, Fisher M, et al. 2002. Analysis of 2,3,7,8-tetrachlorodibenzo-p-dioxin-induced gene expression profile in vivo using pathway-specific cDNA arrays. *Toxicology* 178(3):241-260. [http://doi.org/10.1016/s0300-483x\(02\)00230-5](http://doi.org/10.1016/s0300-483x(02)00230-5).
- Zitko V. 1992. Patterns of 2,3,7,8-substituted chlorinated dibenzodioxins and dibenzofurans in aquatic fauna. *Sci Total Environ* 111(2-3):95-108. [http://doi.org/10.1016/0048-9697\(92\)90349-W](http://doi.org/10.1016/0048-9697(92)90349-W).