5. References

ATSDR 1995. Toxicological Profile for Diethyl Phthalate. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

ATSDR 1997. Toxicological Profile for Di-n-Octyl Phthalate. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

ATSDR. 1998. Toxicological Profile for Chlorinated Dibenzo-p-Dioxins: December 1998. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

ATSDR 2001a Toxicological Profile for Di-n-Butyl Phthalate. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

ATSDR. 2001b. Draft guidance manual for preparation of an interaction profile. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

ATSDR 2002. Toxicological Profile for Di(2-ethylhexyl)phthalate. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

ATSDR. 2004a. Toxicological profile for polybrominated biphenyls and polybrominated diphenyl ethers (Update). Atlanta, GA: Agency for Toxic Substances and Disease Registry.

ATSDR. 2004b. Guidance manual for the assessment of joint toxic action of chemical mixtures. Atlanta, GA: Agency for Toxic Substances and Disease Registry.

ATSDR 2004c. Interaction profile for benzene, toluene, ethylbenzene, and xylenes (BTEX). Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service.

ATSDR 2004d. Public health assessment guidance manual. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.

ATSDR 2012. Addendum to the 1998 Toxicological Profile for Chlorinated Dibenzo-p-Dioxins. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry

Aylward LL, Hays SM. 2002. Temporal trends in human TCDD body burden: decreases over three decades and implications for exposure levels. J Expo Anal Environ Epidemiol 12:319-328.

Brown DJ, Van Overneire I, Goeyens L et al. 2004. Analysis of Ah receptor pathway activation by brominated flame retardants. Chemosphere 55(2004):1509-1518.

CDC (Centers for Disease Control and Prevention). 2006. National Health and Nutrition Examination Survey. Atlanta, GA: CDC/NCEH. Department of Health and Human Services.

Chen G and Bunce NJ. 2003. Polybrominated diphenyl ethers as Ah receptor agonists and antagonists. Toxicological Sciences 76:310-320 (2003).

Huang PC, Kuo PL, Guo YL, Liao PC, Lee CC. 2007. Association between urinary phthalate monesters and thyroid hormones in pregnant women. Hum Reprod. 2007 Oct;22(10):2715-22. Epub 2007 Aug 17.

Koch HM, Presuss R, Angerer J. 2006. Di(2-ethylhexyl)phthalate (DEHP): human metabolism and internal exposure-- an update and latest results. Int J Androl(1):155-65; discussion 181-5.

Meeker JD, Calafat AM, Hauser R. 2007. Di(2-ethylhexyl) phthalate metabolites may alter thyroid hormone levels in men. Environ Health Perspect 2007 Jul 115(7):1029-34.

Peters AK, van Londen K, Bergman A et al. 2004. Effects of polybrominated diphenyl ethers on basal and TCDD-induced Ethoxyresorufin activity and cytochrome P450-1A1 expression in MCF-7, HepG2, and H4IIE cells. Toxicological Sciences 82:488-496(2004).

Peters AK, Sanderson JT, Bergman A, Van den Berg, M. 2006a. Antagonism of TCDD-induced Ethoxyresorufin-O-deethylation activity by polybrominated diphenyl ethers (PBDEs) in primary cyomolgus monkey (Macaca fascicularis) hepatocytes. Toxicology Letters 164(2006):123-132.

Peters AK, Nijmeijer S, Gradin K. et al. 2006b. Interactions of polybrominated diphenyl ethers with the aryl hydrocarbon receptor pathway. Toxicological Sciences 92(1):133-142(2006).

Pohl, HR. 2009. Chemical mixtures: Validation of weight-of-evidence predictions for interactions. The 29th International Symposium on Halogenated Persistent Organic Pollutants. Aug 23-28, 2009, Beijing, China.

Rider CV, Furr JR, Wilson VS et al. 2010. Cumulative effects of in utero administration of mixtures of reproductive toxicants that disrupt common target tissues via diverse mechanisms of toxicity. International Journal Andrology 33:443-462.

Sanders JM, Burka LT, Smith CS et al. 2005. Differential expression of CYP1A1, 2B, and 3A genes in the F344 rat following exposure to a polybrominated diphenyl ether mixture or individual components. Toxicological Sciences 88(1):127-133 (2005).

Schecter A, Papke O, Chi Tung K et al. 2005. Polybrominated diphenyl ether flame retardants in the U.S. population: current levels, temporal trends, and comparison with dioxins, dibenzofurans, and polychlorinated biphenyls. Journal of Occupational and Environmental Medicine 47(3):199-211.

Tanida T, Warita K, Ishihara K. et al. 2009. Fetal and neonatal exposure to three typical environmental chemicals with different mechanisms of action: Mixed exposure to phenol, phthalate, and dioxin cancels the effects of sole exposure on mouse midbrain dopaminergic nuclei. Toxicology Letters 189: 40-47.

Tomaszewski, KA., Montgomery CA, and Melnick RL. 1988. Modulation of 2,3,7,8-Tetrachlorobenzo-p-dioxin toxicity in F344 rats by di(2-ethylhexyl)phthalate. Chem Biol Interactions 65(1988):205-222.

Van den Berg M, Birnbaum L, Denison M et al. 2006. The 2005 World Health Organization reevaluation of human and mammalian toxic equivalency factors for dioxins and dioxin-like compounds. Toxicological Sciences 93(2):223-241(2006).