

# Security- Widefield

El Paso County | Colorado



INFORMATION TO PROTECT OUR COMMUNITIES

## Per- and Polyfluoroalkyl Substances (PFAS) Exposure Assessment

### Appendix A, B, and C



National Center  
for Environmental Health  
Agency for Toxic Substances  
and Disease Registry

## Appendix A: Additional Tables

Table A1. Dust sample results from Security-Widefield EA compared to results from other U.S. studies (nanograms per gram)

PFAS	Security-Widefield EA			Fraser et al. (2013) Household Dust—MA*		Karásková et al. (2016) Household Dust—U.S.†		Wu et al. (2015) Household Dust—CA Homes with Young Children‡		Wu et al. (2015) Household Dust—CA Homes with Older Adults Only‡		Scher et al. (2018) Household Dust—MN§	
	RL	GM	Range	GM	Range	Median	Range	GM	Range	GM	Range	Median	Range
PFBS	0.449–4.43	3.25	ND–67.9	NA	4.98–4.98	0.9	<0.73 <sup>¶</sup> –2.6	–	–	–	–	<5	<5–58
PFPeS	0.452–4.46	NA	ND–28.0	–	–	–	–	–	–	–	–	–	–
PFHxS	0.449–4.43	3.53	ND–267	NA	6.05–430	8.7	1.4–84.4	3.47	ND**–7,490	3.77	ND**–1,050	18	<5–790
PFHpS	0.449–4.43	NA	ND–3.25	–	–	<0.42 <sup>¶</sup>	<0.42 <sup>¶</sup> –2.9	–	–	–	–	–	–
PFOS	0.449–4.43	12.2	2.89–96.0	26.9	14.1–280	14.1	5.7–239	29.0	ND**–6,670	34.6	ND**–1,040	67	8.4–2,000
PFDS	0.449–4.43	NA	ND–9.83	–	–	2.8	0.5–9.8	–	–	–	–	–	–
PFDoS	0.449–4.43	NA	ND–16.3	–	–	–	–	–	–	–	–	–	–
PFBA	1.80–17.8	11.0	ND–160	13.9	4.89–999	–	–	–	–	–	–	24	<5–200
PFPeA	0.899–8.86	NA	ND–10.6	NA	5.39–249	1.7	<0.76 <sup>¶</sup> –24.8	–	–	–	–	6.2	<5–66
PFHxA	0.542–4.43	6.54	1.38–34.2	8.65	4.85–1,380	6.5	2.5–190	–	–	–	–	29	5.4–240
PFHpA	0.449–4.43	3.51	ND–22.2	12.0	4.93–586	3.6	0.9–86.7	–	–	–	–	23	<5–260
PFOA	0.449–4.43	7.99	ND–65.1	23.7	5.71–894	9.0	2.9–318	41.4	ND**–2,360	45.0	ND**–728	51	9.9–970
PFNA	0.449–4.43	6.70	ND–36.8	10.9	6.21–1,420	3.9	1.1–62.9	13.3	ND**–1,910	14.7	ND**–883	26	<5–450
PFDA	0.449–4.43	3.92	ND–13.4	NA	6.97–26.8	1.8	0.4–64.0	8.51	ND**–2,520	7.76	ND**–355	13	<5–370
PFUnA	0.449–4.43	NA	ND–12.2	NA	10.8–39.4	1.2	<1.06 <sup>¶</sup> –13.1	–	–	–	–	7.2	ND–67
PFDoA	0.449–4.43	NA	ND–10.9	NA	5.09–13.3	0.6	<0.72 <sup>¶</sup> –9.0	–	–	–	–	8.2	ND–190
PFTTrA	0.449–4.43	NA	ND–5.10	NA	10.3–10.3	ND <sup>¶</sup>	ND <sup>¶</sup> –2.1	–	–	–	–	–	–
PFTA	0.449–4.43	NA	ND–8.31	NA	11.2–11.2	0.8	<1.15 <sup>¶</sup> –3.0	–	–	–	–	–	–

PFAS	Security-Widefield EA			Fraser et al. (2013) Household Dust—MA*		Karásková et al. (2016) Household Dust—U.S.†		Wu et al. (2015) Household Dust—CA Homes with Young Children‡		Wu et al. (2015) Household Dust—CA Homes with Older Adults Only‡		Scher et al. (2018) Household Dust—MN§	
	RL	GM	Range	GM	Range	Median	Range	GM	Range	GM	Range	Median	Range
PFOSA	0.449–4.43	NA	ND–3.13	–	–	–	–	–	–	–	–	–	–
N-MeFOSE	0.517–5.10	NA	ND–5.20	–	–	0.6	0.6–0.6	–	–	–	–	–	–
MeFOSAA	0.449–4.43	2.35	ND–38.7	–	–	–	–	–	–	–	–	–	–
N-MeFOSE	4.49–44.3	26.8	ND–1,440	NA	18–488	1.0	<0.57 <sup>¶</sup> –9.9	–	–	–	–	–	–
EtFOSAA	0.449–4.43	3.08	ND–12.9	–	–	–	–	–	–	–	–	–	–
N-EtFOSE	3.36–33.1	NA	ND–150	NA	12.2–3,280	<0.34 <sup>¶</sup>	<0.34 <sup>¶</sup> –93.9	–	–	–	–	–	–
FtS 6:2	1.62–16.0	NA	ND–54.7	–	–	–	–	–	–	–	–	–	–
FtS 8:2	1.80–17.8	NA	ND–12.6	–	–	–	–	–	–	–	–	–	–

RL = reporting limit, GM = geometric mean, ng/g = nanograms per gram, NA = not applicable (i.e., too few detected results to calculate a GM), ND = not detected, — = PFAS was not measured as part of the study

\* This study evaluated dust samples collected from homes, offices, and vehicles in the greater Boston, Massachusetts, area between January and March of 2009. This table presents results for dust samples collected in the main living areas of 30 homes.

† This study evaluated dust samples collected from living rooms and bedrooms from homes in Canada, the Czech Republic, and the United States during the spring and summer of 2013. The results presented in this table are from the 14 homes in the United States.

‡ As part of this study, dust samples were collected between 2007 and 2009 from carpet or area rugs in the main living areas of homes in California with and without young children residing in the home. This table presents results separately for dust samples collected in the 82 homes with young children and the 42 homes with older adults only.

§ As part of this study, dust samples were collected between July and September 2010 from 19 homes located in cities with PFAS-contaminated drinking water in Minnesota. Samples were collected at each home from an entryway to the yard as well as in an interior living space (e.g., family room, living room). The results presented in this table are for dust samples collected in interior living spaces only.

¶ Value was less than author-specified method detection limit. For this study, method detection limits varied because they were defined as mean concentration of procedural blanks plus three times the standard deviation of blank response. Values included in this table represent the upper bound of the method detection limit for a given PFAS, unless noted by “ND” (i.e., for PFTra). For PFTra, the upper bound method detection limit was greater than the maximum detected value. For PFTra, the method detection limits ranged from 0.48 to 2.32 ng/g.

\*\* Reporting limits for dust not specified in Wu et al. (2015).

**Table A2. Comparison values for PFAS measured in blood from other exposure assessments**

PFAS/Population	Reference	Geometric Mean for Blood (µg/L)
<b>PFHxS</b>		
Manufacturing workers, Decatur, AL	Olsen et al. 2003	180.0
<b>Security-Widefield EA<sup>†</sup></b>	This EA	10.6
Montgomery and Bucks Counties, PA	PA DOH 2019	6.6
Decatur, AL	ATSDR 2013	6.4
Little Hocking Water Association, OH	Frisbee et al. 2009	5.7*
Portsmouth, NH	NH DPHS 2016	4.1
Westhampton Beach/Quogue Area, NY	NYDOH 2019	3.0
General U.S. population (NHANES 1999/2000)	CDC 2019	2.1
General U.S. population (NHANES 2015/2016)	CDC 2019	1.2
<b>PFOS</b>		
Manufacturing workers, Decatur, AL	Olsen et al. 2003	941.0
Decatur, AL	ATSDR 2013	39.8
General U.S. population (NHANES 1999/2000)	CDC 2019	30.4
Little Hocking Water Association, OH	Frisbee et al. 2009	23.5*
Montgomery and Bucks Counties, PA	PA DOH 2019	10.2
Portsmouth, NH	NH DPHS 2016	8.6
Westhampton Beach/Quogue Area, NY	NYDOH 2019	6.6
<b>Security-Widefield EA<sup>†</sup></b>	This EA	6.2
General U.S. population (NHANES 2015/2016)	CDC 2019	4.7
<b>PFOA</b>		
Manufacturing workers, Decatur, AL	Olsen et al. 2003	899.0
Little Hocking Water Association, OH	Frisbee et al. 2009	227.6*
Decatur, AL	ATSDR 2013	16.3
General U.S. population (NHANES 1999/2000)	CDC 2019	5.2
Montgomery and Bucks Counties, PA	PA DOH 2019	3.1
Portsmouth, NH	NH DPHS 2016	3.1
<b>Security-Widefield EA<sup>†</sup></b>	This EA	2.1
General U.S. population (NHANES 2015/2016)	CDC 2019	1.6
Westhampton Beach/Quogue Area, NY	NYDOH 2019	1.5

µg/L = micrograms per liter

\* The study reported medians instead of geometric means.

<sup>†</sup> Unadjusted geometric means from the Security-Widefield EA are included in this table for comparison.

## Appendix B: Additional Background Statistics

As described in the main body of this report, all statistical analyses (e.g., correlations, geometric means, univariate linear regression models, multivariate linear regression models) were completed in SAS version 9.4 (SAS Institute, Cary, NC) following the methods outlined in the study protocol. Several key details on these methods are provided below.

- Consistent with NHANES methodology and per the EA protocol, all non-detect observations were substituted with a value equal to the LOD divided by the square root of 2. Geometric means were not reported for PFAS with 40% or more non-detect observations. Additional information on the effect of this substitution method is provided below.
- Geometric means, 95% confidence intervals around geometric means, and percentiles were calculated with the SURVEYMEANS procedure in SAS. In this procedure, percentiles are based on the population cumulative distribution function.
- Univariate and multivariate regression analyses were conducted with the SURVEYREG procedure in SAS. Multivariate regressions were conducted using a backwards stepwise approach. “Interactions” were only considered when there was a suspected relationship between two variables. Due to the skewed distribution of PFAS blood levels, log transformed ( $\log_{10}$ ) values were used as dependent variables in all linear regression analyses.
- For this EA, all eligible residents within the randomly selected households were invited to participate. This means a single household may have multiple participants. To account for the one-stage cluster sampling design used for this EA, household IDs were assigned to each participant. All statistics were calculated while accounting for clustering at the household level by including this household ID variable in a CLUSTER statement in SAS survey procedures. Additional information on the effect of clustering is provided below.
- A finite population correction was applied by including the total number of households in the sampling frame in a TOTAL statement in the SAS survey procedures. For this EA, a total of 10,783 households were identified within the sampling frame. A finite population correction corrects the standard errors when sampling without replacement from a finite population and is recommended when sample size is greater than 5% of the population being sampled.
- A p-value of less than 0.05 was used to identify statistically significant associations in regression models and 95% confidence limits were provided for all estimated geometric means.
- Age-adjusted statistics were calculated using the POSTSTRATA statement in the PROC SURVEYREG procedure in SAS. For age adjustments to the sampling frame population, the number of people in the sampling frame in each age category starting from “Under 10 years” and increasing by 5-year age intervals (10–14 years, 15–19 years, etc. through 80+) was calculated from census block data from 2010 and was used as poststratum totals (`_PSTOTAL_`). Similarly, for age-adjustments to the NHANES population, estimates of the U.S. population in each age category starting from 12-14 years and increasing by 5-year age intervals (15-20 years, 25-30 years etc. through 80+) were calculated.

### **Additional details on non-detect observations**

As noted, all results reported below the LOD were substituted with a value equal to the LOD divided by the square root of 2. For blood, all PFAS and all samples were reported from the laboratory with an LOD of 0.1  $\mu\text{g/L}$ , and non-detect observations were therefore substituted with a value equal to 0.071  $\mu\text{g/L}$ .

The same method was applied to urine results (LOD=0.1 µg/L) and dust (LOD varies by PFAS and sample); no summary statistics were computed for tap water for this EA due to low detection frequency.

The study protocol also notes that a sensitivity analysis of aggregate PFAS blood data should be performed using other statistical methods to account for censoring. More specifically, for datasets in which less than 50% of the data are censored (i.e., not detected), the Kaplan-Meier method should be used to calculate summary statistics; and for data sets with between 50% and 80% censored results, maximum likelihood estimation should be used. Only high sample percentiles should be reported for data sets with more than 80% censoring. Given that no nationally representative comparison values using these methods are available, results of this sensitivity analyses should only be used as a comparison to results obtained using the simpler substitution method described above.

Based on these criteria, ATSDR compared geometric means for all PFAS measured in blood (except Sb-PFOA) using the two alternate substitution methods. As shown in Table B1, there is little to no difference in geometric mean estimates when using these methods, and alternative substitution methods would therefore have no effect on the conclusions of this report. This is expected for these data due to the single censoring threshold for all PFAS and blood samples [Helsel 2009].

**Table B1. Comparison of geometric mean blood levels with various substitution methods**

PFAS	Geometric Mean Calculated with LOD/Square Root of 2 (µg/L)	Geometric Mean Calculated with Kaplan Meier Approach (µg/L)	Geometric Mean Calculated with Maximum Likelihood Estimation (µg/L)
PFHxS	10.63	10.63	10.70
n-PFOS	3.98	3.99	4.02
sm-PFOS	2.11	2.12	2.12
n-PFOA	2.04	2.05	2.05
sb-PFOA	NA*	NA*	NA*
PFNA	0.29	0.29	0.29
PFDA	0.12	0.14	0.12
PFUnA	0.09	0.11	0.09
MeFOSAA	0.13	0.15	0.13

LOD = limit of detection, µg/L = micrograms per liter, NA = not applicable

\* LOD does not meet the threshold set in EA protocol for sensitivity analyses (<20%).

### More details on precision and clustering for PFAS blood data

As noted in the study protocol, this investigation was designed to estimate mean concentrations of PFAS in blood for the sampling frame population, with a given level of precision. The target sample size for this EA was based on a desired precision of 15% and 5% level of significance. Table B2 presents the estimated precision for the mean of the log transformed (ln) PFAS concentrations. This was calculated as the difference between the upper confidence interval of ln(PFAS) and the mean ln(PFAS), divided by mean ln(PFAS). Precision estimates ranged from 3.9% to 16%. Except for Sm-PFOS, these values are all below the desired precision of 15% used to determine the target sample size for this EA. The collected data met the precision target specified in the EA protocol. Additional information on target precision is provided in the study protocol.

Note that throughout the main body of the report and Appendix C, geometric means are presented with 95% confidence intervals and regression modeling results are presented with p-values. These statistics provide further insight into the precision of those estimates.

To quantify the effect of clustering and to compare the results of this EA to the assumptions used to determine the target sample size for the EA (listed in the protocol), ATSDR calculated the intra-cluster correlation coefficient (ICC) and design effect for each PFAS that was detected in at least 60% of blood samples (Table B2). ICCs were estimated using variance components from the MIXED procedure in SAS. In brief, a mixed model was run for each PFAS while treating clusters (i.e., households) as a random effect. The ICC was calculated as the ratio of the variance attributable to the random effect (households) divided by the total of the random effect and error variances. The design effect was calculated using the DEFF option in the MODEL statement of the SURVEYREG procedure in SAS. This provides an estimate of the ratio of the actual variance to the variance computed under the assumption of simple random sampling. This information, along with the average number of study participants per house, was then used to calculate the effective sample size for each PFAS. This statistic provides an estimate of the sample size that would be required to achieve the same level of precision if a simple random sample design was used.

The target sample size for this EA was 395 people, based on (1) an ICC of 0.54 for PFOS calculated from data collected as part of biomonitoring study conducted by the New York State Department of Health and the Pennsylvania Department of Health, (2) a design effect of 2.1, and (3) an effective sample size of 188 people. Refer to the study protocol for more details on how these values were derived.

**Table B2. Statistics related to clustering in blood data (all participants)**

PFAS	Household ICC (Unitless)	Design Effect (Unitless)	Effective Sample Size (n)	Standard Deviation of ln(PFAS) (µg/L)	Precision of Mean ln(PFAS) (%)
PFHxS	0.50	1.55	224	1.38	6.2
PFOS	0.33	1.51	229	1.11	6.4
n-PFOS	0.36	1.52	227	1.15	8.8
Sm-PFOS	0.30	1.52	228	1.13	16.0
PFOA	0.26	1.30	267	0.85	11.7
n-PFOA	0.27	1.28	270	0.90	13.2
Sb-PFOA	NA*	NA*	NA*	NA*	NA*
PFNA	0.38	1.46	238	0.84	7.1
PFDA	0.59	1.73	200	0.78	3.9
PFUnA	NA*	NA*	NA*	NA*	NA*
MeFOSAA	0.81	1.73	200	0.97	5.1

µg/L = micrograms per liter, NA = not applicable

\* Per the protocol, geometric means were not calculated for PFAS detected in less than 60% of samples.

Blood ICCs for this EA ranged from 0.26 to 0.81, suggesting weak to strong correlation depending on the compound. The design effects ranged from 1.28 to 1.73, all of which are lower than the assumed design effect of 2.1. Effective sample size estimates ranged from 200 to 270. The design effect in this EA is smaller than that assumed in the protocol in part because of a smaller standard deviation of ln(PFAS)

(the protocol assumed a standard deviation of 1.63), and because of a smaller number of people per household. In this EA the average number of people per household was 1.8 (compared to 3.0, assumed in the protocol).

# Appendix C: PFAS Blood Levels by Demographics and Exposure Characteristics

This appendix provides geometric mean blood PFAS concentrations and 95% confidence intervals stratified by demographic or exposure characteristics for the six PFAS with detection frequencies above 60% (i.e., PFHxS, PFOS, PFOA, PFNA, PFDA, and MeFOSAA). Also included are univariate regressions, multivariate regressions, and box and whisker plots. For each regression, the outputs shown are coefficient estimates, p-values, and marginal effects. The coefficient represents the increase in PFAS blood levels (in units of  $\log_{10}[\mu\text{g/L}]$ ) per unit increase of the independent variable shown on the left side of the table for continuous variables, or when comparing to the reference category for categorical variables. The p-value indicates the significance of the results. Generally, p-values less than 0.05 indicate significant results. The marginal effect is the percent change in PFAS blood levels (in units of  $\mu\text{g/L}$ ) per unit increase of the continuous variables, or in comparison to the reference category for categorical variables.

## Contents

Table C1. Adult blood PFAS geometric means (GM), 95% lower confidence intervals (LCI), and 95% upper confidence intervals (UCI) in micrograms per liter <sup>*,†,‡</sup> .....	C4
Table C1 continued. Adult geometric means (GM), 95% lower confidence intervals (LCI), and 95% upper confidence intervals (UCI) in micrograms per liter <sup>*,†,‡</sup> .....	C8
Table C2. Child blood PFAS geometric means (GM), lower confidence intervals (LCI), and upper confidence intervals (UCI) in micrograms per liter <sup>*,†,‡</sup> .....	C12
Table C2 continued. Child blood PFAS geometric means (GM), lower confidence intervals (LCI), and upper confidence intervals (UCI) in micrograms per liter <sup>*,†,‡</sup> .....	C15
Table C3. Adult univariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME) <sup>*</sup> .....	C18
Table C3 continued. Adult univariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME) <sup>*</sup> .....	C22
Table C4. Child univariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME) .....	C26
Table C4 continued. Child univariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME) .....	C28
Table C5. PFHxS adult multivariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME) .....	C29
Table C6. PFHxS adult female multivariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME) .....	C29
Table C7. PFHxS adult male multivariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME) .....	C30
Table C8. PFOS adult multivariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME) .....	C30
Table C9. PFOS adult female multivariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME) .....	C31

Table C10. PFOS adult male multivariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME) .....	C31
Table C11. PFOA adult multivariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME) .....	C32
Table C12. PFOA adult male multivariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME) .....	C32
Figure C1. Boxplot of adult blood (serum) PFAS concentrations by age .....	C33
Figure C2. Boxplot of adult blood (serum) PFAS concentrations by sex.....	C33
Figure C3. Boxplot of adult blood (serum) PFAS concentrations by race and ethnicity .....	C34
Figure C4. Boxplot of adult blood (serum) PFAS concentrations by body mass index .....	C34
Figure C5. Boxplot of adult blood (serum) PFAS concentrations by years in current home .....	C35
Figure C6. Boxplot of adult blood (serum) PFAS concentrations by years in sampling frame (past 20 years).....	C35
Figure C7. Boxplot of adult blood (serum) PFAS concentrations by cups of tap water drank at home ...	C36
Figure C8. Boxplot of adult blood (serum) PFAS concentrations by water filter type.....	C36
Figure C9. Boxplot of adult blood (serum) PFAS concentrations by kidney disease history .....	C37
Figure C10. Boxplot of adult blood (serum) PFAS concentrations by blood donation frequency .....	C37
Figure C11. Boxplot of adult blood (serum) PFAS concentrations by home cleaning frequency .....	C38
Figure C12. Boxplot of adult blood (serum) PFAS concentrations by stain-resistant product use.....	C38
Figure C13. Boxplot of adult blood (serum) PFAS concentrations by frequency of contact with soil.....	C39
Figure C14. Boxplot of adult blood (serum) PFAS concentrations by local fruit and vegetable consumption .....	C39
Figure C15. Boxplot of adult blood (serum) PFAS concentrations by local fish consumption.....	C40
Figure C16. Boxplot of adult blood (serum) PFAS concentrations by local milk consumption .....	C40
Figure C17. Boxplot of adult blood (serum) PFAS concentrations by fast food consumption frequency .....	C41
Figure C18. Boxplot of adult blood (serum) PFAS concentrations by presence of carpet in home .....	C41
Figure C19. Boxplot of adult blood (serum) PFAS concentrations by occupational exposure .....	C42
Figure C20. Boxplot of adult female blood (serum) PFAS concentrations by breastfeeding history .....	C42
Figure C21. Boxplot of adult female blood (serum) PFAS concentrations by breastfeeding duration.....	C43
Figure C22. Boxplot of adult female blood (serum) PFAS concentrations by biological children variable.....	C43
Figure C23. Boxplot of adult female blood (serum) PFAS concentrations by number of biological children .....	C44
Figure C24. Boxplot of child blood (serum) PFAS concentrations by age.....	C44
Figure C25. Boxplot of child blood (serum) PFAS concentrations by sex .....	C45
Figure C26. Boxplot of child blood (serum) PFAS concentrations by body mass index.....	C45

Figure C27. Boxplot of child blood (serum) PFAS concentrations by race and ethnicity combined ..... C46

Figure C28. Boxplot of child blood (serum) PFAS concentrations by water consumption at current home ..... C46

Figure C29. Boxplot of child blood (serum) PFAS concentrations by water consumption at school ..... C47

Figure C30. Boxplot of child blood (serum) PFAS concentrations by length of residency in sampling frame..... C47

Figure C31. Boxplot of child blood (serum) PFAS concentrations by frequency of contact with soil ..... C48

Figure C32. Boxplot of child blood (serum) PFAS concentrations by local fruit and vegetable consumption ..... C48

Figure C33. Boxplot of child blood (serum) PFAS concentrations by drinking formula reconstituted with tap water..... C49

Figure C34. Boxplot of child blood (serum) PFAS concentrations by duration of drinking formula reconstituted with tap water ..... C49

Figure C35. Boxplot of child blood (serum) PFAS concentrations by history of breastfeeding..... C50

Figure C36. Boxplot of child blood (serum) PFAS concentrations by breastfeeding duration ..... C50

**Table C1. Adult blood PFAS geometric means (GM), 95% lower confidence intervals (LCI), and 95% upper confidence intervals (UCI) in micrograms per liter<sup>\*,†,‡</sup>**

Variable	Category	Frequency <sup>§</sup>	PFHxS			PFOS			PFOA		
			GM	LCI	UCI	GM	LCI	UCI	GM	LCI	UCI
<b>All Adults</b>		318	<b>11.50</b>	10.03	13.19	<b>6.59</b>	5.88	7.38	<b>2.25</b>	2.06	2.45
Age (years)	18 to <50	97	<b>6.35</b>	5.03	8.03	<b>4.53</b>	3.71	5.54	<b>1.67</b>	1.46	1.91
	50+	221	<b>14.93</b>	12.85	17.34	<b>7.76</b>	6.90	8.74	<b>2.56</b>	2.32	2.82
Sex	Female	175	<b>10.63</b>	8.96	12.61	<b>5.62</b>	4.86	6.49	<b>2.14</b>	1.90	2.41
	Male	143	<b>12.67</b>	10.65	15.08	<b>8.01</b>	6.92	9.27	<b>2.38</b>	2.12	2.67
Body Mass Index (kilograms per square meter)	<25	72	<b>12.90</b>	9.36	17.78	<b>7.14</b>	5.69	8.96	<b>2.38</b>	1.92	2.95
	25 to <30	113	<b>10.30</b>	8.41	12.62	<b>6.19</b>	5.20	7.36	<b>2.07</b>	1.84	2.32
	30 to <35	77	<b>13.51</b>	10.70	17.06	<b>7.34</b>	5.92	9.11	<b>2.48</b>	2.09	2.94
	35+	53	<b>10.30</b>	7.74	13.71	<b>5.95</b>	4.75	7.45	<b>2.21</b>	1.79	2.73
Race and ethnicity combined	White alone, not Hispanic	224	<b>12.24</b>	10.48	14.29	<b>6.65</b>	5.88	7.51	<b>2.32</b>	2.11	2.55
	Not White, or Hispanic	87	<b>9.91</b>	7.68	12.80	<b>6.01</b>	4.76	7.60	<b>2.00</b>	1.66	2.40
Length of residence at current address (years)	<10	80	<b>4.98</b>	3.79	6.54	<b>4.26</b>	3.47	5.23	<b>1.60</b>	1.37	1.87
	10 to <20	82	<b>12.57</b>	10.13	15.58	<b>6.25</b>	5.01	7.79	<b>2.21</b>	1.89	2.59
	20 to <30	77	<b>13.53</b>	10.49	17.44	<b>7.93</b>	6.31	9.95	<b>2.62</b>	2.20	3.13
	30+	79	<b>20.92</b>	17.10	25.59	<b>9.04</b>	7.73	10.56	<b>2.77</b>	2.41	3.19
Total length of residence in sampling frame over the past 20 years (years)	<10	56	<b>4.23</b>	3.24	5.53	<b>4.25</b>	3.57	5.07	<b>1.61</b>	1.34	1.93
	10 to <15	34	<b>10.75</b>	7.76	14.89	<b>6.73</b>	4.54	9.98	<b>2.08</b>	1.54	2.80
	15 to 20	228	<b>14.86</b>	12.76	17.30	<b>7.31</b>	6.40	8.35	<b>2.47</b>	2.23	2.72
Current and primary source of drinking water	Public water system	217	<b>12.92</b>	11.03	15.13	<b>6.98</b>	6.09	8.00	<b>2.37</b>	2.13	2.63
	Bottled water	101	<b>8.97</b>	7.05	11.41	<b>5.81</b>	4.85	6.97	<b>2.01</b>	1.74	2.32
Public Water Supply	Other	5	NA	NA	NA	NA	NA	NA	NA	NA	NA
	Security WD	203	<b>11.58</b>	9.73	13.78	<b>5.97</b>	5.20	6.87	<b>2.19</b>	1.97	2.44
	Widefield WSD	109	<b>11.38</b>	9.04	14.31	<b>7.97</b>	6.63	9.57	<b>2.38</b>	2.05	2.76
	Security Mobile Home Park	1	NA	NA	NA	NA	NA	NA	NA	NA	NA

Variable	Category	Frequency <sup>§</sup>	PFHxS			PFOS			PFOA		
			GM	LCI	UCI	GM	LCI	UCI	GM	LCI	UCI
Tap water consumption at current home (average cups per day)	0	45	<b>6.45</b>	4.54	9.17	<b>4.58</b>	3.51	5.97	<b>1.63</b>	1.32	2.03
	>0 to <2	13	<b>11.56</b>	6.35	21.02	<b>6.36</b>	4.51	8.97	<b>2.06</b>	1.41	3.00
	2 to <4	40	<b>13.47</b>	9.89	18.35	<b>8.05</b>	5.92	10.96	<b>2.36</b>	1.87	2.98
	4 to <6	60	<b>10.99</b>	8.64	13.99	<b>6.79</b>	5.66	8.16	<b>2.31</b>	1.98	2.71
	6 to <8	44	<b>10.88</b>	8.18	14.48	<b>5.30</b>	4.02	6.98	<b>2.18</b>	1.77	2.68
	8+	114	<b>14.57</b>	11.48	18.49	<b>7.65</b>	6.27	9.33	<b>2.54</b>	2.19	2.93
Current use of filter or treatment device for tap water at home	None, no filter or treatment device	83	<b>13.18</b>	10.31	16.84	<b>7.70</b>	6.46	9.18	<b>2.28</b>	1.92	2.71
	None, drink bottled water only	45	<b>6.33</b>	4.38	9.13	<b>4.38</b>	3.34	5.74	<b>1.55</b>	1.27	1.89
	Use at least one filter or treatment device	190	<b>12.49</b>	10.55	14.79	<b>6.78</b>	5.83	7.88	<b>2.44</b>	2.20	2.71
History of kidney disease	No	292	<b>11.48</b>	9.98	13.21	<b>6.51</b>	5.80	7.31	<b>2.27</b>	2.08	2.48
	Yes	20	<b>9.90</b>	5.81	16.87	<b>7.61</b>	4.57	12.66	<b>2.24</b>	1.62	3.10
Frequency of blood donation	Never/rarely	305	<b>11.47</b>	9.96	13.21	<b>6.52</b>	5.79	7.33	<b>2.24</b>	2.05	2.45
	Once or more a year	12	<b>12.59</b>	7.29	21.75	<b>9.02</b>	6.35	12.83	<b>2.41</b>	1.68	3.44
Frequency of house cleaning	A few times per month or less	186	<b>12.65</b>	10.72	14.94	<b>6.30</b>	5.48	7.24	<b>2.22</b>	2.00	2.48
	Three times per week or more	132	<b>10.06</b>	8.10	12.49	<b>7.01</b>	5.86	8.39	<b>2.28</b>	1.99	2.61
Frequency of stain-resistant product use	Never	285	<b>11.77</b>	10.20	13.60	<b>6.55</b>	5.80	7.39	<b>2.28</b>	2.09	2.49
	Rarely or more frequently	33	<b>9.41</b>	6.17	14.36	<b>6.97</b>	5.23	9.29	<b>1.97</b>	1.41	2.75
Frequency of direct contact with soil at locations within the sampling frame	A few times per year or less	105	<b>10.57</b>	8.19	13.62	<b>6.42</b>	5.19	7.95	<b>2.11</b>	1.78	2.49
	A few times per month	109	<b>13.42</b>	10.87	16.58	<b>6.83</b>	5.80	8.05	<b>2.53</b>	2.21	2.89
	Three times per week or more	104	<b>10.66</b>	8.58	13.25	<b>6.51</b>	5.34	7.94	<b>2.12</b>	1.84	2.45

Variable	Category	Frequency <sup>s</sup>	PFHxS			PFOS			PFOA		
			GM	LCI	UCI	GM	LCI	UCI	GM	LCI	UCI
Consumption of fruits and vegetables from locations within the sampling frame	No	161	<b>10.90</b>	8.99	13.23	<b>5.97</b>	5.07	7.03	<b>2.13</b>	1.89	2.41
	Yes	154	<b>12.20</b>	10.17	14.64	<b>7.41</b>	6.37	8.62	<b>2.38</b>	2.11	2.70
Consumption of local fish (i.e., fish caught within the sampling frame)	No	306	<b>11.51</b>	10.01	13.23	<b>6.54</b>	5.83	7.33	<b>2.25</b>	2.05	2.45
	Yes	11	<b>10.42</b>	5.62	19.30	<b>7.83</b>	4.60	13.31	<b>2.27</b>	1.61	3.22
Frequency of local milk consumption (i.e., milk from animals within the sampling frame)	Never	306	<b>11.51</b>	9.99	13.27	<b>6.74</b>	6.02	7.55	<b>2.28</b>	2.08	2.49
	Rarely or more frequently	5	NA	NA	NA	NA	NA	NA	NA	NA	NA
Frequency of fast food consumption	Three times per week or more	80	<b>9.94</b>	7.61	12.98	<b>5.34</b>	4.23	6.73	<b>1.97</b>	1.64	2.38
	A few times per month	205	<b>11.72</b>	9.89	13.88	<b>6.96</b>	6.12	7.90	<b>2.32</b>	2.10	2.56
	A few times per year or less	33	<b>14.63</b>	9.84	21.76	<b>7.83</b>	5.70	10.76	<b>2.55</b>	1.96	3.30
Presence of carpeting in bedroom, living room, or kitchen	No	85	<b>13.01</b>	10.59	16.00	<b>7.39</b>	5.98	9.13	<b>2.40</b>	2.11	2.73
	Yes	233	<b>11.00</b>	9.30	13.00	<b>6.32</b>	5.54	7.20	<b>2.19</b>	1.97	2.44
Occupational exposures (count of jobs with potential PFAS exposures)	None	277	<b>12.11</b>	10.45	14.04	<b>6.64</b>	5.85	7.54	<b>2.24</b>	2.04	2.46
	One or more	41	<b>8.13</b>	6.15	10.74	<b>6.24</b>	5.10	7.63	<b>2.30</b>	1.86	2.86

Variable	Category	Frequency <sup>§</sup>	PFHxS			PFOS			PFOA		
			GM	LCI	UCI	GM	LCI	UCI	GM	LCI	UCI
<b>Females only</b>											
Biological children	No	27	<b>7.82</b>	5.69	10.74	<b>4.90</b>	3.89	6.17	<b>1.84</b>	1.42	2.39
	Yes	147	<b>11.32</b>	9.35	13.69	<b>5.76</b>	4.91	6.76	<b>2.21</b>	1.93	2.53
Number of biological children	0	27	<b>7.82</b>	5.69	10.74	<b>4.90</b>	3.89	6.17	<b>1.84</b>	1.42	2.39
	1	33	<b>8.39</b>	5.94	11.84	<b>4.91</b>	3.73	6.45	<b>1.88</b>	1.48	2.40
	2	60	<b>11.75</b>	8.92	15.49	<b>5.64</b>	4.40	7.23	<b>2.27</b>	1.84	2.80
	3+	54	<b>13.03</b>	9.27	18.31	<b>6.50</b>	4.97	8.50	<b>2.37</b>	1.88	2.98
Breastfeeding or previously breastfed children	No	83	<b>10.63</b>	8.53	13.24	<b>5.56</b>	4.71	6.57	<b>2.13</b>	1.80	2.51
	Yes	91	<b>10.74</b>	8.29	13.90	<b>5.67</b>	4.52	7.10	<b>2.17</b>	1.82	2.59
Total duration of breastfeeding for all children (months)	0	84	<b>10.52</b>	8.46	13.08	<b>5.56</b>	4.72	6.55	<b>2.12</b>	1.79	2.50
	>0 to <6	21	<b>17.72</b>	11.51	27.26	<b>7.23</b>	4.57	11.45	<b>2.44</b>	1.77	3.37
	6 to <12	19	<b>7.47</b>	3.92	14.25	<b>4.36</b>	2.56	7.42	<b>1.91</b>	1.34	2.73
	12 to <18	17	<b>10.02</b>	5.16	19.48	<b>6.09</b>	4.08	9.08	<b>2.27</b>	1.49	3.44
	18+	34	<b>9.99</b>	6.95	14.34	<b>5.45</b>	3.88	7.65	<b>2.12</b>	1.61	2.79

\* Several variables that were collected in the questionnaire are not included in these tables. These variables may not be included because they did not have sufficient variability or were not associated with PFAS blood concentrations in preliminary analyses. These variables include full-time vs. part-time residence, behavior change questions, and occupational history in specific industries.

+ Geometric means and confidence levels are not shown for categories with fewer than 10 responses.

\* Detection limits for all PFAS are 0.1 micrograms per liter (µg/L).

§ Some frequency counts may not sum to the total because of missing values. Some variable categories that were presented in the questionnaire were collapsed into larger variable categories.

Table C1 continued. Adult geometric means (GM), 95% lower confidence intervals (LCI), and 95% upper confidence intervals (UCI) in micrograms per liter<sup>\*,†,‡</sup>

Variable	Category	Frequency <sup>§</sup>	PFNA			PFDA			MeFOSAA		
			GM	LCI	UCI	GM	LCI	UCI	GM	LCI	UCI
<b>All Adults</b>		318	<b>0.30</b>	0.27	0.32	<b>0.12</b>	0.11	0.13	<b>0.14</b>	0.12	0.15
Age (years)	18 to <50	97	<b>0.23</b>	0.20	0.26	<b>0.12</b>	0.10	0.14	<b>0.11</b>	0.09	0.12
	50+	221	<b>0.33</b>	0.30	0.37	<b>0.12</b>	0.11	0.14	<b>0.16</b>	0.14	0.17
Sex	Female	175	<b>0.28</b>	0.24	0.32	<b>0.13</b>	0.11	0.14	<b>0.14</b>	0.12	0.15
	Male	143	<b>0.32</b>	0.29	0.35	<b>0.12</b>	0.11	0.13	<b>0.14</b>	0.12	0.16
Body Mass Index (kilograms per square meter)	<25	72	<b>0.31</b>	0.26	0.39	<b>0.13</b>	0.11	0.16	<b>0.17</b>	0.13	0.21
	25 to <30	113	<b>0.29</b>	0.25	0.33	<b>0.12</b>	0.11	0.14	<b>0.14</b>	0.12	0.16
	30 to <35	77	<b>0.33</b>	0.28	0.38	<b>0.13</b>	0.11	0.15	<b>0.13</b>	0.11	0.16
	35+	53	<b>0.25</b>	0.21	0.31	<b>0.11</b>	0.09	0.12	<b>0.12</b>	0.10	0.14
Race and ethnicity combined	White alone, not Hispanic	224	<b>0.28</b>	0.26	0.31	<b>0.12</b>	0.11	0.13	<b>0.14</b>	0.12	0.16
	Not White, or Hispanic	87	<b>0.31</b>	0.26	0.38	<b>0.14</b>	0.12	0.17	<b>0.13</b>	0.11	0.16
Length of residence at current address (years)	<10	80	<b>0.28</b>	0.23	0.33	<b>0.13</b>	0.11	0.16	<b>0.10</b>	0.09	0.12
	10 to <20	82	<b>0.28</b>	0.23	0.34	<b>0.13</b>	0.11	0.15	<b>0.13</b>	0.11	0.15
	20 to <30	77	<b>0.34</b>	0.29	0.39	<b>0.13</b>	0.11	0.15	<b>0.15</b>	0.13	0.19
	30+	79	<b>0.29</b>	0.25	0.34	<b>0.11</b>	0.10	0.12	<b>0.19</b>	0.15	0.24
Total length of residence in sampling frame over the past 20 years (years)	<10	56	<b>0.29</b>	0.23	0.37	<b>0.14</b>	0.12	0.17	<b>0.10</b>	0.09	0.12
	10 to <15	34	<b>0.29</b>	0.22	0.39	<b>0.15</b>	0.11	0.20	<b>0.10</b>	0.08	0.13
	15 to 20	228	<b>0.30</b>	0.27	0.33	<b>0.12</b>	0.11	0.13	<b>0.15</b>	0.14	0.17
Current and primary source of drinking water	Public water system	217	<b>0.29</b>	0.27	0.33	<b>0.12</b>	0.11	0.14	<b>0.13</b>	0.12	0.15
	Bottled water	101	<b>0.30</b>	0.25	0.35	<b>0.12</b>	0.10	0.14	<b>0.15</b>	0.12	0.18

Variable	Category	Frequency <sup>§</sup>	PFNA			PFDA			MeFOSAA		
			GM	LCI	UCI	GM	LCI	UCI	GM	LCI	UCI
Public Water Supply	Other	5	NA	NA	NA	NA	NA	NA	NA	NA	NA
	Security WD	203	<b>0.29</b>	0.26	0.32	<b>0.12</b>	0.11	0.14	<b>0.13</b>	0.12	0.15
	Widefield WSD	109	<b>0.31</b>	0.26	0.36	<b>0.12</b>	0.10	0.14	<b>0.15</b>	0.12	0.18
	Security Mobile Home Park	1	NA	NA	NA	NA	NA	NA	NA	NA	NA
Tap water consumption at current home (average cups per day)	0	45	<b>0.27</b>	0.22	0.33	<b>0.11</b>	0.10	0.14	<b>0.12</b>	0.09	0.16
	>0 to <2	13	<b>0.33</b>	0.24	0.46	<b>0.14</b>	0.10	0.20	<b>0.11</b>	0.08	0.14
	2 to <4	40	<b>0.33</b>	0.26	0.41	<b>0.13</b>	0.10	0.16	<b>0.15</b>	0.12	0.20
	4 to <6	60	<b>0.31</b>	0.26	0.36	<b>0.12</b>	0.10	0.14	<b>0.14</b>	0.12	0.17
	6 to <8	44	<b>0.27</b>	0.22	0.33	<b>0.11</b>	0.10	0.13	<b>0.14</b>	0.11	0.18
	8+	114	<b>0.30</b>	0.26	0.35	<b>0.13</b>	0.11	0.15	<b>0.14</b>	0.12	0.17
Current use of filter or treatment device for tap water at home	None, no filter or treatment device	83	<b>0.30</b>	0.26	0.35	<b>0.12</b>	0.10	0.14	<b>0.15</b>	0.12	0.19
	None, drink bottled water only	45	<b>0.26</b>	0.20	0.34	<b>0.12</b>	0.10	0.14	<b>0.12</b>	0.09	0.16
	Use at least one filter or treatment device	190	<b>0.30</b>	0.27	0.34	<b>0.13</b>	0.11	0.14	<b>0.14</b>	0.12	0.15
History of kidney disease	No	292	<b>0.29</b>	0.27	0.32	<b>0.12</b>	0.11	0.13	<b>0.14</b>	0.12	0.15
	Yes	20	<b>0.36</b>	0.26	0.49	<b>0.14</b>	0.11	0.18	<b>0.17</b>	0.12	0.24
Frequency of blood donation	Never/rarely	305	<b>0.30</b>	0.27	0.32	<b>0.12</b>	0.11	0.13	<b>0.14</b>	0.12	0.15
	Once or more a year	12	<b>0.33</b>	0.23	0.46	<b>0.12</b>	0.09	0.17	<b>0.18</b>	0.09	0.35
Frequency of house cleaning	A few times per month or less	186	<b>0.28</b>	0.25	0.31	<b>0.11</b>	0.10	0.12	<b>0.14</b>	0.12	0.16
	Three times per week or more	132	<b>0.32</b>	0.28	0.37	<b>0.14</b>	0.12	0.16	<b>0.14</b>	0.12	0.16
Frequency of stain-resistant product use	Never	285	<b>0.29</b>	0.26	0.32	<b>0.12</b>	0.11	0.13	<b>0.13</b>	0.12	0.15
	Rarely or more frequently	33	<b>0.37</b>	0.28	0.50	<b>0.15</b>	0.12	0.19	<b>0.20</b>	0.14	0.29

Variable	Category	Frequency <sup>§</sup>	PFNA			PFDA			MeFOSAA		
			GM	LCI	UCI	GM	LCI	UCI	GM	LCI	UCI
Frequency of direct contact with soil at locations within the sampling frame	A few times per year or less	105	<b>0.28</b>	0.24	0.33	<b>0.12</b>	0.10	0.13	<b>0.14</b>	0.11	0.16
	A few times per month	109	<b>0.33</b>	0.29	0.37	<b>0.13</b>	0.11	0.14	<b>0.15</b>	0.13	0.17
	Three times per week or more	104	<b>0.28</b>	0.24	0.33	<b>0.13</b>	0.11	0.15	<b>0.13</b>	0.11	0.15
Consumption of fruits and vegetables from locations within the sampling frame	No	161	<b>0.28</b>	0.24	0.31	<b>0.11</b>	0.10	0.12	<b>0.14</b>	0.12	0.16
	Yes	154	<b>0.32</b>	0.28	0.36	<b>0.14</b>	0.12	0.16	<b>0.14</b>	0.12	0.16
Consumption of local fish (i.e., fish caught within the sampling frame)	No	306	<b>0.29</b>	0.27	0.32	<b>0.12</b>	0.11	0.13	<b>0.14</b>	0.13	0.15
	Yes	11	<b>0.37</b>	0.26	0.51	<b>0.14</b>	0.09	0.24	<b>0.11</b>	0.07	0.17
Frequency of local milk consumption (i.e., milk from animals within the sampling frame)	Never	306	<b>0.30</b>	0.27	0.33	<b>0.12</b>	0.11	0.13	<b>0.14</b>	0.13	0.15
	Rarely or more frequently	5	NA	NA	NA	NA	NA	NA	NA	NA	NA
Frequency of fast food consumption	Three times per week or more	80	<b>0.26</b>	0.22	0.31	<b>0.11</b>	0.10	0.13	<b>0.11</b>	0.10	0.14
	A few times per month	205	<b>0.31</b>	0.27	0.34	<b>0.13</b>	0.11	0.14	<b>0.15</b>	0.13	0.17
	A few times per year or less	33	<b>0.32</b>	0.25	0.40	<b>0.13</b>	0.11	0.17	<b>0.14</b>	0.10	0.18
Presence of carpeting in bedroom, living room, or kitchen	No	85	<b>0.29</b>	0.25	0.34	<b>0.12</b>	0.10	0.14	<b>0.12</b>	0.10	0.14
	Yes	233	<b>0.30</b>	0.27	0.33	<b>0.12</b>	0.11	0.14	<b>0.14</b>	0.13	0.16

Variable	Category	Frequency <sup>§</sup>	PFNA			PFDA			MeFOSAA		
			GM	LCI	UCI	GM	LCI	UCI	GM	LCI	UCI
Occupational exposures (count of jobs with potential PFAS exposures)	None	277	<b>0.29</b>	0.26	0.32	<b>0.12</b>	0.11	0.13	<b>0.14</b>	0.13	0.16
	One or more	41	<b>0.36</b>	0.30	0.43	<b>0.12</b>	0.10	0.15	<b>0.13</b>	0.10	0.17
<b>Females only</b>											
Biological children	No	27	<b>0.28</b>	0.20	0.39	<b>0.13</b>	0.10	0.18	<b>0.11</b>	0.09	0.14
	Yes	147	<b>0.28</b>	0.24	0.32	<b>0.12</b>	0.11	0.14	<b>0.14</b>	0.12	0.16
Number of biological children	0	27	<b>0.28</b>	0.20	0.39	<b>0.13</b>	0.10	0.18	<b>0.11</b>	0.09	0.14
	1	33	<b>0.28</b>	0.22	0.37	<b>0.13</b>	0.10	0.16	<b>0.16</b>	0.12	0.21
	2	60	<b>0.29</b>	0.24	0.35	<b>0.13</b>	0.11	0.16	<b>0.12</b>	0.10	0.14
	3+	54	<b>0.26</b>	0.21	0.33	<b>0.12</b>	0.10	0.14	<b>0.15</b>	0.12	0.19
Breastfeeding or previously breastfed children	No	83	<b>0.29</b>	0.24	0.34	<b>0.13</b>	0.11	0.15	<b>0.14</b>	0.12	0.17
	Yes	91	<b>0.27</b>	0.23	0.32	<b>0.13</b>	0.11	0.14	<b>0.13</b>	0.11	0.15
Total duration of breastfeeding for all children (months)	0	84	<b>0.28</b>	0.24	0.34	<b>0.13</b>	0.11	0.14	<b>0.14</b>	0.12	0.17
	>0 to <6	21	<b>0.30</b>	0.21	0.42	<b>0.12</b>	0.09	0.14	<b>0.15</b>	0.12	0.19
	6 to <12	19	<b>0.28</b>	0.19	0.41	<b>0.13</b>	0.10	0.18	<b>0.13</b>	0.09	0.18
	12 to <18	17	<b>0.26</b>	0.18	0.36	<b>0.14</b>	0.10	0.19	<b>0.13</b>	0.09	0.18
	18+	34	<b>0.26</b>	0.20	0.34	<b>0.12</b>	0.10	0.15	<b>0.12</b>	0.10	0.16

\* Several variables that were collected in the questionnaire are not included in these tables. These variables may not be included because they did not have sufficient variability or were not associated with PFAS blood concentrations in preliminary analyses. These variables include full-time vs. part-time residence, behavior change questions, and occupational history in specific industries.

<sup>†</sup> Geometric means and confidence levels are not shown for categories with fewer than 10 responses.

<sup>\*</sup> Detection limits for all PFAS are 0.1 micrograms per liter (µg/L).

<sup>§</sup> Some frequency counts may not sum to the total because of missing values. Some variable categories that were presented in the questionnaire were collapsed into larger variable categories.

Table C2. Child blood PFAS geometric means (GM), lower confidence intervals (LCI), and upper confidence intervals (UCI) in micrograms per liter<sup>\*,†,‡</sup>

Variable	Category	Frequency <sup>§</sup>	PFHxS			PFOS			PFOA		
			GM	LCI	UCI	GM	LCI	UCI	GM	LCI	UCI
<b>All Children</b>		28	<b>4.35</b>	3.09	6.13	<b>3.21</b>	2.53	4.06	<b>1.24</b>	1.07	1.45
Age (years)	3 to <12	9	NA	NA	NA	NA	NA	NA	NA	NA	NA
	12 to <18	19	<b>4.24</b>	2.55	7.07	<b>3.37</b>	2.53	4.50	<b>1.19</b>	0.97	1.45
Sex	Female	18	<b>4.54</b>	3.07	6.70	<b>3.04</b>	2.23	4.14	<b>1.25</b>	0.99	1.57
	Male	10	<b>4.04</b>	1.87	8.74	<b>3.53</b>	2.35	5.31	<b>1.24</b>	0.94	1.63
Body mass index (kilograms per square meter)	<15	1	NA	NA	NA	NA	NA	NA	NA	NA	NA
	15 to 20	9	NA	NA	NA	NA	NA	NA	NA	NA	NA
	20 to <25	14	<b>3.67</b>	2.00	6.75	<b>2.99</b>	2.16	4.15	<b>1.25</b>	0.94	1.66
	25+	3	NA	NA	NA	NA	NA	NA	NA	NA	NA
Birth order	First born	9	NA	NA	NA	NA	NA	NA	NA	NA	NA
	Second born	12	<b>4.10</b>	2.62	6.41	<b>3.32</b>	2.33	4.72	<b>1.15</b>	0.95	1.39
	Third born	6	NA	NA	NA	NA	NA	NA	NA	NA	NA
	Fifth born	1	NA	NA	NA	NA	NA	NA	NA	NA	NA
Race and ethnicity combined	White, not Hispanic	15	<b>4.01</b>	2.23	7.20	<b>3.16</b>	2.19	4.55	<b>1.21</b>	0.92	1.59
	Not White, or Hispanic	12	<b>4.77</b>	2.86	7.97	<b>3.44</b>	2.44	4.85	<b>1.31</b>	1.10	1.56
Water consumption at current home (average cups per day)	0 to <2	8	NA	NA	NA	NA	NA	NA	NA	NA	NA
	2 to <4	5	NA	NA	NA	NA	NA	NA	NA	NA	NA
	4+	15	<b>4.74</b>	2.99	7.52	<b>3.92</b>	2.91	5.29	<b>1.28</b>	1.03	1.59
Water consumption at school (average cups per day)	0 to <1	3	NA	NA	NA	NA	NA	NA	NA	NA	NA
	1 to <2	5	NA	NA	NA	NA	NA	NA	NA	NA	NA
	2 to <3	8	NA	NA	NA	NA	NA	NA	NA	NA	NA
	3+	12	<b>4.67</b>	3.05	7.16	<b>2.97</b>	2.19	4.03	<b>1.16</b>	0.91	1.46

Variable	Category	Frequency <sup>s</sup>	PFHxS			PFOS			PFOA		
			GM	LCI	UCI	GM	LCI	UCI	GM	LCI	UCI
Length of residency in sampling frame (years)	<6	3	NA	NA	NA	NA	NA	NA	NA	NA	NA
	6 to <12	12	<b>3.81</b>	1.89	7.70	<b>3.03</b>	2.10	4.36	<b>1.30</b>	0.95	1.78
	12 to <18	13	<b>5.39</b>	3.84	7.57	<b>3.45</b>	2.53	4.71	<b>1.22</b>	0.98	1.51
Frequency of direct contact with soil at locations within the sampling frame	A few times per year or less	4	NA	NA	NA	NA	NA	NA	NA	NA	NA
	A few times per month	11	<b>4.55</b>	2.77	7.49	<b>3.69</b>	2.61	5.21	<b>1.26</b>	0.94	1.68
	Three times per week or more	13	<b>4.36</b>	2.80	6.78	<b>2.79</b>	2.03	3.82	<b>1.24</b>	0.94	1.65
Consumption of fruits and vegetables from locations within the sampling frame	No	11	<b>3.24</b>	2.08	5.04	<b>2.84</b>	2.15	3.75	<b>1.13</b>	0.93	1.37
	Yes	17	<b>5.28</b>	3.11	8.97	<b>3.47</b>	2.33	5.17	<b>1.33</b>	1.01	1.74
Frequency of local milk consumption (i.e., milk from animals within the sampling frame)	Rarely or more frequently	7	NA	NA	NA	NA	NA	NA	NA	NA	NA
	Never	21	<b>3.99</b>	2.45	6.48	<b>3.36</b>	2.49	4.54	<b>1.31</b>	1.07	1.60
Drank formula reconstituted with tap water	No	18	<b>5.90</b>	4.21	8.27	<b>3.95</b>	2.98	5.25	<b>1.49</b>	1.26	1.78
	Yes	10	<b>2.52</b>	1.26	5.03	<b>2.20</b>	1.81	2.67	<b>0.90</b>	0.82	0.97
Duration of drinking formula reconstituted with tap water duration (months)	<7	23	<b>5.49</b>	4.10	7.34	<b>3.45</b>	2.56	4.64	<b>1.34</b>	1.10	1.63
	7 to <13	4	NA	NA	NA	NA	NA	NA	NA	NA	NA
	19+	1	NA	NA	NA	NA	NA	NA	NA	NA	NA
Currently breastfeeding or previously breastfed	No	5	NA	NA	NA	NA	NA	NA	NA	NA	NA
	Yes	23	<b>4.41</b>	2.77	7.02	<b>3.55</b>	2.70	4.65	<b>1.33</b>	1.12	1.59

Variable	Category	Frequency <sup>§</sup>	PFHxS			PFOS			PFOA		
			GM	LCI	UCI	GM	LCI	UCI	GM	LCI	UCI
Breastfeeding duration (months)	<7	15	<b>3.24</b>	1.77	5.92	<b>2.59</b>	1.88	3.56	<b>1.06</b>	0.86	1.32
	7 to <19	10	<b>7.24</b>	5.79	9.04	<b>4.37</b>	3.31	5.77	<b>1.51</b>	1.16	1.96
	19+	3	NA	NA	NA	NA	NA	NA	NA	NA	NA
Public Water System	Security WD	21	<b>3.56</b>	2.25	5.63	<b>2.74</b>	2.14	3.51	<b>1.12</b>	0.94	1.33
	Widefield WSD	7	NA	NA	NA	NA	NA	NA	NA	NA	NA

\* Several variables that were collected in the questionnaire are not included in these tables. These variables may not be included because they did not have sufficient variability or were not associated with PFAS blood concentrations in preliminary analyses. These variables include full-time vs. part-time residence and school attendance.

† Geometric means and confidence levels are not shown for categories with fewer than 10 responses.

‡ Detection limits for all PFAS are 0.1 micrograms per liter (µg/L).

§ Some frequency counts may not sum to the total because of missing values. Some variable categories that were presented in the questionnaire were collapsed into larger variable categories.

Table C2 continued. Child blood PFAS geometric means (GM), lower confidence intervals (LCI), and upper confidence intervals (UCI) in micrograms per liter<sup>\*,†,‡</sup>

Variable	Category	Frequency <sup>§</sup>	PFNA			PFDA			MeFOSAA		
			GM	LCI	UCI	GM	LCI	UCI	GM	LCI	UCI
<b>All Children</b>		28	<b>4.35</b>	3.09	6.13	<b>3.21</b>	2.53	4.06	<b>1.24</b>	1.07	1.45
Age (years)	3 to <12	9	NA	NA	NA	NA	NA	NA	NA	NA	NA
	12 to <18	19	<b>0.19</b>	0.15	0.25	<b>0.13</b>	0.09	0.19	<b>0.10</b>	0.07	0.14
Sex	Female	18	<b>0.18</b>	0.14	0.23	<b>0.13</b>	0.09	0.17	<b>0.09</b>	0.07	0.13
	Male	10	<b>0.23</b>	0.17	0.30	<b>0.11</b>	0.08	0.15	<b>0.10</b>	0.07	0.14
Body mass index (kilograms per square meter)	<15	1	NA	NA	NA	NA	NA	NA	NA	NA	NA
	15 to 20	9	NA	NA	NA	NA	NA	NA	NA	NA	NA
	20 to <25	14	<b>0.19</b>	0.14	0.25	<b>0.11</b>	0.09	0.13	<b>0.12</b>	0.08	0.17
	25+	3	NA	NA	NA	NA	NA	NA	NA	NA	NA
Birth order	First born	9	NA	NA	NA	NA	NA	NA	NA	NA	NA
	Second born	12	<b>0.17</b>	0.12	0.23	<b>0.12</b>	0.08	0.16	<b>0.11</b>	0.07	0.17
	Third born	6	NA	NA	NA	NA	NA	NA	NA	NA	NA
	Fifth born	1	NA	NA	NA	NA	NA	NA	NA	NA	NA
Race and ethnicity combined	White, Not Hispanic	15	<b>0.20</b>	0.15	0.27	<b>0.11</b>	0.08	0.14	<b>0.10</b>	0.07	0.16
	Not White, or Hispanic	12	<b>0.19</b>	0.16	0.22	<b>0.14</b>	0.08	0.24	<b>0.08</b>	0.07	0.10
Water consumption at current home (average cups per day)	0 to <2	8	NA	NA	NA	NA	NA	NA	NA	NA	NA
	2 to <4	5	NA	NA	NA	NA	NA	NA	NA	NA	NA
	4+	15	<b>0.19</b>	0.15	0.25	<b>0.13</b>	0.09	0.20	<b>0.10</b>	0.07	0.15
Water consumption at school (average cups per day)	0 to <1	3	NA	NA	NA	NA	NA	NA	NA	NA	NA
	1 to <2	5	NA	NA	NA	NA	NA	NA	NA	NA	NA
	2 to <3	8	NA	NA	NA	NA	NA	NA	NA	NA	NA
	3+	12	<b>0.18</b>	0.14	0.23	<b>0.12</b>	0.08	0.16	<b>0.09</b>	0.07	0.11

Variable	Category	Frequency <sup>s</sup>	PFNA			PFDA			MeFOSAA		
			GM	LCI	UCI	GM	LCI	UCI	GM	LCI	UCI
Length of residency in sampling frame (years)	<6	3	NA	NA	NA	NA	NA	NA	NA	NA	NA
	6 to <12	12	<b>0.21</b>	0.16	0.27	<b>0.11</b>	0.08	0.14	<b>0.11</b>	0.06	0.18
	12 to <18	13	<b>0.18</b>	0.14	0.23	<b>0.13</b>	0.09	0.21	<b>0.09</b>	0.08	0.10
Frequency of direct contact with soil at locations within the sampling frame	A few times per year or less	4	NA	NA	NA	NA	NA	NA	NA	NA	NA
	A few times per month	11	<b>0.17</b>	0.13	0.23	<b>0.12</b>	0.09	0.17	<b>0.10</b>	0.06	0.16
	Three times per week or more	13	<b>0.21</b>	0.19	0.25	<b>0.12</b>	0.09	0.17	<b>0.09</b>	0.07	0.11
Consumption of fruits and vegetables from locations within the sampling frame	No	11	<b>0.15</b>	0.12	0.18	<b>0.09</b>	0.08	0.10	<b>0.10</b>	0.06	0.16
	Yes	17	<b>0.24</b>	0.20	0.28	<b>0.15</b>	0.10	0.22	<b>0.09</b>	0.07	0.12
Frequency of local milk consumption (i.e., milk from animals within the sampling frame)	Rarely or more frequently	7	NA	NA	NA	NA	NA	NA	NA	NA	NA
	Never	21	<b>0.19</b>	0.15	0.24	<b>0.11</b>	0.09	0.14	<b>0.10</b>	0.07	0.14
Drank formula reconstituted with tap water	No	18	<b>0.20</b>	0.16	0.26	<b>0.13</b>	0.09	0.19	<b>0.08</b>	0.07	0.09
	Yes	10	<b>0.18</b>	0.12	0.26	<b>0.10</b>	0.09	0.12	<b>0.12</b>	0.06	0.22
Duration of drinking formula reconstituted with tap water duration (months)	<7	23	<b>0.20</b>	0.17	0.23	<b>0.13</b>	0.09	0.18	<b>0.08</b>	0.07	0.09
	7 to <13	4	NA	NA	NA	NA	NA	NA	NA	NA	NA
	19+	1	NA	NA	NA	NA	NA	NA	NA	NA	NA
Currently breastfeeding or previously breastfed	No	5	NA	NA	NA	NA	NA	NA	NA	NA	NA
	Yes	23	<b>0.20</b>	0.16	0.25	<b>0.13</b>	0.09	0.18	<b>0.10</b>	0.07	0.13

Variable	Category	Frequency <sup>§</sup>	PFNA			PFDA			MeFOSAA		
			GM	LCI	UCI	GM	LCI	UCI	GM	LCI	UCI
Breastfeeding duration (months)	<7	15	<b>0.19</b>	0.14	0.25	<b>0.10</b>	0.08	0.13	<b>0.10</b>	0.06	0.16
	7 to <19	10	<b>0.20</b>	0.15	0.27	<b>0.16</b>	0.10	0.26	<b>0.09</b>	0.08	0.10
	19+	3	NA	NA	NA	NA	NA	NA	NA	NA	NA
Public Water System	Security WD	21	<b>0.19</b>	0.16	0.23	<b>0.11</b>	0.08	0.15	<b>0.10</b>	0.07	0.14
	Widefield WSD	7	NA	NA	NA	NA	NA	NA	NA	NA	NA

\* Several variables that were collected in the questionnaire are not included in these tables. These variables may not be included because they did not have sufficient variability or were not associated with PFAS blood concentrations in preliminary analyses. These variables include full-time vs. part-time residence and school attendance.

† Geometric means and confidence levels are not shown for categories with fewer than 10 responses.

‡ Detection limits for all PFAS are 0.1 micrograms per liter (µg/L).

§ Some frequency counts may not sum to the total because of missing values. Some variable categories that were presented in the questionnaire were collapsed into larger variable categories.

**Table C3. Adult univariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME)\***

Variable	Category	PFHxS			PFOS			PFOA		
		Coef.	p-val	ME (%)	Coef.	p-val	ME (%)	Coef.	p-val	ME (%)
Age	NA—continuous variable	0.011	<0.001	2.5	0.008	<0.001	1.9	0.006	<0.001	1.3
Sex	Male	0.076	0.1007	19.2	0.154	0.0002	42.6	0.046	0.1953	11.1
	Female	—	—	—	—	—	—	—	—	—
Body mass index	NA—continuous variable	-0.002	0.5597	-0.6	-0.004	0.3407	-0.8	0.000	0.9657	0.0
Race and ethnicity Combined	Not White, or Hispanic	-0.091	0.1498	-19.0	-0.043	0.4381	-9.5	-0.066	0.1467	-14.1
	White alone, not Hispanic	—	—	—	—	—	—	—	—	—
Length of residence at current address (years)	NA—continuous variable	0.016	<0.001	3.7	0.009	<0.001	2.1	0.007	<0.001	1.6
Total length of residence in sampling frame over the past 20 years (years)	NA—continuous variable	0.041	<0.001	9.8	0.017	<0.001	4.1	0.014	0.0001	3.3
Current and primary source of drinking water	Bottled water	-0.158	0.0126	-30.6	-0.080	0.1010	-16.8	-0.070	0.0709	-15.0
	Public water system	—	—	—	—	—	—	—	—	—
Tap water consumption at current home (average cups per day)	NA—continuous variable	0.016	0.0037	3.7	0.009	0.0160	2.0	0.008	0.0435	1.7

Variable	Category	PFHxS			PFOS			PFOA		
		Coef.	p-val	ME (%)	Coef.	p-val	ME (%)	Coef.	p-val	ME (%)
Current use of filter or treatment device for tap water at home	None, no filter or treatment device	—	—	—	—	—	—	—	—	—
	None, drink bottled water only	-0.319	0.0011	-52.0	-0.245	0.0005	-43.1	-0.167	0.0050	-31.9
	Use at least one filter or treatment device	-0.023	0.7191	-5.2	-0.055	0.2661	-12.0	0.030	0.5008	7.1
History of kidney disease	Yes	-0.065	0.5929	-13.8	0.067	0.5572	16.8	-0.006	0.9406	-1.3
	No	—	—	—	—	—	—	—	—	—
Frequency of blood donation	Once or more a Year	0.040	0.7450	9.7	0.141	0.0825	38.4	0.031	0.7042	7.4
	Never/rarely	—	—	—	—	—	—	—	—	—
Frequency of house cleaning	Three times per week or more	-0.100	0.0897	-20.5	0.047	0.3414	11.3	0.011	0.7626	2.6
	A few times per month or less	—	—	—	—	—	—	—	—	—
Frequency of stain-resistant product use	Rarely or more frequently	-0.097	0.3220	-20.1	0.027	0.6859	6.5	-0.064	0.4037	-13.7
	Never	—	—	—	—	—	—	—	—	—
Frequency of direct contact with soil at locations within the sampling frame	A few times per month	0.104	0.1444	27.1	0.027	0.6432	6.4	0.079	0.0819	20.0
	Three times per week or more	0.004	0.9571	0.9	0.006	0.9230	1.4	0.004	0.9411	0.8
	A few times per year or less	—	—	—	—	—	—	—	—	—

Variable	Category	PFHxS			PFOS			PFOA		
		Coef.	p-val	ME (%)	Coef.	p-val	ME (%)	Coef.	p-val	ME (%)
Consumption of fruits and vegetables from locations within the sampling frame	Yes	0.049	0.3890	11.9	0.094	0.0568	24.1	0.049	0.2031	11.9
	No	—	—	—	—	—	—	—	—	—
Consumption of local fish (i.e., fish caught within the sampling frame)	Yes	-0.043	0.7542	-9.5	0.078	0.5046	19.7	0.005	0.9450	1.3
	No	—	—	—	—	—	—	—	—	—
Frequency of fast food consumption	A few times per month	-0.097	0.3083	-19.9	-0.052	0.4911	-11.2	-0.041	0.4904	-9.1
	Three times per week or more	-0.168	0.1116	-32.1	-0.166	0.0525	-31.8	-0.111	0.1162	-22.5
	A few times per year or less	—	—	—	—	—	—	—	—	—
Presence of carpeting in bedroom, living room, or kitchen	Yes	-0.073	0.2044	-15.5	-0.068	0.2095	-14.5	-0.039	0.2841	-8.6
	No	—	—	—	—	—	—	—	—	—
Occupational exposures (count of jobs with potential PFAS exposures)	One or more occupational exposures	-0.173	0.0122	-32.9	-0.027	0.6019	-6.1	0.012	0.8087	2.9
	None	—	—	—	—	—	—	—	—	—
Biological children	Yes	0.161	0.0447	44.8	0.070	0.2229	17.5	0.079	0.2249	19.9
	No	—	—	—	—	—	—	—	—	—
Number of biological children	NA—continuous variable	0.069	0.0027	17.2	0.044	0.0194	10.6	0.029	0.1136	6.8
Breastfeeding or previously breastfed children	Yes	0.004	0.9534	1.0	0.008	0.8950	1.8	0.009	0.8703	2.1
	No	—	—	—	—	—	—	—	—	—

Variable	Category	PFHxS			PFOS			PFOA		
		Coef.	p-val	ME (%)	Coef.	p-val	ME (%)	Coef.	p-val	ME (%)
Total duration of breastfeeding for all biological children (months)	NA—continuous variable	-0.002	0.455	-0.5	-0.001	0.818	-0.1	-0.001	0.743	-0.2
Public Water Supply	Widefield WSD	-0.008	0.9033	-1.7	0.125	0.0142	33.3	0.035	0.3943	8.4
	Security WD	—	—	—	—	—	—	—	—	—

\* Not all categorical variables included in Table C1 are included in Table C3: variable categories that had fewer than 10 responses were not included in the regressions (Table C3). This variable is frequency of local milk consumption.

Table C3 continued. Adult univariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME)\*

Variable	Category	PFNA			PFDA			MeFOSAA		
		Coef.	p-val	ME (%)	Coef.	p-val	ME (%)	Coef.	p-val	ME (%)
Age	NA—continuous variable	0.005	<0.001	1.2	0.001	0.3799	0.2	0.006	<0.001	1.4
Sex	Male	0.061	0.0584	15.0	-0.022	0.3993	-5.0	0.013	0.6812	3.0
	Female	—	—	—	—	—	—	—	—	—
Body mass index	NA—continuous variable	-0.005	0.0847	-1.1	-0.005	0.0098	-1.2	-0.005	0.1565	-1.1
Race and ethnicity Combined	Not White, or Hispanic	0.043	0.3435	10.4	0.081	0.0488	20.5	-0.015	0.7531	-3.3
	White alone, not Hispanic	—	—	—	—	—	—	—	—	—
Length of residence at current address (years)	NA—continuous variable	0.001	0.3137	0.3	-0.002	0.0766	-0.5	0.008	<0.001	1.8
Total length of residence in sampling frame over the past 20 years (years)	NA—continuous variable	0.000	0.9770	0.0	-0.009	0.0135	-2.0	0.015	<0.001	3.6
Current and primary source of drinking water	Bottled water	0.004	0.9304	0.9	-0.021	0.5562	-4.7	0.035	0.4940	8.3
	Public water system	—	—	—	—	—	—	—	—	—
Tap water consumption at current home (average cups per day)	NA—continuous variable	0.002	0.5430	0.4	0.003	0.2028	0.7	0.004	0.3014	0.9

Variable	Category	PFNA			PFDA			MeFOSAA		
		Coef.	p-val	ME (%)	Coef.	p-val	ME (%)	Coef.	p-val	ME (%)
Current use of filter or treatment device for tap water at home	None, no filter or treatment device	—	—	—	—	—	—	—	—	—
	None, drink bottled water only	-0.063	0.3064	-13.5	-0.020	0.6931	-4.6	-0.089	0.2121	-18.6
	Use at least one filter or treatment device	-0.003	0.9424	-0.7	0.015	0.7069	3.5	-0.041	0.4343	-9.0
History of kidney disease	Yes	0.090	0.1907	23.1	0.054	0.3768	13.2	0.093	0.2657	24.0
	No	—	—	—	—	—	—	—	—	—
Frequency of blood donation	Once or more a Year	0.043	0.5789	10.4	-0.001	0.9882	-0.3	0.110	0.4760	28.7
	Never/rarely	—	—	—	—	—	—	—	—	—
Frequency of house cleaning	Three times per week or more	0.062	0.0933	15.4	0.082	0.0168	20.8	-0.008	0.8561	-1.8
	A few times per month or less	—	—	—	—	—	—	—	—	—
Frequency of stain-resistant product use	Rarely or more frequently	0.110	0.0988	28.8	0.092	0.1035	23.7	0.183	0.0267	52.3
	Never	—	—	—	—	—	—	—	—	—
Frequency of direct contact with soil at locations within the sampling frame	A few times per month	0.069	0.1278	17.3	0.040	0.2867	9.6	0.035	0.4759	8.4
	Three times per week or more	0.000	0.9928	0.1	0.043	0.2623	10.5	-0.011	0.8173	-2.4
	A few times per year or less	—	—	—	—	—	—	—	—	—

Variable	Category	PFNA			PFDA			MeFOSAA		
		Coef.	p-val	ME (%)	Coef.	p-val	ME (%)	Coef.	p-val	ME (%)
Consumption of fruits and vegetables from locations within the sampling frame	Yes	0.064	0.0946	15.9	0.096	0.0058	24.6	0.019	0.6410	4.5
	No	—	—	—	—	—	—	—	—	—
Consumption of local fish (i.e., fish caught within the sampling frame)	Yes	0.097	0.1959	25.1	0.069	0.5253	17.3	-0.099	0.2994	-20.4
	No	—	—	—	—	—	—	—	—	—
Frequency of fast food consumption	A few times per month	-0.018	0.7427	-4.1	-0.029	0.5540	-6.5	0.038	0.5567	9.2
	Three times per week or more	-0.088	0.1663	-18.3	-0.083	0.1355	-17.4	-0.075	0.2828	-15.9
	A few times per year or less	—	—	—	—	—	—	—	—	—
Presence of carpeting in bedroom, living room, or kitchen	Yes	0.015	0.7209	3.5	0.007	0.8664	1.6	0.077	0.0654	19.5
	No	—	—	—	—	—	—	—	—	—
Occupational exposures (count of jobs with potential PFAS exposures)	One or more occupational exposures	0.094	0.0351	24.2	0.000	0.9937	-0.1	-0.042	0.4966	-9.2
	None	—	—	—	—	—	—	—	—	—
Biological children	Yes	0.000	0.9971	0.1	-0.033	0.6142	-7.3	0.095	0.0846	24.4
	No	—	—	—	—	—	—	—	—	—
Number of biological children	NA—continuous variable	-0.003	0.8716	-0.7	-0.012	0.4252	-2.7	0.024	0.1840	5.6
Breastfeeding or previously breastfed children	Yes	-0.022	0.6821	-4.9	0.003	0.9503	0.6	-0.032	0.5187	-7.0
	No	—	—	—	—	—	—	—	—	—

Variable	Category	PFNA			PFDA			MeFOSAA		
		Coef.	p-val	ME (%)	Coef.	p-val	ME (%)	Coef.	p-val	ME (%)
Total duration of breastfeeding for all biological children (months)	NA—continuous variable	-0.002	0.316	-0.4	-0.000	0.963	-0.0	-0.002	0.259	-0.4
Public Water Supply	Widefield WSD	0.027	0.5229	6.4	-0.014	0.7158	-3.2	0.042	0.3797	10.1
	Security WD	—	—	—	—	—	—	—	—	—

\* Not all categorical variables included in Table C1 are included in Table C3: variable categories that had fewer than 10 responses were not included in the regressions (Table C3). This variable is frequency of local milk consumption.

**Table C4. Child univariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME)**

ParamModel	Parameter	PFHxS			PFOS			PFOA		
		Coef.	p-val	ME (%)	Coef.	p-val	ME (%)	Coef.	p-val	ME (%)
Age	NA—continuous variable	0.001	0.963	0.2	0.005	0.704	1.1	-0.008	0.419	-1.9
Sex	Male	-0.050	0.7753	-10.9	0.066	0.4871	16.3	-0.004	0.9554	-1.0
	Female	—	—	—	—	—	—	—	—	—
Body mass index (kilograms per square meter)	NA—continuous variable	0.009	0.3863	2.0	-0.001	0.7951	-0.3	-0.005	0.4697	-1.3
Race and ethnicity combined	Not White alone, or Hispanic	0.076	0.6380	19.1	0.037	0.7190	8.9	0.035	0.6054	8.3
	White alone, not Hispanic	—	—	—	—	—	—	—	—	—
Water consumption at current home (average cups per day)	NA—continuous variable	0.007	0.5195	1.6	0.006	0.6764	1.4	0.001	0.9490	0.1
Water consumption at school (average cups per day)	NA—continuous variable	0.028	0.1519	6.6	0.001	0.9611	0.2	-0.003	0.8406	-0.7
Length of residency in sampling frame (years)	NA—continuous variable	0.030	0.1092	7.3	0.008	0.4670	1.8	-0.002	0.7934	-0.5
Drank formula reconstituted with tap water	Yes	-0.370	0.0383	-57.3	-0.255	0.0037	-44.4	-0.223	0.0002	-40.1
	No	—	—	—	—	—	—	—	—	—
Drank formula reconstituted with tap water duration (months)	NA—continuous variable	-0.032	0.052	-7.2	-0.018	0.052	-4.0	-0.014	0.047	-3.3

ParamModel	Parameter	PFHxS			PFOS			PFOA		
		Coef.	p-val	ME (%)	Coef.	p-val	ME (%)	Coef.	p-val	ME (%)
Breastfeeding duration (months)	NA—continuous variable	0.009	0.251	2.0	0.010	0.041	2.4	0.009	0.018	2.2
Public Water Supply	Widefield WSD	0.349	0.0047	123.6	0.273	0.0073	87.5	0.180	0.0047	51.3
	Security WD	—	—	—	—	—	—	—	—	—

\* Not all categorical variables included in Table C2 are also included in Table C4: variable categories that had fewer than 10 responses were not included in the regressions (Table C4). These variables include birth order, frequency of direct contact with soil at locations within the sampling frame, frequency of local fruit and vegetable consumption, frequency of local milk consumption, and currently breastfeeding or previously breastfed.

Table C4 continued. Child univariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME)

ParamModel	Parameter	PFNA			PFDA			MeFOSAA		
		Coef.	p-val	ME (%)	Coef.	p-val	ME (%)	Coef.	p-val	ME (%)
Age	NA—continuous variable	-0.021	0.683	-4.8	0.120	0.117	31.9	0.108	0.132	28.3
Sex	Male	0.096	0.2310	24.7	-0.053	0.2253	-11.4	0.018	0.8376	4.3
	Female	—	—	—	—	—	—	—	—	—
Body mass index (kilograms per square meter)	NA—continuous variable	-0.013	0.0708	-3.0	-0.014	0.1728	-3.1	0.005	0.5913	1.1
Race and ethnicity combined	Not White alone, or Hispanic	-0.013	0.8476	-3.0	0.123	0.3129	32.6	-0.095	0.3223	-19.7
	White alone, not Hispanic	—	—	—	—	—	—	—	—	—
Water consumption at current home (average cups per day)	NA—continuous variable	-0.009	0.0771	-2.1	0.010	0.0274	2.3	0.002	0.7708	0.5
Water consumption at school (average cups per day)	NA—continuous variable	-0.003	0.8722	-0.6	-0.003	0.8111	-0.7	-0.013	0.5486	-2.9
Length of residency in sampling frame (years)	NA—continuous variable	-0.017	0.0674	-3.8	0.011	0.3720	2.7	-0.015	0.1847	-3.4
Drank formula reconstituted with tap water	Yes	-0.053	0.5864	-11.4	-0.103	0.0974	-21.2	0.159	0.2269	44.2
	No	—	—	—	—	—	—	—	—	—
Drank formula reconstituted with tap water duration (months)	NA—continuous variable	-0.001	0.881	-0.2	-0.009	0.132	-2.1	0.014	0.145	3.3
Breastfeeding duration (months)	NA—continuous variable	0.001	0.850	0.2	0.002	0.615	0.5	-0.001	0.743	-0.3

\* Not all categorical variables in included in Table C2 are also included in Table C4: variable categories that had fewer than 10 responses were not included in the regressions (Table C4). These variables include birth order, frequency of direct contact with soil at locations within the sampling frame, frequency of local fruit and vegetable consumption, frequency of local milk consumption, and currently breastfeeding or previously breastfed.

**Table C5. PFHxS adult multivariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME)**

Parameter	Coef.	p-val	ME (%)
Age (continuous)	0.011	<.0001	2.5
Sex: male* (categorical)	0.381	0.0111	140.4
Age x sex: male*,† (continuous)	-0.005	0.0379	-1.2
Years in sampling frame in the past 20 years (continuous)	0.030	<.0001	7.1
Kidney disease history‡: yes (categorical)	-0.216	0.0464	-39.2
Occupational exposure history§: yes (categorical)	-0.140	0.0235	-27.6
Filter use**: none, use bottled water only	-0.195	0.0385	-36.2
Filter use**: one or more filter or treatment device	0.002	0.9665	0.5

Model statistics: R<sup>2</sup> = 0.3005 p-value = <0.0001, n = 312, n-households = 186, intercept = -0.020

\* Reference category is adult participants who identified as female.

† This variable is an interaction term between age and sex.

‡ Reference category is adult participants who did not report a history of kidney disease.

§ Reference category is adult participants who did not report a history of occupational exposure.

\*\* Reference category is adult participants who reported using no filter or treatment device.

**Table C6. PFHxS adult female multivariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME)**

Parameter	Coef.	p-val	ME (%)
Age (continuous)	0.010	<.0001	2.4
Years in sampling frame in the past 20 years (continuous)	0.031	<.0001	7.3
Filter use*: none, use bottled water only	-0.191	0.0282	-35.6
Filter use*: one or more filter or treatment device	-0.056	0.444	-12.1

Model statistics: R<sup>2</sup> = 0.3035 p-value = <0.0001, n = 175, n-households = 151, intercept = 0.003

\* Reference category is adult female participants who reported using no filter or treatment device.

**Table C7. PFHxS adult male multivariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME)**

Parameter	Coef.	p-val	ME (%)
Age (continuous)	0.006	0.0050	1.4
Years in sampling frame in the past 20 years (continuous)	0.033	<.0001	7.8
Kidney disease history*: yes (categorical)	-0.356	0.0314	-55.9
Occupational exposure history†: yes (categorical)	-0.143	0.0468	-28.1

Model statistics: R<sup>2</sup> = 0.2681 p-value = <0.0001, n = 138, n-households = 123, intercept = 0.283

\* Reference category is adult male participants who did not report a history of kidney disease.

† Reference category is adult male participants who did not report a history of occupational exposure.

**Table C8. PFOS adult multivariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME)**

Parameter	Coef.	p-val	ME (%)
Age (continuous)	0.007	<.0001	1.7
Sex: male* (categorical)	0.152	0.0001	42.0
Years in sampling frame in the past 20 years (continuous)	0.009	0.0194	2.2
Filter use†: none, use bottled water only	-0.176	0.0105	-33.4
Filter use†: one or more filter or treatment device	-0.041	0.3580	-9.0
Frequency of house cleaning‡: three times per week or more (categorical)	0.094	0.0437	24.1
Consumption of fruits and vegetables from locations within the sampling frame§: yes (categorical)	0.092	0.0431	23.5

Model statistics: R<sup>2</sup> = 0.2149, p-value = <0.0001, n = 315, n-households = 186, intercept = 0.143

\* Reference category is adult participants who identified as female.

† Reference category is adult participants who reported using no filter or treatment device.

‡ Reference category is adult participants who reported cleaning their house a few times per month or less.

§ Reference category is adult participants who reported not consuming fruits or vegetables from locations within the sampling frame.

**Table C9. PFOS adult female multivariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME)**

Parameter	Coef.	p-val	ME (%)
Age (continuous)	0.010	<.0001	2.3
Filter use*: none, use bottled water only	-0.180	0.0060	-34.0
Filter use*: one or more filter or treatment device	-0.033	0.6053	-7.3
Frequency of house cleaning†: three times per week or more (categorical)	0.149	0.0120	40.9

Model statistics: R2 = 0.1838, p-value = <0.0001, n = 175, n-households = 151, intercept = 0.169

\* Reference category is adult female participants who reported using no filter or treatment device.

† Reference category is adult female participants who reported cleaning their house a few times per month or less.

**Table C10. PFOS adult male multivariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME)**

Parameter	Coef.	p-val	ME (%)
Age (continuous)	0.006	0.0006	1.4
Years in sampling frame in the past 20 years (continuous)	0.013	0.0449	3.0
Consumption of fruits and vegetables from locations within the sampling frame*: yes (categorical)	0.138	0.0330	37.3

Model statistics: R2 = 0.1630, p-value = <0.0001, n = 142, n-households = 126, intercept = 0.296

\* Reference category is adult male participants who reported not consuming fruits of vegetables from locations within the sampling frame.

**Table C11. PFOA adult multivariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME)**

Parameter	Coef.	p-val	ME (%)
Age (continuous)	0.004	<.0001	1.0
Years in sampling frame in the past 20 years (continuous)	0.009	0.0117	2.0
Filter use*: none, use bottled water only	-0.122	0.0399	-24.4
Filter use*: one or more filter or treatment device	0.049	0.2435	12.0

Model statistics: R2 = 0.1275, p-value = <0.0001, n = 318, n-households = 188, intercept = -0.060

\* Reference category is adult participants who reported using no filter or treatment device.

**Table C12. PFOA adult male multivariate regression results including coefficient estimate (Coef.), p-value (p-val), and marginal effect (ME)**

Parameter	Coef.	p-val	ME (%)
Age (continuous)	0.003	0.0203	0.6
Years in sampling frame in the past 20 years (continuous)	0.012	0.0063	2.7

Model statistics: R2 = 0.0768, p-value = <0.0001, n = 143, n-households = 127, intercept = 0.036

## Box and Whisker Plots (or Boxplots)

Figure C1. Boxplot of adult blood (serum) PFAS concentrations by age




Figure C2. Boxplot of adult blood (serum) PFAS concentrations by sex




Figure C3. Boxplot of adult blood (serum) PFAS concentrations by race and ethnicity




Figure C4. Boxplot of adult blood (serum) PFAS concentrations by body mass index




Figure C5. Boxplot of adult blood (serum) PFAS concentrations by years in current home




Figure C6. Boxplot of adult blood (serum) PFAS concentrations by years in sampling frame (past 20 years)




Figure C7. Boxplot of adult blood (serum) PFAS concentrations by cups of tap water drank at home




Figure C8. Boxplot of adult blood (serum) PFAS concentrations by water filter type




Figure C9. Boxplot of adult blood (serum) PFAS concentrations by kidney disease history




Figure C10. Boxplot of adult blood (serum) PFAS concentrations by blood donation frequency




Figure C11. Boxplot of adult blood (serum) PFAS concentrations by home cleaning frequency




Figure C12. Boxplot of adult blood (serum) PFAS concentrations by stain-resistant product use




Figure C13. Boxplot of adult blood (serum) PFAS concentrations by frequency of contact with soil




Figure C14. Boxplot of adult blood (serum) PFAS concentrations by local fruit and vegetable consumption




Figure C15. Boxplot of adult blood (serum) PFAS concentrations by local fish consumption




Figure C16. Boxplot of adult blood (serum) PFAS concentrations by local milk consumption




Figure C17. Boxplot of adult blood (serum) PFAS concentrations by fast food consumption frequency




Figure C18. Boxplot of adult blood (serum) PFAS concentrations by presence of carpet in home




Figure C19. Boxplot of adult blood (serum) PFAS concentrations by occupational exposure




Figure C20. Boxplot of adult female blood (serum) PFAS concentrations by breastfeeding history




Figure C21. Boxplot of adult female blood (serum) PFAS concentrations by breastfeeding duration




Figure C22. Boxplot of adult female blood (serum) PFAS concentrations by biological children variable




Figure C23. Boxplot of adult female blood (serum) PFAS concentrations by number of biological children




Figure C24. Boxplot of child blood (serum) PFAS concentrations by age




Figure C25. Boxplot of child blood (serum) PFAS concentrations by sex




Figure C26. Boxplot of child blood (serum) PFAS concentrations by body mass index




Figure C27. Boxplot of child blood (serum) PFAS concentrations by race and ethnicity combined




Figure C28. Boxplot of child blood (serum) PFAS concentrations by water consumption at current home




Figure C29. Boxplot of child blood (serum) PFAS concentrations by water consumption at school




Figure C30. Boxplot of child blood (serum) PFAS concentrations by length of residency in sampling frame




Figure C31. Boxplot of child blood (serum) PFAS concentrations by frequency of contact with soil




Figure C32. Boxplot of child blood (serum) PFAS concentrations by local fruit and vegetable consumption




Figure C33. Boxplot of child blood (serum) PFAS concentrations by drinking formula reconstituted with tap water




Figure C34. Boxplot of child blood (serum) PFAS concentrations by duration of drinking formula reconstituted with tap water




Figure C35. Boxplot of child blood (serum) PFAS concentrations by history of breastfeeding




Figure C36. Boxplot of child blood (serum) PFAS concentrations by breastfeeding duration

