

9. REFERENCES

- Abedin Z, Cook RC, Jr, Milberg RM. 1980. Cardiac toxicity of perchloroethylene (a dry cleaning agent). *South Med J* 73:1081-1083.
- Abrahamsson K, Ekdahl A, Cohen J, et al. 1995. Marine algae-a source of trichloroethylene and perchloroethylene. *Limnol Oceanogr* 40(7):1321-1326.
- Abreu LDV, Johnson PC. 2005. Effect of vapor source-building separation and building construction on soil vapor intrusion as studied with a three-dimensional numerical model. *Environ Sci Technol* 39(12):4550-4561.
- ACGIH. 1991. Documentation of the threshold limit values and biological exposure indices. Sixth edition. Cincinnati, OH: American Conference of Governmental Industrial Hygienists.
- ACGIH. 1995. Threshold limit values (TLVs) for chemical substances and physical agents and biological exposure indices (BEIs) for 1995-1996. Perchloroethylene. Cincinnati, OH: American Conference of Governmental Industrial Hygienists.
- ACGIH. 2012. Tetrachloroethylene. Threshold limit values for chemical substances and physical agents and biological exposure indices. Cincinnati, OH: American Conference of Governmental Industrial Hygienists, 55, 110.
- Adgate JL, Church TR, Ryan AD, et al. 2004. Outdoor, indoor, and personal exposure to VOCs in children. *Environ Health Perspect* 112(14):1386-1392.
- Adlercreutz H. 1995. Phytoestrogens: Epidemiology and a possible role in cancer protection. *Environ Health Perspect* 103(Suppl 7):103-112.
- Aggazzotti G, Fantuzzi G, Predieri G, et al. 1994a. Indoor exposure to perchloroethylene (PCE) in individuals living with dry cleaning workers. *Sci Total Environ* 156:133-137.
- Aggazzotti G, Fantuzzi G, Righi E, et al. 1994b. Occupational and environmental exposure to perchloroethylene (PCE) in dry cleaners and their family members. *Arch Environ Health* 49(6):487-493.
- Ahlborg G Jr. 1990. Pregnancy outcome among women working in laundries and dry cleaning shops using tetrachloroethylene. *Am J Ind Med* 17:567-575.
- AIHA. 2011. Emergency response planning guidelines (ERPG). Fairfax, VA: American Industrial Hygiene Association.
<http://www.aiha.org/INSIDEAIHA/GUIDELINEDEVELOPMENT/ERPG/Pages/default.aspx>. April 24, 2013.
- Albee RR, Mattsson JL, Bradley GJ, et al. 1991. Acute neurophysiologic effects of 1,1,2,2-tetrachloroethylene in rats. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8E. OTS0540058. EPA88920003407.

*Not cited in text

9. REFERENCES

- Altman PL, Dittmer DS. 1974. Biological handbooks: Biology data book. Vol. III. 2nd ed. Bethesda, MD: Federation of American Societies of Experimental Biology.
- Altmann L, Bottger A, Wiegand H. 1990. Neurophysiological and psychophysical measurements reveal effects of acute low-level organic solvent exposure in humans. *Int Arch Occup Environ Health* 62:493-499.
- Altmann L, Florian Neuhann H, Kramer U, et al. 1995. Neurobehavioral and neurophysiological outcome of chronic low-level tetrachloroethene exposure measured in neighborhoods of dry cleaning shops. *Environ Res* 69:83-89.
- Altmann L, Wiegand H, Bottger A, et al. 1992. Neurobehavioral and neurophysiological outcomes of acute repeated perchloroethylene exposure. *Appl Psychol* 41(3):269-279.
- Anders MW, Lash L, Dekant W, et al. 1988. Biosynthesis and biotransformation of glutathione S-conjugates to toxic metabolites. *CRC Crit Rev Toxicol* 18:311-341.
- Andersen A, Barlow L, Engeland A, et al. 1999. Work-related cancer in the Nordic countries. *Scand J Work Environ Health* 25(Suppl 2):1-116.
- Andersen ME, Clewell HJ, Gargas ML, et al. 1987. Physiologically based pharmacokinetics and the risk assessment process for methylene chloride. *Toxicol Appl Pharmacol* 87(2):185-205.
- Andersen ME, Krishnan K. 1994. Relating *in vitro* to *in vivo* exposures with physiologically based tissue dosimetry and tissue response models. In: Salem H, ed. *Animal test alternatives: Refinement, reduction, and replacement*. New York, NY: Marcel Dekker, Inc., 9-25.
- Andrasik I, Cloutet D. 1990. Monitoring solvent vapors in dry cleaning plants. *International Fabricare Institute Focus on Drycleaning* 14(3):1-8.
- Andrys C, Hanovcová I, Chýlková V, et al. 1997. Immunological monitoring of dry-cleaning shop workers. Exposure to tetrachloroethylene. *Cent Eur J Public Health* 5(3):136-142.
- Anttila A, Pukkala E, Sallmen M, et al. 1995. Cancer incidence among Finnish workers exposed to halogenated hydrocarbons. *J Occup Environ Med* 37(7):797-806.
- Antoine SR, DeLeon IR, O'Dell-Smith RM. 1986. Environmentally significant volatile organic pollutants in human blood. *Bull Environ Contam Toxicol* 36:364-371.
- Aoki A, Suzuki H, Kawabata Y, et al. 1994. Effect of perchloroethylene inhalation on nasal mucosa in mice. *Eur Arch Otorhinolaryngol* 251:361-365.
- APHA. 1992. Standard methods for the examination of water and wastewater, 18th ed. American Public Health Association, American Water Works Association and Pollution Control Federation, Washington, DC, 6-42-6-47.

9. REFERENCES

- Aranda-Rodriguez R, Cabecinha A, Harvie J, et al. 2015. A method for quantification of volatile organic compounds in blood by SPME-GC-MS/MS with broader application: From non-occupational exposure population to exposure studies. *J Chromatogr B Analyt Technol Biomed Life Sci* 992:76-85. 10.1016/j.jchromb.2015.04.020.
- Aranyi C, O'Shea WJ, Graham JA, et al. 1986. The effects of inhalation of organic chemical air contaminants on murine lung host defenses. *Fundam Appl Toxicol* 6:713-720.
- Aschengrau A, Ozonoff D, Paulu C, et al. 1993. Cancer risk and tetrachloroethylene-contaminated drinking water in Massachusetts. *Arch Environ Health* 48(5):284-292.
- Aschengrau A, Paulu C, Ozonoff D. 1998. Tetrachloroethylene-contaminated drinking water and the risk of breast cancer. *Environ Health Perspect* 106(Suppl 4):947-953.
- Aschengrau A, Rogers S, Ozonoff D. 2003. Perchloroethylene-contaminated drinking water and the risk of breast cancer: Additional results from Cape Cod, Massachusetts, USA. *Environ Health Perspect* 111(2):167-173.
- Aschengrau A, Weinberg JM, Janulewicz PA, et al. 2009. Prenatal exposure to tetrachloroethylene-contaminated drinking water and the risk of congenital anomalies: A retrospective cohort study. *Environ Health* 8:44.
- Aschengrau A, Weinberg JM, Janulewicz PA, et al. 2011. Affinity for risky behaviors following prenatal and early childhood exposure to tetrachloroethylene (PCE)-contaminated drinking water: A retrospective cohort study. *Environ Health* 10:102.
- Aschengrau A, Weinberg JM, Janulewicz PA, et al. 2012. Occurrence of mental illness following prenatal and early childhood exposure to tetrachloroethylene (PCE)-contaminated drinking water: A retrospective cohort study. *Environ Health* 11:2.
- Aschengrau A, Weinberg J, Rogers S, et al. 2008. Prenatal exposure to tetrachloroethylene-contaminated drinking water and the risk of adverse birth outcomes. *Environ Health Perspect* 116(6):814-820.
- Aschengrau A, Winter MR, Vieira VM, et al. 2015. Long-term health effects of early life exposure to tetrachloroethylene (PCE)-contaminated drinking water: A retrospective cohort study. *Environ Health* 14:36. 10.1186/s12940-015-0021-z.
- Ashley DL, Bonin MA, Cardinali FL, et al. 1992. Determining volatile organic compounds in human blood from a large sample population by using purge and trap gas chromatography/mass spectrometry. *Anal Chem* 64:1021-1029.
- Atkinson R. 1985. Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds under atmospheric conditions. *Chem Rev* 85:69-133.
- Atkinson R, Carter WPL. 1984. Kinetics and mechanisms of the gas-phase reactions of ozone with organic compounds under atmospheric conditions. *Chem Rev* 84:437-470.
- ATSDR. 1989. Decision guide for identifying substance-specific data needs related to toxicological profiles; Notice. Agency for Toxic Substances and Disease Registry. *Fed Regist* 54(174):37618-37634.

9. REFERENCES

- ATSDR. 1997. Toxicological profile for trichloroethylene. U.S. Department of Health and Human Services, Agency for Toxic Substances and Disease Registry, TP-92/18.
- ATSDR. 1998. Volatile organic compounds in drinking water and adverse pregnancy outcome. United State Marine Corps Base, Camp Lejeune, North Carolina. U.S. Department of Health and Human Services, Public Health Service.
- ATSDR. 2004. Interaction profile for 1,1,1-trichloroethane, 1,1-dichloroethane, trichloroethylene and tetrachloroethylene. Atlanta, GA: U.S. Department of Health and Human Services, Agency for Toxic Substances and Disease Registry.
- ATSDR. 2006a. Health consultation. Evaluation of tetrachloroethylene vapor intrusion into buildings located above a contaminated aquifer. Atlanta, GA: U.S. Department of Health and Human Services, Agency for Toxic Substances and Disease Registry, Division of Health Assessment and Consultation.
- ATSDR. 2006b. Toxicological profile for vinyl chloride. Atlanta, GA: Agency for Toxic Substances and Disease Registry, U.S. Department of Health and Human Services.
- ATSDR. 2007a. Analyses of groundwater flow, contaminant fate and transport, and distribution of drinking water at Tarawa terrace and vicinity, U.S. Marine Corps Base Camp Lejeune, North Carolina: Historical reconstruction and present-day conditions. Chapter E: Occurrence of contaminants in groundwater. Atlanta, GA: U.S. Department of Health and Human Services, Agency for Toxic Substances and Disease Registry, Division of Health Assessment and Consultation. http://www.atsdr.cdc.gov/sites/lejeune/docs/ChapterE_TarawaTerrace_revised0108.pdf. August 27, 2013.
- ATSDR. 2007b. Interaction profile for carbon monoxide, formaldehyde, methylene chloride, nitrogen dioxide, and tetrachloroethylene. Atlanta, GA: U.S. Department of Health and Human Services, Agency for Toxic Substances and Disease Registry.
- ATSDR. 2013. Volatile organic compounds in drinking water. Camp Lejeune. Atlanta, GA: Centers for Disease Control and Prevention, Agency for Toxic Substances and Disease Registry.
- ATSDR. 2014. Medical Management Guidelines (MMGs). Tetrachloroethylene. Atlanta, GA: Agency for Toxic Substances and Disease Registry. <https://www.atsdr.cdc.gov/mmg/mmg.asp?id=261&tid=48.pdf>. September 3, 2018.
- ATSDR. 2016. Addendum to the toxicological profile for vinyl chloride. Atlanta, GA: Agency for Toxic Substances and Disease Registry, U.S. Department of Health and Human Services.
- ATSDR. 2017a. Tetrachloroethylene. Full SPL data. Substance priority list (SPL) resource page. Agency for Toxic Substances and Disease Registry, Centers for Disease Control and Prevention. <http://www.atsdr.cdc.gov/SPL/resources/index.html>. May 29, 2018.
- ATSDR. 2017b. ATSDR Assessment of the evidence for the drinking water contaminants at Camp Lejeune and specific cancers and other diseases. Agency for Toxic Substances and Disease Registry. https://www.atsdr.cdc.gov/sites/lejeune/docs/atsdr_summary_of_the_evidence_for_causality_tce_pce_508.pdf. October 22, 2018.

9. REFERENCES

- ATSDR. 2018. Agency for Toxic Substances and Disease Registry case studies in environmental medicine. Tetrachloroethylene toxicity. Atlanta, GA: Agency for Toxic Substances and Disease Registry. <https://www.atsdr.cdc.gov/csem/csem.asp?csem=36&po=0.pdf>. September 3, 2018.
- Azimi M, Bahrami MR, Rezaei Hachesu V, et al. 2017. Primary DNA damage in dry cleaners with perchlorethylene exposure. *Int J Occup Environ Med* 8(4):224-231. <http://doi.org/10.15171/ijoem.2017.1089>.
- Bagnell PC, Ellenberger HA. 1977. Obstructive jaundice due to a chlorinated hydrocarbon in breast milk. *Can Med Assoc J* 117:1047-1048.
- Bale AS, Barone S, Scott CS, et al. 2011. A review of potential neurotoxic mechanisms among three chlorinated organic solvents. *Toxicol Appl Pharmacol* 255(1):113-126.
- Bale AS, Meacham CA, Benignus VA, et al. 2005. Volatile organic compounds inhibit human and rat neuronal nicotinic acetylcholine receptors expressed in *Xenopus* oocytes. *Toxicol Appl Pharmacol* 205(1):77-88.
- Barber LB II, Thurman EM, Schroeder MP. 1988. Long-term fate of organic micropollutants in sewage-contaminated groundwater. *Environ Sci Technol* 22:205-211.
- Barkley J, Bunch J, Bursey JT, et al. 1980. Gas chromatography/mass spectrometry computer analysis of volatile halogenated hydrocarbons in man and his environment. A multimedia environmental study. *Biomed Mass Spectrom* 7:139-147.
- Barnes DG, Dourson M. 1988. Reference dose (RfD): Description and use in health risk assessments. *Regul Toxicol Pharmacol* 8(4):471-486.
- Barrows ME, Petrocelli SR, Macek KJ, et al. 1980. Bioconcentration and elimination of selected water pollutants by bluegill sunfish (*Lepomis macrochirus*). In: Haque R, ed. Dynamics, exposure and hazard assessment of toxic chemicals. Ann Arbor, MI: Ann Arbor Science, 379-392.
- Bartels MJ. 1994. Quantitation of the tetrachloroethylene metabolite N-acetyl-S-(trichlorovinyl) cysteine in rat urine via negative ion chemical ionization gas chromatography/tandem mass spectrometry. *Biol Mass Spectrom* 23:689-694.
- Bartsch H, Malaveille C, Barbin A, et al. 1979. Mutagenic and alkylating metabolites of haloethylenes, chlorobutadienes, and dichlorobutenes produced by rodent or human liver tissues. *Arch Toxicol* 41:249-277.
- Bayer CW, Black MS. 1987. Capillary chromatographic analysis of volatile organic compounds in the indoor environment. *J Chromatogr Sci* 25(2):60-64.
- Baykut G, Voigt A. 1992. Spray extraction of volatile organic compounds from aqueous systems into the gas phase for gas chromatography/mass spectrometry. *Anal Chem* 64:677-681.
- Begerow J, Jermann E, Keles T, et al. 1996. Screening method for the determination of 28 volatile organic compounds in indoor and outdoor air at environmental concentrations using dual-column capillary gas chromatography with tandem electron-capture-flame ionization detection. *J Chromatogr A* 749(1-2):181-191.

9. REFERENCES

- Benane SG, Blackman CF, House DE. 1996. Effect of perchloroethylene and its metabolites on intercellular communication in clone 9 rat liver cells. *J Toxicol Environ Health* 48(5):427-437.
- Benfenati E, Facchini G, Pierucci P, et al. 1996. Identification of organic contaminants in leachates from industrial waste landfills. *Trends Anal Chem* 15(8):305-310.
- Benignus VA, Boyes WK, Geller AM, et al. 2009. Long-term perchloroethylene exposure: A meta-analysis of neurobehavioral deficits in occupationally and residentially exposed groups. *J Toxicol Environ Health A* 72(13):824-831.
- Bentley P, Calder I, Elcombe C, et al. 1993. Hepatic peroxisome proliferation in rodents and its significance for humans. *Food Chem Toxicol* 31(11):857-907.
- Bergamaschi E, Mutti A, Bocchi MC, et al. 1992. Rat model of perchloroethylene-induced renal dysfunctions. *Environ Res* 59:427-439.
- Berger GS, ed. 1994. Epidemiology of endometriosis. In: *Endometriosis: Modern surgical management of endometriosis*. New York, NY: Springer-Verlag, 3-7.
- Berger T, Horner CM. 2003. *In vivo* exposure of female rats to toxicants may affect oocyte quality. *Reprod Toxicol* 17(3):273-281.
- Berman E, Schlicht M, Moser VC, et al. 1995. A multidisciplinary approach to toxicological screening: I. Systemic toxicity. *J Toxicol Environ Health* 45:127-143.
- *Birner G, Richling C, Henschler D, et al. 1994. Metabolism of tetrachloroethylene in rats: Identification of N-(dichloroacetyl)-L-lysine and N-(trichloroacetyl)-L-lysine as protein adducts. *Chem Res Toxicol* 7:724-732.
- Birner G, Rutkowska A, Dekant W. 1996. N-Acetyl-S-(1,2,2-trichlorovinyl)-L-cysteine and 2,2,2-trichloroethanol: Two novel metabolites of tetrachloroethene in humans after occupational exposure. *Drug Metab Dispos* 24(1):41-48.
- Blair A, Decoufle P, Grauman D. 1979. Causes of death among laundry and dry cleaning workers. *Am J Public Health* 69:508-511.
- Blair A, Petralia SA, Stewart PA. 2003. Extended mortality follow-up of a cohort of dry-cleaners. *Ann Epidemiol* 13(1):50-56.
- Blair A, Stewart PA, Tolbert PE, et al. 1990. Cancer and other causes of death among a cohort of dry cleaners. *Br J Ind Med* 47:162-168.
- Blount BC, Kobelski RJ, McElprang DO, et al. 2006. Quantification of 31 volatile organic compounds in whole blood using solid-phase microextraction and gas chromatography-mass spectrometry. *J Chromatogr B* 832(2):292-301.
- Boekhold AJ, Van Der Schee HA, Kaandorp BH. 1989. [Rapid gas-chromatographic determination of trichloroethylene and/or tetrachloroethylene in lettuce by direct head-space analysis]. *Z Lebensm Unters Forsch* 189:550-553. (German)

9. REFERENCES

- Bogen KT, Colston BW Jr., Machicao LK. 1992. Dermal absorption of diluted aqueous chloroform, trichloroethylene, and tetrachloroethylene in hairless guinea pigs. *Fundam Appl Toxicol* 18:30-39.
- Boice JD, Jr., Marano DE, Fryzek JP, et al. 1999. Mortality among aircraft manufacturing workers. *Occup Environ Med* 56(9):581-597.
- Bois FY, Zeise L, Tozer TN. 1990. Precision and sensitivity of pharmacokinetic models for cancer risk assessments: Tetrachloroethylene in mice, rats, and humans. *Toxicol Appl Pharmacol* 102:300-315.
- Bosco MG, Figa-Talamanca I, Salerno S. 1986. Health and reproductive status of female workers in dry cleaning shops. *Int Arch Occup Environ Health* 59:295-301.
- Bouwer EJ, McCarty PL. 1982. Removal of trace chlorinated organic compounds by activated carbon and fixed-film bacteria. *Environ Sci Technol* 16:836-843.
- Bouwer EJ, Rittmann BE, McCarty PL. 1981. Anaerobic degradation of halogenated 1- and 2-carbon organic compounds. *Environ Sci Technol* 15:596-599.
- Bove FJ, Fulcomer MC, Klotz JB. 1995. Public drinking water contamination and birth outcomes. *Am J Epidemiol* 141:850-862.
- Bove FJ, Ruckart PZ, Maslia M, et al. 2014a. Evaluation of mortality among marines and navy personnel exposed to contaminated drinking water at USMC Base Camp Lejeune: A retrospective cohort study. *Environ Health* 13(1):10. <http://doi.org/10.1186/1476-069X-13-10>.
- Bove FJ, Ruckart PZ, Maslia M, et al. 2014b. Mortality study of civilian employees exposed to contaminated drinking water at USMC Base Camp Lejeune: A retrospective cohort study. *Environ Health* 13:68. <http://doi.org/10.1186/1476-069x-13-68>.
- Boverhof DR, Krieger SM, Hotchkiss JA, et al. 2013. Assessment of the immunotoxic potential of trichloroethylene and perchloroethylene in rats following inhalation exposure. *J Immunotoxicol* 10(3):311-320.
- Boyes WK, Bercegeay M, Oshiro WM, et al. 2009. Acute perchloroethylene exposure alters rat visual-evoked potentials in relation to brain concentrations. *Toxicol Sci* 108(1):159-172.
- Briving C, Jacobson I, Hamberger A, et al. 1986. Chronic effects of perchloroethylene and trichloroethylene on the gerbil brain amino acids and glutathione. *Neurotoxicology* 7:101-108.
- Brodkin CA, Daniell W, Checkoway H, et al. 1995. Hepatic ultrasonic changes in workers exposed to perchloroethylene. *Occup Environ Med* 52:679-685.
- *Bronstein AC, Currence PL. 1988. Flammable liquids. In: *Emergency care for hazardous materials exposure*. Washington, DC: The C.V. Mosby Company, 93-94.
- Bronzetti G, Bauer C, Corsi R, et al. 1983. Genetic and biochemical studies on perchloroethylene *in vitro* and *in vivo*. *Mutat Res* 116:323-331.
- *Brouwer M, Koeman T, van den Brandt PA, et al. 2015. Occupational exposures and Parkinson's disease mortality in a prospective Dutch cohort. *Occup Environ Med* 72(6):448-455. [10.1136/oemed-2014-102209](https://doi.org/10.1136/oemed-2014-102209).

9. REFERENCES

- Brown DP, Kaplan SD. 1987. Retrospective cohort mortality study of dry cleaning workers using perchloroethylene. *J Occup Med* 29:535-541.
- Brown RP, Delp MD, Lindstedt SL, et al. 1997. Physiologically parameter values for physiologically based pharmacokinetic models. *Toxicol Ind Health* 13(4):407-484.
- Brugnone F, Perbellini L, Giuliari C, et al. 1994. Blood and urine concentrations of chemical pollutants in the general population. *Med Lav* 85(5):370-389.
- Buben JA, O'Flaherty EJ. 1985. Delineation of the role of metabolism in the hepatotoxicity of trichloroethylene and perchloroethylene: A dose-effect study. *Toxicol Appl Pharmacol* 78:105-122.
- Budavari S, ed. 1989. Tetrachloroethylene. In: *The Merck index*. 11th ed. Rahway, NJ: Merck and Co., Inc., 1449.
- Bukowski J. 2014. Critical review of the epidemiologic literature regarding the association between congenital heart defects and exposure to trichloroethylene. *Crit Rev Toxicol* 44(7):581-589. 10.3109/10408444.2014.910755.
- Bundschuh I, Herbot C, Fels LM, et al. 1993. Renal fibronectin excretion as a marker for renal environmental toxins. *Contrib Nephrol* 101:177-184.
- Burk T, Zarus G. 2013. Community exposures to chemicals through vapor intrusion: A review of past Agency for Toxic Substances and Disease Registry public health evaluations. *J Environ Health* 75(9):36-41.
- Byczkowski JZ, Fisher JW. 1994. Lactational transfer of tetrachloroethylene in rats. *Risk Anal* 14(3):339-349.
- Byczkowski JZ, Fisher JW. 1995. A computer program linking physiologically based pharmacokinetic model with cancer risk assessment for breast-fed infants. *Comput Methods Programs Biomed* 46:155-163.
- Byczkowski JZ, Kinkead ER, Leahy HF, et al. 1994. Computer simulation of the lactational transfer of tetrachloroethylene in rats using a physiologically based model. *Toxicol Appl Pharmacol* 125:228-236.
- Byers VS, Levin AS, Ozonoff DM, et al. 1988. Association between clinical symptoms and lymphocyte abnormalities in a population with chronic domestic exposure to industrial solvent-contaminated domestic water supply and a high incidence of leukemia. *Cancer Immunol Immunother* 27:77-81.
- Cabirol N, Jacob F, Perrier J, et al. 1998. Complete degradation of high concentrations of tetrachloroethylene by a methanogenic consortium in a fixed-bed reactor. *J Biotechnol* 62(2):133-141.
- Cabirol N, Perrier J, Jacob F, et al. 1996. Role of methanogenic and sulfate-reducing bacteria in the reductive dechlorination of tetrachloroethylene in mixed culture. *Bull Environ Contam Toxicol* 56(5):817-824.
- Cai S-X, Huang M-Y, Chen Z, et al. 1991. Subjective symptom increase among dry cleaning workers exposed to tetrachloroethylene vapor. *Ind Health* 29:111-121.

9. REFERENCES

- Cal EPA. 1999. Public health goal for trichloroethylene in drinking water. California Environmental Protection Agency, Office of Environmental Health Hazard Assessment.
- Cal EPA. 2001. Public health goal for tetrachloroethylene in drinking water. California Environmental Protection Agency, Office of Environmental Health Hazard Assessment.
- Callen DF, Wolf CR, Philpot RM. 1980. Cytochrome P-450 mediated genetic activity and cytotoxicity of seven halogenated aliphatic hydrocarbons in *Saccharomyces cerevisiae*. *Mutat Res* 77:55-63.
- Calvert GM, Ruder AM, Petersen MR. 2011. Mortality and end-stage renal disease incidence among dry cleaning workers. *Occup Environ Med* 68(10):709-716.
- *Cano MI, Pollan M. 2001. Non-Hodgkin's lymphomas and occupation in Sweden. *Int Arch Occup Environ Health* 74(6):443-449.
- Carney EW, Thorsrud BA, Dugard PH, et al. 2006. Developmental toxicity studies in Crl:CD (SD) rats following inhalation exposure to trichloroethylene and perchloroethylene. *Birth Defects Res B Dev Reprod Toxicol* 77(5):405-412.
- Carpenter CP. 1937. The chronic toxicity of tetrachloroethylene. *J Ind Hyg Toxicol* 19:323-336.
- Carwile JL, Mahalingaiah S, Winter MR, et al. 2014. Prenatal drinking-water exposure to tetrachloroethylene and ischemic placental disease: A retrospective cohort study. *Environ Health* 13:72. 10.1186/1476-069x-13-72.
- Cavalleri A, Gobba F. 1998. Reversible color vision loss in occupational exposure to metallic mercury. *Environ Res* 77:173-177.
- Cavalleri A, Gobba F, Paltrinieri M, et al. 1994. Perchloroethylene exposure can induce colour vision loss. *Neurosci Lett* 179:162-166.
- CDC. 2015. Fourth national report on human exposure to environmental chemicals, updated tables (February 2015). Centers for Disease Control and Prevention. http://www.cdc.gov/biomonitoring/pdf/FourthReport_UpdatedTables_Feb2015.pdf. December 14, 2015.
- CDPHE. 2002. Fact sheet. Dry cleaners and PCE. Colorado Department of Public Health and Environment.
- CDPHE. 2006. Dry cleaner remediation guidance document. Colorado Department of Public Health and Environment.
- Cederberg H, Henriksson J, Binderup ML. 2010. DNA damage detected by the alkaline comet assay in the liver of mice after oral administration of tetrachloroethylene. *Mutagenesis* 25(2):133-138.
- CEPA. 1993. Canadian Environmental Protection Act. Priority substances list assessment report: Tetrachloroethylene. Government of Canada, Environment Canada, Health Canada, 10-14, 29-30.
- CEPA. 2001. Public health goal for tetrachloroethylene in drinking water. California Environmental Protection Agency, Office of Environmental Health Hazard Assessment.

9. REFERENCES

- Chai JC, Miura N. 2004. Field vapor extraction test and long-term monitoring at a PCE contaminated site. *J Hazard Mater* 110:85-92.
- Chang YC, Hatsu M, Jung K, et al. 1998. Degradation of a variety of halogenated aliphatic compounds by an anaerobic mixed culture. *J Ferment Bioengineer* 86(4):410-412.
- Chang YC, Ikeutsu K, Toyama T, et al. 2011. Isolation and characterization of tetrachloroethylene- and cis-1,2-dichloroethylene-dechlorinating propionibacteria. *J Ind Microbiol Biotechnol* 38(10):1667-1677.
- Chang YM, Tai CF, Yang SC, et al. 2003. A cohort mortality study of workers exposed to chlorinated organic solvents in Taiwan. *Ann Epidemiol* 13(9):652-660.
- *Chang YM, Tai CF, Yang SC, et al. 2005. Cancer incidence among workers potentially exposed to chlorinated solvents in an electronics factory. *J Occup Health* 47(2):171-180.
- Chapman JAW, Connolly JG, Rosenbaum L. 1981. Occupational bladder cancer: A case-control study. In: Connolly JE, ed. *Carcinoma of the bladder*. New York, NY: Raven Press, 45-54.
- Charbotel B, Massardier-Pilonchery A, Fort E, et al. 2013. Occupational trichloroethylene exposure and cervical pathology: A case-control study. *Ann Occup Hyg* 57(3):407-416. 10.1093/annhyg/mes075.
- Chaudhuri RN, Mukerji AK. 1947. Death following administration of tetrachloroethylene. *Ind Med Gaz* 82:115-116.
- C&EN. 1994. Facts and figures for the chemical industry. *Chemical and Engineering News*, July 4, 1994.
- Chemical Products Synopsis. 1985. Perchloroethylene. Mannsville Chemical Products Corp., Cortland, NY.
- Chemical Profile. 1995. Perchloroethylene. *Chemical Marketing Reporter*, March 13, 1995.
- Chen CW, Blancato JN. 1987. Role of pharmacokinetic modeling in risk assessment: Perchloroethylene as an example. In: *Pharmacokinetics in risk assessment, drinking water and health*. Vol. 8. Washington, DC: National Academy of Sciences, National Research Council, 369-390.
- Chen KM, Dallas CE, Muralidhara S, et al. 1993. Analyses of volatile C₂ haloethanes and haloethenes in tissues: Sample preparation and extraction. *J Chromatogr* 612:199-208.
- Chen SJ, Wang JL, Chen JH, et al. 2002. Possible involvement of glutathione and p53 in trichloroethylene- and perchloroethylene-induced lipid peroxidation and apoptosis in human lung cancer cells. *Free Radic Biol Med* 33(4):464-472.
- Chiappini L, Delery L, Leoz E, et al. 2009. A first French assessment of population exposure to tetrachloroethylene from small dry-cleaning facilities. *Indoor Air* 19:226-233.
- Chilton PJ, Lawrence AR, Barker A. 1990. Chlorinated solvents in chalk aquifers: Some preliminary observations on behavior and transport. In: Telford T, ed. *International Chalk Symposium*. London, England, 605-610.

9. REFERENCES

- Chiu WA, Ginsberg GL. 2011. Development and evaluation of a harmonized physiologically based pharmacokinetic (PBPK) model for perchloroethylene toxicokinetics in mice, rats, and humans. *Toxicol Appl Pharmacol* 253(3):203-234.
- Chiu WA, Micallef S, Monster AC, et al. 2007. Toxicokinetics of inhaled trichloroethylene and tetrachloroethylene in humans at 1 ppm: Empirical results and comparisons with previous studies. *Toxicol Sci* 95(1):23-36.
- Chiu WA, Okino MS, Evans MV. 2009. Characterizing uncertainty and population variability in the toxicokinetics of trichloroethylene and metabolites in mice, rats, and humans using an updated database, physiologically based pharmacokinetic (PBPK) model, and Bayesian approach. *Toxicol Appl Pharmacol* 241:36-60.
- Chodola GR, Biswas N, Bewtra JK, et al. 1989. Fate of selected volatile organic substances in aqueous environment. *Water Pollut Res J Can* 24:119-142.
- Christensen JM, Rasmussen K, Koppen B. 1988. Automatic headspace gas chromatographic method for the simultaneous determination of trichloroethylene and metabolites in blood and urine. *J Chromatogr* 442:317-323.
- Christensen KY, Vizcaya D, Richardson H, et al. 2013. Risk of selected cancers due to occupational exposure to chlorinated solvents in a case-control study in Montreal. *J Occup Environ Med* 55(2):198-208. [10.1097/JOM.0b013e3182728eab](https://doi.org/10.1097/JOM.0b013e3182728eab).
- Cichocka D, Nikolausz M, Haest PJ, et al. 2010. Tetrachloroethene conversion to ethene by a dehalococoides-containing enrichment culture from Bitterfeld. *FEMS Microbiol Ecol* 72(2):297-310.
- Cichocki JA, Furuya S, Konganti K, et al. 2017a. Impact of nonalcoholic fatty liver disease on toxicokinetics of tetrachloroethylene in mice. *J Pharmacol Exp Ther* 361(1):17-28. <http://doi.org/10.1124/jpet.116.238790>.
- Cichocki JA, Furuya S, Luo YS, et al. 2017b. Nonalcoholic fatty liver disease is a susceptibility factor for perchloroethylene-induced liver effects in mice. *Toxicol Sci* 159(1):102-113. <http://doi.org/10.1093/toxsci/kfx120>.
- Cichocki JA, Furuya S, Venkatratnam A, et al. 2017c. Characterization of variability in toxicokinetics and toxicodynamics of tetrachloroethylene using the collaborative cross mouse population. *Environ Health Perspect* 125(5):057006. <http://doi.org/10.1289/ehp788>.
- Cichocki JA, Guyton KZ, Guha N, et al. 2016. Target organ metabolism, toxicity, and mechanisms of trichloroethylene and perchloroethylene: Key similarities, differences, and data gaps. *J Pharmacol Exp Ther* 359(1):110-123. <http://doi.org/10.1124/jpet.116.232629>.
- Class TH, Ballschmiter K. 1986. Chemistry of organic traces in air. VI: Distribution of chlorinated C1-C4 hydrocarbons in air over the northern and southern Atlantic Ocean. *Chemosphere* 15:413-427.
- Clewell HJ, Andersen ME. 1985. Risk assessment extrapolations and physiological modeling. *Toxicol Ind Health* 1(4):111-131.
- Clewell HJ, Gentry PR, Covington TR, et al. 2004. Evaluation of the potential impact of age- and gender-specific pharmacokinetic differences on tissue dosimetry. *Toxicol Sci* 79(2):381-393.

9. REFERENCES

- CMR. 2008. Chemical profile: Perchloroethylene. Chemical Marketing Reporter. <http://www.icis.com/Articles/2008/08/25/9150899/chemical-profile-perchloroethylene.html>. March 11, 2013.
- Cohn P, J Klotz, F Bove, et al. 1994. Drinking water contamination and the incidence of leukemia and non-Hodgkin's lymphoma. *Environ Health Perspect* 102(6-7):556-561.
- Coler HR, Rossmiller HR. 1953. Tetrachloroethylene exposure in a small industry. *AMA Arch Ind Hyg Occup Med* 8:227-233.
- *Corbin M, McLean D, Mannetje A, et al. 2011. Lung cancer and occupation: A New Zealand cancer registry-based case-control study. *Am J Ind Med* 54(2):89-101. 10.1002/ajim.20906.
- Cornish HH, Adefuin J. 1966. Ethanol potentiation of halogenated aliphatic solvent toxicity. *Am Ind Hyg Assoc J* 27:57-61.
- Cornish HH, Ling BP, Barth ML. 1973. Phenobarbital and organic solvent toxicity. *Am Ind Hyg Assoc J* 34:487-492.
- Costa AK, Ivanetich KM. 1980. Tetrachloroethylene metabolism by the hepatic microsomal cytochrome P-450 system. *Biochem Pharmacol* 29:2863-2869.
- Costa LG, Aschner M, Vitalone A, et al. 2004. Developmental neuropathology of environmental agents. *Ann Rev Pharmacol Toxicol* 44:87-110. 10.1146/annurev.pharmtox.44.101802.121424.
- Cramer PH, Boggess KE, Hosenfeld JM, et al. 1988. Determination of organic chemicals in human whole blood: Preliminary method development for volatile organics. *Bull Environ Contam Toxicol* 40(4):612-618.
- Cupitt LT. 1987. Atmospheric persistence of eight air toxics. Environmental Protection Agency, Research Triangle Park, NC, Atmospheric Sciences Research Lab. EPA600S387004.
- Daft JL. 1988. Rapid determination of fumigant and industrial chemical residues in food. *J Assoc Off Anal Chem* 71:748-760.
- Daft JL. 1989. Determination of fumigants and related chemicals in fatty and nonfatty foods. *J Agric Food Chem* 37:560-564.
- *Dallas CE, Chen XM, Muralidhara S, et al. 1994a. Use of tissue disposition data from rats and dogs to determine species differences in input parameters for physiological model for perchloroethylene. *Environ Res* 67:54-67.
- Dallas CE, Chen XM, Muralidhara S, et al. 1995. Physiologically based pharmacokinetic model useful in prediction of the influence of species, dose, and exposure route on perchloroethylene pharmacokinetics. *J Toxicol Environ Health* 44:301-317.
- Dallas CE, Chen XM, O'Barr K, et al. 1994b. Development of a physiologically based pharmacokinetic model for perchloroethylene using tissue concentration - time data. *Toxicol Appl Pharmacol* 128:50-59.

9. REFERENCES

- Dallas CE, Muralidhara S, Chen XM, et al. 1994c. Use of a physiologically based model to predict systemic uptake and respiratory elimination of perchloroethylene. *Toxicol Appl Pharmacol* 128:60-68.
- *Daniel JW. 1963. The metabolism of ³⁶Cl-labeled trichloroethylene and tetrachloroethylene in the rat. *Biochem Pharmacol* 12:795-802.
- *Dart RC. 2004. Halogenated hydrocarbons-halogenated solvents. In: *Medical toxicology*. 3rd ed. Philadelphia, PA: Lippincott Williams and Wilkins, 1339-1341.
- Dawson HE, McAlary T. 2009. A compilation of statistics for VOCs from post-1990 indoor air concentration studies in North American residences unaffected by subsurface vapor intrusion. *Ground Water Monit Remed* 29(1):60-69.
- De Ceaurriz J, Desiles JP, Bonnet P, et al. 1983. Concentration-dependent behavioral changes in mice following short-term inhalation exposure to various industrial solvents. *Toxicol Appl Pharmacol* 67:383-389.
- Dehon B, Humbert L, Devisme L, et al. 2000. Tetrachloroethylene and trichloroethylene fatality: Case report and simple headspace SPME-capillary gas chromatographic determination in tissues. *J Anal Toxicol* 24(1):22-26.
- Deipser A, Stegmann, R. 1994. The origin and fate of volatile trace components in municipal solid waste landfills. *Waste Manag Res* 12:129-139.
- Dekant W. 1986. Metabolic conversion of tri- and tetrachloroethylene: Formation and deactivation of genotoxic intermediates. In: Chambers PL, Gehring P, Sakai F, eds. *New concepts and developments in toxicology*. Elsevier Science Publishers, 211-221.
- Dekant W, Martens G, Vamavakas S, et al. 1987. Bioactivation of tetrachloroethylene. Role of glutathione s-transferase-catalyzed conjugation versus cytochrome P-450 dependent phospholipid alkylation. *Drug Metab Dispos* 15:702-709.
- Dekant W, Metzler M, Henschler D. 1986. Identification of S-1,2,2-trichlorovinyl-n-acetylcysteine as a urinary metabolite of tetrachloroethylene: Bioactivation through glutathione conjugation as a possible explanation of its nephrocarcinogenicity. *J Biochem Toxicol* 1:57-72.
- Delfino RJ, Gong HH, Linn WS, et al. 2003. Respiratory symptoms and peak expiratory flow in children with asthma in relation to volatile organic compounds in exhaled breath and ambient air. *J Expo Anal Environ Epidemiol* 13:348-363.
- De Raat K. 2003. The Nordic Expert Group for criteria documentation of health risks from chemicals and the Dutch Expert Committee on occupational standards. Tetrachloroethylene (PER). Stockholm, Sweden: Nordic Council of Ministers.
- DeWulf J, Van Langehove H, Heireman B. 1998. The air/water exchange of volatile organic compounds from waters in the transient and turbulent regime. *Water Res* 32(7):2106-2112.
- DHHS. 1994. Seventh annual report on carcinogens. Summary 1994. U.S. Department of Health and Human Services, Public Health Service. Report to the National Institute of Environmental Health Sciences, Research Triangle Park, NC, by Technical Resources, Inc., Rockville, MD, 372-376.

9. REFERENCES

- Diamadopoulos DS, Koukouraki E. 1998. The effect of humic acid on the transport of volatile chlorinated hydrocarbons in soil. *Water Res* 32(11):3325-3330.
- Dias CM, Menezes HC, Cardeal ZL. 2017. Use of exhaled air as an improved biomonitoring method to assess perchloroethylene short-term exposure. *Environ Res* 156:108-112.
- Dickson AG, Riley JP. 1976. The distribution of short-chain halogenated aliphatic hydrocarbons in some marine organisms. *Mar Pollut Bull* 7:167-169.
- Dietz EA, Singley KF. 1979. Determination of chlorinated hydrocarbons in water by headspace gas chromatography. *Anal Chem* 51(11):1809-1814.
- Dilling WL. 1977. Interphase transfer processes. II. Evaporation rates of chloromethanes, ethanes, ethylenes, propanes, and propylenes from dilute aqueous solutions. Comparisons with theoretical predictions. *Environ Sci Technol* 11:405-409.
- Dilling WL, Tefertiller NB, Kallos GJ. 1975. Evaporation rates and reactivities of methylene chloride, chloroform, 1,1,1-trichloroethane, trichloroethylene, tetrachloroethylene, and other chlorinated compounds in dilute aqueous solutions. *Environ Sci Technol* 9:833-838.
- DiStefano TD, Gossett JM, Zinder SH. 1991. Reductive dechlorination of high concentrations of tetrachloroethene to ethene by an anaerobic enrichment culture in the absence of methanogenesis. *Appl Environ Microbiol* 57(8):2287-2292.
- DOE. 2012. Protective action criteria (PAC). Oak Ridge, TN: U.S. Department of Energy and Subcommittee on Consequence Assessment and Protective Actions (SCAPA). <http://orise.orau.gov/emi/scapa/chem-pacs-teels/default.htm>. April 24, 2013.
- Dosemeci M, Cocco P, Chow WH. 1999. Gender differences in risk of renal cell carcinoma and occupational exposures to chlorinated aliphatic hydrocarbons. *Am J Ind Med* 36:54-59.
- Doust HG, Huang J. 1992. The fate and transport of hazardous chemicals in the subsurface environment. *Water Sci Technol* 25:169-176.
- Dow. 2008. Product safety assessment perchloroethylene. Dow Chemical Company.
- *Dow Chemical. 1983. Initial submission: Perchloroethylene solvent formulation: Acute toxicological properties and industrial handling hazards. Submitted to the U.S. Environmental Protection Agency under TSCA, Section 8e. OTS0574294.
- Doyle P, Roman E, Beral V, et al. 1997. Spontaneous abortion in dry cleaning workers potentially exposed to perchloroethylene. *Occup Environ Med* 54(12):848-853.
- Duh RW, Asal NR. 1984. Mortality among laundry and dry cleaning workers in Oklahoma. *Am J Public Health* 74:1278-1280.
- Dyksen JE, Hess AF III. 1982. Alternatives for controlling organics in groundwater supplies. *J Am Water Works Assoc* 74:394-402.

9. REFERENCES

- Earnest GS. 1995. In-depth survey report: Chemical exposures and fire safety in commercial dry cleaners at A-One Cleaners, Cincinnati, Ohio. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Report No. ECTB 201-18a.
- Earnest GS. 1996. Evaluation and control of perchloroethylene exposure during dry cleaning. *Appl Occup Environ Hyg* 11(2):125-132.
- Earnest GS, Spencer AB. 1995. In-depth survey report: Perchloroethylene exposures in commercial dry cleaners at Brown's Cleaners, Santa Monica, California. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Report No. ECTB 201-16a.
- Earnest GS, Moran AJ, Spencer AB, et al. 1995a. In-depth survey report: Control of perchloroethylene exposures in commercial dry cleaners at Appearance Plus Cleaners, Cincinnati, Ohio. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Report No. ECTB: 201-15a.
- Earnest GS, Moran AJ, Spencer AB. 1995b. In-depth survey report: Perchloroethylene exposures in commercial dry cleaners at Tuchman Cleaners (shop #27), Fishers, Indiana. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Report No. ECTB: 201-17a.
- Earnest GS, Spencer AB, Smith SS, et al. 1995c. In-depth survey report: Perchloroethylene exposures and ergonomic risk factors in commercial dry cleaners at Tuchman Cleaners (shop #24), Carmel, Indiana. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Report No. ECTB: 201-19a.
- Ebrahim AS, Babakrishnan K, Sakthisekaran D. 1996. Perchloroethylene-induced alterations in glucose metabolism and their prevention by 2-deoxy-D-glucose and vitamin E in mice. *J Appl Toxicol* 16(4):339-348.
- Ebrahim AS, Babu E, Thirunavukkarasu C, et al. 2001. Protective role of vitamin E, 2-deoxy-D-glucose, and taurine on perchloroethylene induced alterations in ATPases. *Drug Chem Toxicol* 24(4):429-437.
- Ebrahim AS, Gopalakrishnan R, Murugesan A, et al. 1995. *In vivo* effect of vitamin E on serum and tissue glycoprotein levels in perchloroethylene induced cytotoxicity. *Mol Cell Biol* 144:13-18.
- Echeverria D, White RF, Sampaio C. 1995. A behavioral evaluation of PCE exposure in patients and dry cleaners: A possible relationship between clinical and preclinical effects. *J Occup Environ Med* 37(6):667-680.
- E.I. Dupont. 1969. Cardiac sensitization in dogs. Submitted to the U.S. Environmental Protection Agency under TSCA, Section 8d. OTS0514917.
- Ek CJ, Dziegielewska KM, Habgood MD, et al. 2012. Barriers in the developing brain and neurotoxicology. *Neurotoxicology* 33(3):586-604. 10.1016/j.neuro.2011.12.009.
- *Ellenhorn MJ, Barceloux DG. 1988. Airborne toxins. In: *Medical toxicology: Diagnosis and treatment of human poisoning*. New York, NY: Elsevier, 841-843.

9. REFERENCES

- Emara AM, Abo El-Noor MM, Hassan NA, et al. 2010. Immunotoxicity and hematotoxicity induced by tetrachloroethylene in Egyptian dry cleaning workers. *Inhal Toxicol* 22(2):117-124.
- Emmert B, Bunger J, Keuch K, et al. 2006. Mutagenicity of cytochrome P450 2E1 substrates in the Ames test with the metabolic competent *S. typhimurium* strain YG7108pin3ERb₅. *Toxicology* 228(1):66-76.
- Entz RC, Diachenko GW. 1988. Residues of volatile halocarbons in margarines. *Food Addit Contam* 5:267-276.
- Entz RC, Hollifield HC. 1982. Headspace gas chromatographic analysis of foods for volatile halocarbons. *J Agric Food Chem* 30:84-88.
- EPA. 1980. Hazardous waste management system: Identification and listing of hazardous waste: Interim rule. U.S. Environmental Protection Agency. Fed Regist 45:78530-78550.
- EPA. 1982a. Methods for organic chemical analysis of municipal and industrial wastewater. Test method: Purgeable halocarbons - Method 601 and Purgeables - Method 624. Cincinnati, OH: U.S. Environmental Protection Agency, Environmental Monitoring and Support Laboratory. EPA600/482057, 601-1 to 601-10 and 624-1 to 624-12.
- EPA. 1982b. Test methods for evaluating solid waste. Physical/chemical methods. Method 8010-Halogenated volatile organics. Washington, DC: U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response, 8010-1 to 8010-12.
- EPA. 1985. Designation, reportable quantities, and notification. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 302.
- EPA. 1987. Health advisories for 25 organics. U.S. Environmental Protection Agency, Office of Drinking Water. PB87-2355781AS.
- EPA. 1988. Toxic chemical release reporting: Community right-to-know. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 372. Fed Regist 53(30):4500-4554.
- EPA. 1990. Interim methods for development of inhalation reference concentrations. Washington, DC: U.S. Environmental Protection Agency, Office of Health and Environmental Assessment, Office of Research and Development. EPA600/890066A. PB90238890.
- EPA. 1991. Hazardous waste and superfund programs; amendment to preambles published at 54 FR 33418 (August 14, 1989) and 54 FR 50968 (December 11, 1989). U.S. Environmental Protection Agency. Fed Regist 56:643-645.
- EPA. 1995. EPA Office of Compliance sector notebook project. Profile of the dry cleaning industry. Washington, DC: U.S. Environmental Protection Agency, Office of Compliance, Office of Enforcement and Compliance Assurance. EPA310R95001.
<http://www.epa.gov/compliance/resources/publications/assistance/sectors/notebooks/dryclng.pdf>.
October 14, 2014.

9. REFERENCES

- EPA. 1996. Air quality; revision to definition of volatile organic compounds- exclusion of perchloroethylene. Fed Regist 61(26):4587-4591. <https://www.gpo.gov/fdsys/pkg/FR-1996-02-07/html/96-2495.htm>. June 07, 2018.
- EPA. 1997. Special report on environmental endocrine disruption: An effects assessment and analysis. Washington, DC. U.S. Environmental Protection Agency. EPA630R96012.
- EPA. 1998. RCRA waste minimization PBT priority chemical list. U.S. Environmental Protection Agency. Fed Regist 63 FR 60332. <http://www.gpo.gov/fdsys>. April 24, 2013.
- EPA. 1999a. Compendium of methods for the determination of toxic organic compounds in ambient air. Second edition. Compendium Method T0-15. Determination of volatile organic compounds (VOCs) in air collected in specially-prepared canisters and analyzed by gas chromatography/mass spectrometry (GC/MS). Cincinnati, OH: U.S. Environmental Protection Agency, Office of Research and Development. EPA625R96010b.
- EPA. 1999b. Compendium of methods for the determination of toxic organic compounds in ambient air. Second edition. Compendium Method T0-17. Determination of volatile organic compounds in ambient air using active sampling onto sorbent tubes. Cincinnati, OH: U.S. Environmental Protection Agency, Office of Research and Development. EPA625R96010b.
- EPA. 2005. Toxic chemical release inventory reporting forms and instructions: Revised 2004 version. Section 313 of the Emergency Planning and Community Right-to-Know Act (Title III of the Superfund Amendments and Reauthorization Act of 1986).
- EPA. 2006. National perchloroethylene air emission standards for dry cleaning facilities. Final rule. Fed Regist 71(144):42723-42746.
- EPA. 2007. Definition of hazardous waste. U.S. Environmental Protection Agency 40 CFR 261.3, 34-41. <http://www.gpo.gov/fdsys/pkg/CFR-2007-title40-vol25/pdf/CFR-2007-title40-vol25-sec261-3.pdf> September 9, 2014.
- EPA. 2009a. Drinking water contaminant candidate list. U.S. Environmental Protection Agency. Fed Regist 74 FR 51850:51850-51862. <http://www.gpo.gov/fdsys>. April 24, 2013.
- EPA. 2009b. National primary drinking water regulations. Washington, DC: U.S. Environmental Protection Agency, Office of Ground Water and Drinking Water. EPA816F090004. <http://water.epa.gov/drink/contaminants/upload/mcl-2.pdf>. April 24, 2013.
- EPA. 2009c. National recommended water quality criteria. Washington, DC: U.S. Environmental Protection Agency, Office of Water, Office of Science and Technology. <http://water.epa.gov/scitech/swguidance/standards/criteria/current/index.cfm>. April 24, 2013.
- EPA. 2009d. National primary drinking water regulations. Washington, DC: U.S. Environmental Protection Agency, Office of Ground Water and Drinking Water. EPA816F090004. <http://water.epa.gov/drink/contaminants/upload/mcl-2.pdf>. April 24, 2013.
- EPA. 2011. Background indoor air concentrations of volatile organic compounds in North American residences (1990-2005): A compilation of statistics for assessing vapor intrusion. Washington, DC: U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response. EPA530R10001.

9. REFERENCES

- EPA. 2012a. Toxicological review of tetrachloroethylene (perchloroethylene). Washington, DC: U.S. Environmental Protection Agency. https://cfpub.epg.gov/ncea/iris/iris_documents/documents/toxreviews/0106tr.pdf. May 30, 2018.
- EPA. 2012b. Designated as hazardous substances in accordance with section 311(b)(2)(a) of the Clean Water Act. U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 116.4. <http://www.gpo.gov/fdsys>. April 24, 2013.
- EPA. 2012c. Drinking water standards and health advisories. Washington, DC: U.S. Environmental Protection Agency, Office of Water. EPA822S12001. <http://water.epa.gov/drink/standards/hascience.cfm>. April 24, 2013.
- EPA. 2012d. Identification and listing of hazardous waste. Hazardous constituents. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 261, Appendix VIII. <http://www.gpo.gov/fdsys>. April 24, 2013.
- EPA. 2012e. Reportable quantities of hazardous substances designated pursuant to section 311 of the Clean Water Act. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 117.3. <http://www.gpo.gov/fdsys>. April 24, 2013.
- EPA. 2012f. Standards for owners and operators of hazardous waste TSD facilities. Groundwater monitoring list. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 264, Appendix IX. <http://www.gpo.gov/fdsys>. April 24, 2013.
- EPA. 2012g. Superfund, emergency planning, and community right-to-know programs. Designation, reportable quantities, and notifications. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 302.4. <http://www.gpo.gov/fdsys>. April 24, 2013.
- EPA. 2012h. Superfund, emergency planning, and community right-to-know programs. Toxic chemical release reporting. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 372.65. <http://www.gpo.gov/fdsys>. April 24, 2013.
- EPA. 2012i. Superfund, emergency planning, and community right-to-know programs. Extremely hazardous substances and their threshold planning quantities. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 355, Appendix A. <http://www.gpo.gov/fdsys>. April 24, 2013.
- EPA. 2012j. Toxic Substances Control Act. Chemical lists and reporting periods. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 712.30. <http://www.gpo.gov/fdsys>. April 24, 2013.
- EPA. 2012k. Toxic Substances Control Act. Health and safety data reporting. U.S. Environmental Protection Agency. Code of Federal Regulations. 40 CFR 716.120. <http://www.gpo.gov/fdsys>. April 24, 2013.
- EPA. 2012l. Vapor intrusion database. http://www.epa.gov/oswer/vaporintrusion/vi_data.html. January 24, 2014.
- EPA. 2012m. Petroleum hydrocarbons and chlorinated hydrocarbons differ in their potential for vapor intrusion. Washington, DC: U.S. Environmental Protection Agency, Office of Underground Storage Tanks.

9. REFERENCES

- EPA. 2012n. Conceptual model scenarios for the vapor intrusion pathway. U.S. Environmental Protection Agency. EPA530R10003. <http://www.epa.gov/sites/production/files/2015-09/documents/vi-cms-v11final-2-24-2012.pdf>. February 24, 2016.
- EPA. 2013a. Acute exposure guideline levels (AEGs). Washington, DC: U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics. <http://www.epa.gov/oppt/aegl/>. April 24, 2013.
- EPA. 2013b. Hazardous air pollutants. Clean Air Act. U.S. Environmental Protection Agency. United States Code 42 USC 7412. <http://www.epa.gov/ttn/atw/orig189.html>. April 24, 2013.
- EPA. 2013c. National ambient air quality standards (NAAQS). Washington, DC: U.S. Environmental Protection Agency, Office of Air and Radiation. <http://www.epa.gov/air/criteria.html>. April 24, 2013.
- EPA. 2013d. Inert ingredients permitted for use in nonfood pesticide products. Washington, DC: U.S. Environmental Protection Agency. <http://iaspub.epa.gov/apex/pesticides/f?p=124:1>. April 24, 2013.
- EPA. 2013e. Master testing list. Washington, DC: U.S. Environmental Protection Agency, Office of Pollution and Prevention and Toxics. <http://www.epa.gov/opptintr/chemtest/pubs/mtl.html>. April 24, 2013.
- EPA. 2013f. Non-confidential 2006 IUR records by chemical, including manufacturing, processing and use information. U.S. Environmental Protection Agency. http://cfpub.epa.gov/iursearch/2006_iur_companyinfo.cfm?chemid=3752&outchem=both. July 02, 2013.
- EPA. 2013g. Dry cleaning emission standards. U.S. Environmental Protection Agency. <http://www.epa.gov/drycleaningrule/basic.html>. July 02, 2013.
- EPA. 2013h. Air toxics data. Tetrachloroethylene. Air toxics-data analysis. U.S. Environmental Protection Agency. <http://www.epa.gov/ttnamti1/toxdat.html#data>. March 19, 2013.
- EPA. 2015a. Chemical data access tool (CDAT). 2012 Chemical data reporting (CDR), ethene, 1,1,2,2-tetrachloro-. U.S. Environmental Protection Agency. http://java.epa.gov/oppt_chemical_search/. December 4, 2015.
- EPA. 2015b. Technology transfer network. Clearinghouse for inventories & emissions factors. National Emissions Inventory (NEI) air pollutant emissions trends data. U.S. Environmental Protection Agency. <http://www3.epa.gov/ttnchie1/trends/>. December 7, 2015.
- EPA. 2018. Air quality system (AQS). Data files. Annual summary data. U.S. Environmental Protection Agency. <https://www.epa.gov/aqs>. December 11, 2018.
- Eskenazi B, Fenster L, Hudes M, et al. 1991a. A study of the effect of perchloroethylene exposure on the reproductive outcomes of wives of dry cleaning workers. *Am J Ind Med* 20:593-600.
- Eskenazi B, Wyrobek AJ, Fenster L, et al. 1991b. A study of the effect of perchloroethylene exposure on semen quality in dry cleaning workers. *Am J Ind Med* 20:575-591.
- Everatt R, Slapsyte G, Mierauskiene J, et al. 2013. Biomonitoring study of dry cleaning workers using cytogenetic tests and the comet assay. *J Occup Environ Hyg* 10(11):609-621. 10.1080/15459624.2013.818238.

9. REFERENCES

- *Fagliano J, Berry M, Bove F, et al. 1990. Drinking water contamination and the incidence of leukemia: An ecologic study. *Am J Public Health* 80:1209-1212.
- FDA. 2013. Everything added to food in the United States (EAFUS). Washington, DC: U.S. Food and Drug Administration. <http://www.accessdata.fda.gov/scripts/fcn/fcnavigation.cfm?rpt=eafuslisting>. April 24, 2013.
- *Feldman AL, Johansson AL, Nise G, et al. 2011. Occupational exposure in parkinsonian disorders: A 43-year prospective cohort study in men. *Parkinsonism Relat Disord* 17(9):677-682. 10.1016/j.parkreldis.2011.06.009.
- Feldman RG, Chirico-Post J, Proctor SP. 1988. Blink reflex latency after exposure to tetrachloroethylene in well water. Pittsburgh, PA: Meeting on Epidemiology in Environmental Health held at the First International Symposium on Environmental Epidemiology, June 3-5, 1987. *Arch Environ Health* 42:143-148.
- Ferrario JB, Lawler GC, Deleon IR, et al. 1985. Volatile organic pollutants in biota and sediments of Lake Pontchartrain. *Bull Environ Contam Toxicol* 34:246-255.
- Ferroni C, Selis L, Mutti A, et al. 1992. Neurobehavioral and neuroendocrine effects of occupational exposure to perchloroethylene. *Neurotoxicology* 13:243-248.
- Fisher J, Mahle D, Bankstron L, et al. 1997. Lactational transfer of volatile chemicals in breast milk. *Am Ind Hyg Assoc J* 58(6):425-431.
- Folkes D, Wertz W, Kurtz J, et al. 2009. Observed spatial and temporal distributions of CVOCs at Colorado and New York vapor intrusion sites. *Ground Water Monit Remed* 29(1):70-80.
- Fomon SJ. 1966. Body composition of the infant: Part 1: The male reference infant. In: Faulkner F, ed. *Human development*. Philadelphia, PA: WB Saunders, 239-246.
- Fomon SJ, Haschke F, Ziegler EE, et al. 1982. Body composition of reference children from birth to age 10 years. *Am J Clin Nutr* 35(Suppl 5):1169-1175.
- Forand SP, Lewis-Michl EL, Gomez MI. 2012. Adverse birth outcomes and maternal exposure to trichloroethylene and tetrachloroethylene through soil vapor intrusion in New York State. *Environ Health Perspect* 120(4):616-621.
- Franchini I, Cavotorta A, Falzoi M, et al. 1983. Early indicators of renal damage in workers exposed to organic solvents. *Int Arch Occup Environ Health* 52:1-9.
- Frantz SW, Watanabe PG. 1983. Tetrachloroethylene: Balance and tissue distribution in male Sprague-Dawley rats by drinking water administration. *Toxicol Appl Pharmacol* 69:66-72.
- Fredriksson A, Danielsson BRG, Eriksson P. 1993. Altered behavior in adult mice orally exposed to tri- and tetrachloroethylene as neonates. *Toxicol Lett* 66:13-19.
- Freedman DL, Gossett JM. 1989. Biological reductive dechlorination of tetrachloroethylene and trichloroethylene to ethylene under methanogenic conditions. *Appl Environ Microbiol* 55:2144-2151.

9. REFERENCES

- Friberg L, Kylin B, Nystrom A. 1953. Toxicities of trichloroethylene and tetrachloroethylene and Fujiwara's pyridine-alkali reaction. *Acta Pharmacol Toxicol* 9:303-312.
- Friesel P, Milde G, Steiner B. 1984. Interactions of halogenated hydrocarbons with soils. *Fresenius Z Anal Chem* 319:160-164.
- Gallagher LG, Vieira VM, Ozonoff D, et al. 2011. Risk of breast cancer following exposure to tetrachloroethylene-contaminated drinking water in Cape Cod, Massachusetts: Reanalysis of a case-control study using a modified exposure assessment. *Environ Health* 10:47.
- Garetano G, Gochfeld M. 2000. Factors influencing tetrachloroethylene concentrations in residences above dry-cleaning establishments. *Arch Environ Health* 55(1):59-68.
- Gargas ML, Burgess RJ, Voisard DE, et al. 1986. Partition coefficients of low molecular weight volatile chemicals in various liquids and tissues. Wright-Patterson Air Force Base, OH: Armstrong Aerospace Medical Research Laboratory, Toxic Hazards Division, Biochemistry Branch.
- Gargas ML, Burgess RJ, Voisard DE, et al. 1989. Partition coefficients of low-molecular-weight volatile chemicals in various liquids and tissues. *Toxicol Appl Pharmacol* 98:87-99.
- Garnier R, Bedouin J, Pepin G, et al. 1996. Coin-operated dry cleaning machines may be responsible for acute tetrachloroethylene poisoning: Report of 26 cases including one death. *Clin Toxicol* 34(2):191-197.
- Gay BW Jr, Hanst PL, Bulfalini JJ, et al. 1976. Atmospheric oxidation of chlorinated ethylenes. *Environ Sci Technol* 10:58-67.
- Gearhart JM, Mahle DA, Greene RJ. 1993. Variability of physiologically based pharmacokinetic (PBPK) model parameters and their effects on PBPK model predictions in a risk assessment for perchloroethylene. *Toxicol Lett* 68:131-144.
- Gennari P, Naldi M, Motta R, et al. 1992. Gamma-glutamyltransferase isoenzyme pattern in workers exposed to tetrachloroethylene. *Am J Ind Med* 21:661-671.
- Germolec DR, Yang RSH, Ackermann MF, et al. 1989. Toxicology studies of a chemical mixture of 25 groundwater contaminants: II. Immunosuppression in B6C3F₁ mice. *Fundam Appl Toxicol* 13:377-387.
- Getz KD, Janulewicz PA, Rowe S, et al. 2012. Prenatal and early childhood exposure to tetrachloroethylene and adult vision. *Environ Health Perspect* 120(9):1327-1332.
- Ghantous H, Danielsson BRG, Dencker L, et al. 1986. Trichloroacetic acid accumulates in murine amniotic fluid after tri- and tetrachloroethylene inhalation. *Acta Pharmacol Toxicol* 58:105-114.
- Ghittori S, Imbraini M, Pezzagno G. 1987. The urinary concentration of solvents as a biological indicator of exposure: Proposal for the biological equivalent exposure limit for nine solvents. *Am Ind Hyg Assoc J* 48:786-790.
- *Gilboa SM, Desrosiers TA, Lawson C, et al. 2012. Association between maternal occupational exposure to organic solvents and congenital heart defects, National Birth Defects Prevention Study, 1997-2002. *Occup Environ Med* 69(9):628-635. 10.1136/oemed-2011-100536.

9. REFERENCES

- Gimmi T, Fluehler H, Studer B, et al. 1993. Transport of volatile chlorinated hydrocarbons in unsaturated aggregated media. *Water Air Soil Pollut* 68:291-305.
- Giovannini L, Guglielmi G, Casini T, et al. 1992. Effect of ethanol chronic use on hepatotoxicity in rats exposed to tetrachloroethylene. *Int J Tiss React* 14(6):281-285.
- Giwercman A, Carlsen E, Keiding N, et al. 1993. Evidence for increasing incidence of abnormalities of the human testis: A review. *Environ Health Perspect* 101(Supp 2):65-71.
- *Glass DC, Heyworth J, Thomson AK, et al. 2015. Occupational exposure to solvents and risk of breast cancer. *Am J Ind Med* 58(9):915-922. 10.1002/ajim.22478.
- Gobba F. 2000. Color vision: A sensitive indicator of exposure to neurotoxins. *Neurotoxicology* 21(5):857-862.
- Gobba F, Righi E, Fantuzzi G, et al. 1998. Two-year evolution of perchloroethylene-induced color-vision loss. *Arch Environ Health* 53(3):196-198.
- Gold LS, Milliken K, Stewart P, et al. 2010. Occupation and multiple myeloma: An occupation and industry analysis. *Am J Ind Med* 53:768-779.
- Gold LS, Stewart PA, Milliken K, et al. 2011. The relationship between multiple myeloma and occupational exposure to six chlorinated solvents. *Occup Environ Med* 68(6):391-399.
- Goldberg ME, Johnson HE, Pozzani UC, et al. 1964. Effect of repeated inhalation of vapors of industrial solvents on animal behavior. I. Evaluation of nine solvent vapors on pole-climb performance in rats. *Am Ind Hyg Assoc J* 25:369-375.
- Goldman SM, Quinlan PJ, Ross GW, et al. 2012. Solvent exposures and Parkinson disease risk in twins. *Ann Neurol* 71(6):776-784.
- Goldsworthy TL, Popp JA. 1987. Chlorinated hydrocarbon-induced peroxisomal enzyme activity in relation to species and organ carcinogenicity. *Toxicol Appl Pharmacol* 88:225-233.
- Goldsworthy TL, Lyght O, Burnett VL, et al. 1988. Potential rate of α -2 μ -globulin, protein droplet accumulation, and cell replication in the renal carcinogenicity of rats exposed to trichloroethylene, perchloroethylene, and pentachloroethane. *Toxicol Appl Pharmacol* 96:367-379.
- Gossett JM. 1987. Measurement of Henry's law constants for C₁ and C₂ chlorinated hydrocarbons. *Environ Sci Technol* 21:202-208.
- Green SM, Khan MF, Kaphalia BS, et al. 2001. Immunohistochemical localization of trichloroacylated protein adducts in tetrachloroethene-treated mice. *J Toxicol Environ Health A* 63(2):145-157.
- Green T. 1990. Species differences in carcinogenicity: The role of metabolism in human risk evaluation. *Teratog Carcinog Mutagen* 10:103-113.
- Green T, Odum J, Nash JA, et al. 1990. Perchloroethylene-induced rat kidney tumors: An investigation of the mechanisms involved and their relevance to humans. *Toxicol Appl Pharmacol* 103:77-89.

9. REFERENCES

- Gregersen P. 1988. Neurotoxic effects of organic solvents in exposed workers: Two controlled follow-up studies after 5.5 and 10.6 years. *Am J Ind Med* 14:681-701.
- Gregersen P, Angelso B, Nielsen TE, et al. 1984. Neurotoxic effects of organic solvents in exposed workers: An occupational neuropsychological and neurological investigation. *Am J Ind Med* 5:201-225.
- Greim H, Bonse G, Radwan Z, et al. 1975. Mutagenicity *in vitro* and potential carcinogenicity of chlorinated ethylenes as a function of metabolic oxirane formation. *Biochem Pharmacol* 24:2013-2017.
- Grob K, Frauenfelder C, Artho A. 1990. Uptake by foods of tetrachloroethylene, trichloroethylene, toluene, and benzene from air. *Z Lebensm Unters Forsch* 191:435-441.
- Guberan E, Fernandez J. 1974. Control of industrial exposure to tetrachloroethylene by measuring alveolar concentrations: Theoretical approach using a mathematical model. *Br J Ind Med* 31:159-167.
- Gulyas H, Hemmerling L. 1990. Tetrachloroethene air pollution originating from coin-operated dry cleaning establishments. *Environ Res* 53:90-99.
- Gummin DD. 2015. Hydrocarbons. In: Hoffman RS, Lewin NA, Goldfrank LR, et al., eds. *Goldfrank's toxicologic emergencies*. 10th ed. New York, NY: McGraw-Hill Education, 309-310, 1334.
- Guo Z, Tichenor BA, Mason MA, et al. 1990. The temperature dependence of the emission of perchloroethylene from dry cleaned fabrics. *Environ Res* 52:107-115.
- Guyton KZ, Hogan KA, Scott CS, et al. 2014. Human health effects of tetrachloroethylene: Key findings and scientific issues. *Environ Health Perspect* 122(4):325-334. <http://doi.org/10.1289/ehp.1307359>.
- Guzelian PS, Henry CJ, Olin SS. 1992. Similarities and differences between children and adults: Implications for risk assessment. Washington, DC: International Life Sciences and Press Institute Press.
- *Haddad LM, Winchester JF. 1990. Elimination of poison from the eyes, skin and gastrointestinal tract. In: *Clinical management of poisoning and drug overdose*. Second edition. Philadelphia, PA: W.B. Saunders Company, Harcourt Brace Jovanovich, Inc., 13-20.
- Hadkhale K, Martinsen JI, Weiderpass E, et al. 2017. Occupational exposure to solvents and bladder cancer: A population-based case control study in Nordic countries. *Int J Cancer* 140(8):1736-1746. <http://doi.org/10.1002/ijc.30593>.
- Haerer AF, Udelman HD. 1964. Acute brain syndrome secondary to tetrachloroethylene ingestion. *Am J Psychiatry* 12:78-79.
- Hake CL, Stewart RD. 1977. Human exposure to tetrachloroethylene: Inhalation and skin contact. *Environ Health Perspect* 21:231-238.
- Hamada T, Tanaka H. 1995. Transfer of methyl chloroform, trichloroethylene and tetrachloroethylene to milk, tissues and expired air following intraruminal or oral administration in lactating goats and milk-fed kids. *Environ Pollut* 87(3):313-318.
- Hanioka N, Jinno H, Toyo'oka T, et al. 1995. Induction of rat liver drug-metabolizing enzymes by tetrachloroethylene. *Arch Environ Contam Toxicol* 28:273-280.

9. REFERENCES

- Hardin BD, Bond GP, Sikov MR, et al. 1981. Testing of selected workplace chemicals for teratogenic potential. *Scand J Work Environ Health* 7(Suppl 4):66-75.
- Harkov R, Kebbekus B, Bozzelli JW, et al. 1984. Comparison of selected volatile organic compounds during the summer and winter at urban sites in New Jersey. *Sci Total Environ* 38:259-274.
- Hartmann A, Speit G. 1995. Genotoxic effects of chemicals in the single cell gel (SCG) test with human blood cells in relation to the induction of sister-chromatid exchanges (SCE). *Mutat Res* 346(1):49-56.
- Hartwell TD, Pellizzari ED, Perritt RL, et al. 1987. Comparison of volatile organic levels between sites and seasons for the total exposure assessment methodology (TEAM) study. *Atmos Environ* 21:2413-2424.
- Hasanen E, Soininen V, Pyysalo M, et al. 1979. The occurrence of aliphatic chlorine and bromine compound in automobile exhaust. *Atmos Environ* 13:1217-1219.
- Hattis D, White P, Koch P. 1993. Uncertainties in pharmacokinetic modeling for perchloroethylene: II. Comparison of model predictions with data for a variety of different parameters. *Risk Anal* 13(6):599-610.
- Hattis D, White P, Marmorstein L, et al. 1990. Uncertainties in pharmacokinetic modeling for perchloroethylene. I. Comparison of model structure, parameters, and predictions for low-dose metabolism rates for models derived by different authors. *Risk Anal* 10(3):449-458.
- Haworth S, Lawlor T, Mortelmans K, et al. 1983. Salmonella mutagenicity test results for 250 chemicals. *Environ Mutagen Supplement* 1:3-142.
- Hayes JR, Condie LW Jr, Borzelleca JF. 1986. The subchronic toxicity of tetrachloroethylene (perchloroethylene) administered in the drinking water of rats. *Fundam Appl Toxicol* 7:119-125.
- HazDat. 2007. Trichloroethylene. HazDat Database: ATSDR's Hazardous Substance Release and Health Effects Database. Atlanta, GA: Agency for Toxic Substances and Disease Registry.
- *Health Canada. 2014. Tetrachloroethylene in drinking water. Document for public comment. http://www.hc-sc.gc.ca/ewh-semt/alt_formats/pdf/consult/_2014/tetrachloroethylene/consult-eng.pdf. May 10, 2016.
- Heikes DL, Hopper ML. 1986. Purge and trap method for determination of fumigants in whole grains, milled grain products, and intermediate grain-based foods. *J Assoc Off Anal Chem* 69:990-998.
- *Heineman EF, Cocco P, Gomez MR, et al. 1994. Occupational exposure to chlorinated aliphatic hydrocarbons and risk of astrocytic brain cancer. *Am J Ind Med* 26:155-169.
- Hemminki K, Saloniemi I, Luoma K, et al. 1980. Transplacental carcinogens and mutagens: Childhood cancer, malformation and abortions as risk indicators. *J Toxicol Environ Health* 6:1115-1126.
- Henschler D. 1977. Metabolism and mutagenicity of halogenated olefins: A comparison of structure and activity. *Environ Health Perspect* 21:61-64.

9. REFERENCES

- Hickman JC. 2000. Tetrachloroethylene. In: Kirk Othmer encyclopedia of chemical technology. John Wiley & Sons, Inc., 1-9.
- Hoel DG, Davis DL, Miller AB, et al. 1992. Trends in cancer mortality in 15 industrialized countries, 1969-1986. *J Natl Cancer Inst* 84(5):313-320.
- Hook EB. 1982. Perspectives in mutation epidemiology: 2. Epidemiologic and design aspects of studies of somatic chromosome breakage and sister-chromatic exchange. *Mutat Res* 99:373-382.
- Hoyt SD, Smith VL. 1991. Measurement of toxic organic compounds in ambient air using EPA method TO-14. In: *Capillary chromatography - The applications*. Jennings W, Nikelly JG, eds. Heidelberg: Huthig Buch Verlag, 83-94.
- HSDB. 2013. Tetrachloroethylene. Hazardous Substances Data Bank. National Library of Medicine, <http://toxnet.nlm.nih.gov>. May 09, 2013.
- Hunkeler D, Aravena R, Butler BJ. 1999. Monitoring microbial dechlorination of tetrachloroethene (PCE) in groundwater using compound-specific stable carbon isotope ratios: Microcosm and field studies. *Environ Sci Technol* 33(16):2733-2738.
- IARC. 1979. IARC monographs on the evaluation of the carcinogenic risk of chemicals in humans. Tetrachloroethylene. Lyon, France: World Health Organization, 20:491-514.
- IARC. 2014. Tetrachloroethylene. In: Trichloroethylene, tetrachloroethylene, and some other chlorinated agents. Vol. 106. IARC Monographs on the evaluation of carcinogenic risks to humans. World Health Organization, International Agency for Research on Cancer, 219-351. <https://monographs.iarc.fr/ENG/Monographs/vol106/mono106-F01-F02.pdf>. April 6, 2016.
- ICRP. 2002. Basic anatomical and physiological data for use in radiological protection: Reference values. A report of age-and gender-related differences in the anatomical and physiological characteristics of reference individuals. International Commission on Radiological Protection. *Ann ICRP* 32(Publication 89):5-265.
- Ikeda M, Imamura T. 1973. Biological half-life of trichloroethylene and tetrachloroethylene in human subjects. *Int Arch Arbeitsmed* 31:209-224.
- Ikeda M, Koizuma A, Watanabe T, et al. 1980. Cytogenetic and cytokinetic investigations on lymphocytes from workers occupationally exposed to tetrachloroethylene. *Toxicol Lett* 5:251-256.
- Ikeda M, Ohtsuji H, Imamura T, et al. 1972. Urinary excretion of total trichloro-compounds, trichloroethanol, and trichloroacetic acid as a measure of exposure to trichloroethylene and tetrachloroethylene. *Br J Ind Med* 29:328-333.
- Imbriani M, Ghittori S, Pezzagno G, et al. 1988. Urinary excretion of tetrachloroethylene (perchloroethylene) in experimental and occupational exposure. *Arch Environ Health* 43:292-298.
- IRIS. 2012. Tetrachloroethylene. Integrated Risk Information System. Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/iris/>. April 24, 2013.
- Isalou M, Sleep BE, Liss SN. 1998. Biodegradation of high concentrations of tetrachloroethene in a continuous flow column system. *Environ Sci Technol* 32(22):3579-3585.

9. REFERENCES

- Isenschmid DS, Cassin BJ, Hepler BR, et al. 1998. Tetrachloroethylene intoxication in an autoerotic fatality. *J Forensic Sci* 43(1):231-234.
- Itoh N, Kutsuna S, Ibusuki T. 1994. A product study of the OH radical initiated oxidation of perchloroethylene and trichloroethylene. *Chemosphere* 28(11):2029-2040.
- ITRC. 2003. An introduction to characterizing sites contaminated with DNAPLs. Interstate Technology and Regulatory Council, Dense Nonaqueous Phase Liquids Team.
- Jakobson I, Wahlberg JE, Holmberg B, et al. 1982. Uptake via the blood and elimination of 10 organic solvents following epicutaneous exposure of anesthetized guinea pigs. *Toxicol Appl Pharmacol* 63:181-187.
- James KJ, Stack MA. 1996. The determination of volatile organic compounds in soils using solid phase microextraction with gas chromatography-mass spectrometry. *J High Resolut Chromatogr* 19(9):515-519.
- Jang JY, Droz PO. 1997. Ethnic differences in biological monitoring of several organic solvents. II. A simulation study with a physiologically based pharmacokinetic model. *Int Arch Occup Environ Health* 70(1):41-50.
- Jang JY, Kang S K, Chung HK. 1993. Biological exposure indices of organic solvents for Korean workers. *Int Arch Occup Environ Health* 65:S219-S222.
- Janulewicz PA, White RF, Martin BM, et al. 2012. Adult neuropsychological performance following prenatal and early postnatal exposure to tetrachloroethylene (PCE)-contaminated drinking water. *Neurotoxicol Teratol* 34(3):350-359.
- Janulewicz PA, Killiany RJ, White RF, et al. 2013. Structural magnetic resonance imaging in an adult cohort following prenatal and early postnatal exposure to tetrachloroethylene (PCE)-contaminated drinking water. *Neurotoxicol Teratol* 38:13-20. 10.1016/j.ntt.2013.03.060.
- Janulewicz PA, White RF, Winter MR, et al. 2008. Risk of learning and behavioral disorders following prenatal and early postnatal exposure to tetrachloroethylene (PCE)-contaminated drinking water. *Neurotoxicol Teratol* 30(3):175-185.
- Jeffers PM, Ward LM, Woytowitch LM, et al. 1989. Homogenous hydrolysis rate constants for selected chlorinated methanes, ethanes, ethenes and propanes. *Environ Sci Technol* 23:965-969.
- *Ji J, Hemminki K. 2006. Socioeconomic/occupational risk factors for lymphoproliferative diseases in Sweden. *Ann Epidemiol* 16(5):370-376. 10.1016/j.annepidem.2005.09.002.
- JISA. 1993. Carcinogenicity study of tetrachloroethylene by inhalation in rats and mice. Japan Industrial Safety Association.
- Johnson PC, Ettinger RA. 1991. Heuristic model for predicting the intrusion rate of contaminant vapors into buildings. *Environ Sci Technol* 25(8):1445-1452.

9. REFERENCES

- Johnston JE, Gibson JM. 2013. Spatiotemporal variability of tetrachloroethylene in residential indoor air due to vapor intrusion: A longitudinal, community-based study. *J Expo Sci Environ Epidemiol* [epub ahead of print].
- Johnston JE, Gibson JM. 2014. Spatiotemporal variability of tetrachloroethylene in residential indoor air due to vapor intrusion: a longitudinal, community-based study. *J Expo Sci Environ Epidemiol* 24(6):564-571. 10.1038/jes.2013.13.
- Jonker D, Woutersen RA, Feron VJ. 1996. Toxicity of mixtures of nephrotoxicants with similar or dissimilar mode of action. *Food Chem Toxicol* 34(11-12):1075-1082.
- Jung WT, Fujita M, Sohn DH. 1992. Levels of volatile halogenated hydrocarbons in Tokyo rain and their seasonal, time-series changes. *Jpn J Toxicol Environ Health* 38:490-497.
- Karlsson JE, Rosengren LE, Kjellstrand P, et al. 1987. Effects of low-dose inhalation of three chlorinated aliphatic organic solvents on deoxyribonucleic acid in gerbil brain. *Scand J Work Environ Health* 13:453-458.
- Katz RM, Jowett D. 1981. Female laundry and dry cleaning workers in Wisconsin. A mortality analysis. *Am J Public Health* 71:305-307.
- Kawakami T, Takano T, Araki R. 1988. Synergistic interaction of tri- and tetra-chloroethylene, hypoxia, and ethanol on the atrioventricular conduction of the perfused rat heart. *Ind Health* 26:25-33.
- Kawasaki M. 1980. Experiences with the test scheme under the chemical control law of Japan: An approach to structure-activity correlations. *Ecotoxicol Environ Saf* 4:444-454.
- Kawata K, Fujieda Y. 1993. Volatile chlorinated hydrocarbons in ambient air at Niigata area. *Jpn J Toxicol Environ Health* 39(5):474-479.
- Kawata K, Tanabe A, Saito S, et al. 1997. Screening of volatile organic compounds in river sediment. *Bull Environ Contam Toxicol* 58(6):893-900.
- Kearns GL, Abdel-Rahman SM, Alander SW, et al. 2003. Developmental pharmacology--drug disposition, action, and therapy in infants and children. *N Engl J Med* 349(12):1157-1167. 10.1056/NEJMra035092.
- Keil SL. 1985. Tetrachloroethylene. In: Grayson M, Eckroth P, eds. *Kirk-Othmer concise encyclopedia of chemical technology*. New York, NY: John Wiley and Sons, 754-762.
- Kenaga EE. 1980. Predicted bioconcentration factors and soil sorption coefficients of pesticides and other chemicals. *Ecotoxicol Environ Saf* 4:26-38.
- Kendrick JF. 1929. The treatment of hookworm disease with tetrachloroethylene. *Am J Trop Med* 483-488.
- Kezic S, Monster AC, Kruse J, et al. 2000. Skin absorption of some vaporous solvents in volunteers. *Int Arch Occup Environ Health* 73(6):415-422.
- Kezic S, Monster AC, van de Gevel IA, et al. 2001. Dermal absorption of neat liquid solvents on brief exposures in volunteers. *Am Ind Hyg Assoc J* 62(1):12-18.

9. REFERENCES

- *Kido K, Shiratori T, Watanabe T, et al. 1989. Correlation of tetrachloroethylene in blood and in drinking water: A case of well water pollution. *Bull Environ Contam Toxicol* 43:444-453.
- Kido T, Sugaya C, Ikeuchi R, et al. 2013. The increases in mRNA expressions of inflammatory cytokines by adding cleaning solvent or tetrachloroethylene in the murine macrophage cell line J774.1 evaluated by real-time PCR. *Ind Health* 51:319-323.
- Kim BH, Baek KH, Cho DH, et al. 2010. Complete reductive dechlorination of tetrachloroethene to ethene by anaerobic microbial enrichment culture developed from sediment. *Biotechnol Lett* 32(12):1829-1835.
- Kinkead ER, Leahy HF. 1987. Evaluation of the acute toxicity of selected groundwater contaminants. Government Reports Announcements & Index (GRA&I). ADA1801984.
- Kirchner K, Helf D, Ott P, et al. 1990. The reaction of OH radicals with 1,1-di, tri- and tetrachloroethylene. *Berichte der Bunsengesellschaft für Physikalische Chemie* 94:77-83.
- Kjellstrand P, Bjerkemo M, Adler-Maihofer M, et al. 1985. Effects of solvent exposure on testosterone levels and butyrylcholinesterase activity in mice. *Acta Pharmacol Toxicol* 57:242-249.
- Kjellstrand P, Holmquist B, Kanje M, et al. 1984. Perchloroethylene: Effects on body and organ weights and plasma butyrylcholinesterase activity in mice. *Acta Pharmacol Toxicol* 54:414-424.
- Klaassen CD, Plaa GL. 1966. Relative effects of various chlorinated hydrocarbons on liver and kidney function in mice. *Toxicol Appl Pharmacol* 9:139-151.
- Klaassen CD, Plaa GL. 1967. Relative effects of various chlorinated hydrocarbons on liver and kidney function in dogs. *Toxicol Appl Pharmacol* 10:119-131.
- Kline SA, McCoy EC, Rosenkranz HS, et al. 1982. Mutagenicity of chloroalkene epoxides in bacterial systems. *Mutat Res* 101:115-125.
- Kluwe WM, Abdo KM, Huff J. 1984. Chronic kidney disease and organic chemical exposure: Evaluations of causal relationships in humans and experimental animals. *Fundam Appl Toxicol* 4:899-901.
- Kobayashi S, Hutcheon DE, Regan J. 1982. Cardiopulmonary toxicity of tetrachloroethylene. *J Toxicol Environ Health* 10:23-30.
- Koch R, Schlegelmilch R, Wolf HU. 1988. Genetic effects of chlorinated ethylenes in the yeast *Saccharomyces cerevisiae*. *Mutat Res* 206:209-216.
- Koizumi A. 1989. Potential of physiologically based pharmacokinetics to amalgamate kinetic data of trichloroethylene and tetrachloroethylene obtained in rats and man. *Br J Ind Med* 46(4):239-249.
- Komori M, Nishio K, Kitada M, et al. 1990. Fetus-specific expression of a form of cytochrome P-450 in human livers. *Biochemistry* 29(18):4430-4433.
- Koppel C, Arndt I, Arendt U, et al. 1985. Acute tetrachloroethylene poisoning: Blood elimination kinetics during hyperventilation therapy. *Clin Toxicol* 23:103-115.

9. REFERENCES

- Korpela M. 1988. Organic solvents affect rat synaptosome membrane acetylcholinesterase and adenosine triphosphatase *in vitro*. *Arch Toxicol Suppl* 12:384-386.
- Korpela M, Tahti H. 1986. The effect of selected organic solvents on intact human red cell membrane acetylcholinesterase *in vitro*. *Toxicol Appl Pharmacol* 85:257-262.
- Krause RJ, Lash LH, Elfarra AA. 2003. Human kidney flavin-containing monooxygenases and their potential roles in cystein S-conjugate metabolism and nephrotoxicity. *J Pharmacol Exp Ther* 304(1):185-191.
- Kringstad KP, Ljungquist PO, deSousa F, et al. 1981. Identification and mutagenic properties of some chlorinated aliphatic compounds in the spent liquor from Kraft pulp chlorination. *Environ Sci Technol* 15:562-566.
- Krishnan K, Andersen ME. 1994. Physiologically based pharmacokinetic modeling in toxicology. In: Hayes AW, ed. *Principles and methods of toxicology*. 3rd ed. New York, NY: Raven Press, Ltd., 149-188.
- Krishnan K, Anderson ME, Clewell HJ, et al. 1994. Physiologically based pharmacokinetic modeling of chemical mixtures. In: Yang RSH, ed. *Toxicology of chemical mixtures. Case studies, mechanisms, and novel approaches*. San Diego, CA: Academic Press, 399-437.
- Krost KJ, Pellizzari ED, Walburn SG, et al. 1982. Collection and analysis of hazardous organic emissions. *Anal Chem* 54:810-817.
- Krotoszynski BK, Bruneau GM, O'Neill HJ. 1979. Measurement of chemical inhalation exposure in urban population in the presence of endogenous effluents. *J Anal Toxicol* 3:225-234.
- Krska R, Taga K, Kellner R. 1993. New IR fiber-optic chemical sensor for in situ measurements of chlorinated hydrocarbons in water. *Appl Spectroscopy* 47(9):1484-1487.
- Krumholz LR, Sharp R, Fishbain SS. 1996. A freshwater anaerobe coupling acetate oxidation to tetrachloroethylene dehalogenation. *Appl Environ Microbiol* 62(11):4108-4113.
- Kukongviriyapan V, Kukongviriyapan U, Stacey NH. 1990. Interference with hepatocellular substrate uptake by 1,1,1-trichloroethane and tetrachloroethylene. *Toxicol Appl Pharmacol* 102(1):80-90.
- Kylin B, Reichard H, Sumegi I, et al. 1963. Hepatotoxicity of inhaled trichloroethylene, tetrachloroethylene, and chloroform--single exposure. *Acta Pharmacol Toxicol* 20:16-26.
- Kylin B, Sumegi I, Yllner S. 1965. Hepatotoxicity of inhaled trichloroethylene and tetrachloroethylene--long-term exposure. *Acta Pharmacol Toxicol* 22:379-385.
- Kyrklund T, Haglid K. 1991. Brain lipid composition in guinea pigs after intrauterine exposure to perchloroethylene. *Pharmacol Toxicol* 68:146-148.
- Kyrklund T, Alling C, Kjellstrand P, et al. 1984. Chronic effects of perchloroethylene on the composition of lipid and acyl groups in cerebral cortex and hippocampus of the gerbil. *Toxicol Lett* 22:343-349.

9. REFERENCES

- Kyrklund T, Kjellstrand P, Haglid K. 1987. Brain lipid changes in rats exposed to xylene and toluene. *Toxicology* 45:123-133.
- Kyrklund T, Kjellstrand P, Haglid KG. 1988. Effects of exposure to Freon 11, 1,1,1-trichloroethane or perchloroethylene on the lipid and fatty-acid composition of rat cerebral cortex. *Scand J Work Environ Health* 14:91-94.
- Kyrklund T, Kjellstrand P, Haglid KG. 1990. Long-term exposure of rats to perchloroethylene, with and without a post-exposure solvent-free recovery period: Effects on brain lipids. *Toxicol Lett* 52:279-285.
- Kyyronen P, Taskinen H, Lindbohm ML, et al. 1989. Spontaneous abortions and congenital malformations among women exposed to tetrachloroethylene in dry cleaning. *J Epidemiol Community Health* 43:346-351.
- Lafuente A, Mallol J. 1986. Thioethers in urine during occupational exposure to tetrachloroethylene. *Br J Ind Med* 43:68-69.
- Lagakos SW, Wessen BJ, Zelen M, et al. 1986. An analysis of contaminated well water and health effects in Woburn, Massachusetts. *J Am Stat Assoc* 81:583-614.
- Larson CD, Love OT, Jr., Reynolds G. 1983. Tetrachloroethylene leached from lined asbestos-cement pipe into drinking water. *J Am Water Works Assoc* April 1983:184-190.
- Lash LH, Parker JC. 2001. Hepatic and renal toxicities associated with perchloroethylene. *Pharmacol Rev* 53(2):177-208.
- Lash LH, Putt DA, Huang P, et al. 2007. Modulation of hepatic and renal metabolism and toxicity of trichloroethylene and perchloroethylene by alterations in status of cytochrome P450 and glutathione. *Toxicology* 235(1-2):11-26.
- Lash LH, Qian W, Putt DA, et al. 1998. Glutathione conjugation of perchloroethylene in rats and mice *in vitro*: Sex-, species-, and tissue-dependent differences. *Toxicol Appl Pharmacol* 150(1):49-57.
- Lash LH, Qian W, Putt DA, et al. 2002. Renal toxicity of perchloroethylene and S-(1,2,2-trichlorovinyl)glutathione in rats and mice: Sex- and species-dependent differences. *Toxicol Appl Pharmacol* 179(3):163-171.
- Lash LH, Qian W, Xu Y, et al. 1996. Renal and hepatic metabolism and cytotoxicity of perchloroethylene (PER) by GSH conjugation. *Toxicologist* 30(1):314.
- Lauwerys R, Herbrand J, Buchet JP, et al. 1983. Health surveillance of workers exposed to tetrachloroethylene in dry cleaning shops. *Int Arch Occup Environ Health* 52:69-77.
- Lee LJ, Chan C, Chung C. 2002. Health risk assessment on residents exposed to chlorinated hydrocarbons contaminated in groundwater of a hazardous waste site. *J Toxicol Environ Health Part A* 65:219-235.
- Leeder JS, Kearns GL. 1997. Pharmacogenetics in pediatrics: Implications for practice. *Pediatr Clin North Am* 44(1):55-77.

9. REFERENCES

- Lehmann I, Rehwagen M, Dietz U, et al. 2001. Enhanced *in vivo* IgE production and T cell polarization toward the type 2 phenotype in association with indoor exposure to VOC: Results of the LARS study. *Int J Hyg Environ Health* 204:211-221.
- Lehmann I, Thoeleke A, Rehwagen M, et al. 2002. The influence of maternal exposure to volatile organic compounds on the cytokine secretion profile of neonatal T cells. *Environ Toxicol* 17(3):203-210.
- Leibman KC, Ortiz E. 1977. Metabolism of halogenated ethylenes. *Environ Health Perspect* 21:91-97.
- *Leikin JB, Paloucek FP. 2008. Tetrachloroethylene. In: *Poisoning and toxicology handbook*, 4th ed. Hudson, OH: Lexi-Comp, Inc., 853.
- Leung H. 1993. Physiologically-based pharmacokinetic modelling. In: Ballantyne B, Marrs T, Turner P, eds. *General and applied toxicology*. Vol. 1. New York, NY: Stockton Press, 153-164.
- Levine B, Fierro MF, Goza SW, et al. 1981. A tetrachloroethylene fatality. *J Forensic Sci* 26:206-209.
- LHWMP. 2013. Evaluating vapor leak detectors for use in "PERC" dry cleaners. Final report. King County, WA: Local Hazardous Waste Management Program.
- Lide DR, ed. 1990. *CRC Handbook of chemistry and physics*. 71st ed. Ann Arbor, MI: CRC Press, Inc., 3-242.
- Lide DR. 2008. Tetrachloroethylene. In: *CRC handbook of chemistry and physics*. 88th ed. Boca Raton, FL: CRC Press, 3-470.
- Ligocki MP, Leuenberger C, Pankow JF. 1985. Trace organic compounds in rain-II. Gas scavenging of neutral organic compounds. *Atmos Environ* 19:1609-1617.
- Lillford L, Beevers C, Bowen D, et al. 2010. RE: DNA damage detected by the alkaline comet assay in the liver of mice after oral administration of tetrachloroethylene. *Mutagenesis* 25:133-138.
- Lin YS, Egeghy PP, Rappaport SM. 2008. Relationships between levels of volatile organic compounds in air and blood from the general population. *J Expo Sci Environ Epidemiol* 18(4):421-429. 10.1038/sj.jes.7500635.
- Lindbohm ML, Taskinen H, Sallmén M, et al. 1990. Spontaneous abortions among women exposed to organic solvents. *Am J Ind Med* 17:449-463.
- Ling S, Lindsay WA. 1971. Perchloroethylene burns (Letter). *Br Med J* 3:115.
- Lipworth L, Sonderman JS, Mumma MT, et al. 2011. Cancer mortality among aircraft manufacturing workers: An extended follow-up. *J Occup Environ Med* 53(9):992-1007.
- Livingston AL. 1978. Forage plant estrogens. *J Toxicol Environ Health* 4(2-3):301-324.
- Loizou GD. 2001. The application of physiologically based pharmacokinetic modelling in the analysis of occupational exposure to perchloroethylene. *Toxicol Lett* 124(1-3):59-69.
- *Lorenz H, Weber E, Omlor A, et al. 1990. [Evidence of cerebral lesions caused by tetrachloroethylene]. *Zbl Arbeitsmed* 40:355-364. (German)

9. REFERENCES

- Lovell D. 2010. Is tetrachloroethylene genotoxic or not? *Mutagenesis* 25(5):443-446.
- Lu C, Bjerg PL, Zhang F, et al. 2011. Sorption of chlorinated solvents and degradation products on natural clayey tills. *Chemosphere* 83(11):1467-1474.
- Lucas D, Herve A, Lucas R, et al. 2015. Assessment of exposure to perchloroethylene and its clinical repercussions for 50 dry-cleaning employees. *J Occup Environ Hyg* 12(11):767-773. 10.1080/15459624.2015.1048346.
- Lukaszewski T. 1979. Acute tetrachloroethylene fatality. *Clin Toxicol* 15:411-415.
- Lurker PA, Clark CS, Elia VJ. 1982. Atmospheric release of chlorinated organic compounds from the activated sludge process. *J Water Pollut Control Fed* 54:1566-1573.
- *Lynge E. 1994. Danish cancer registry as a resource for occupational research. *J Occup Med* 36(11):1169-1173.
- Lynge E, Thygesen L. 1990. Primary liver cancer among women in laundry and dry cleaning work in Denmark. *Scand J Work Environ Health* 16:108-112.
- Lynge E, Andersen A, Rylander L, et al. 2006. Cancer in persons working in dry cleaning in the Nordic countries. *Environ Health Perspect* 114(2):213-219.
- Lynge E, Carstensen B, Andersen O. 1995. Primary liver cancer and renal cell carcinoma in laundry and dry cleaning workers in Denmark. *Scand J Work Environ Health* 21:293-295.
- Ma T-H, Sandhu SS, Peng Y, et al. 1992. Synergistic and antagonistic effects on genotoxicity of chemicals commonly found in hazardous waste sites. *Mutat Res* 270:71-77.
- Macca I, Carrieri M, Scapellato ML, et al. 2012. Biological monitoring of exposure to perchloroethylene in dry cleaning workers. *Med Lav* 103(5):382-393.
- MacMahon B. 1986. Comment. *J Am Stat Assoc* 81(395):597-599.
- Mahalingaiah S, Winter MR, Aschengrau A. 2016. Association of prenatal and early life exposure to tetrachloroethylene (PCE) with polycystic ovary syndrome and other reproductive disorders in the Cape Cod health study: A retrospective cohort study. *Reprod Toxicol* 65:87-94. <http://doi.org/10.1016/j.reprotox.2016.07.005>.
- Mahle DA, Gearhart JM, Godfrey RJ, et al. 2004. Determination of partition coefficients for a mixture of volatile organic compounds in rats and humans at different life stages. Wright-Patterson Air Force Base, OH: Air Force Research Laboratory, Applied Biotechnology Branch. ADA436299.
- Mahle DA, Gearhart JM, Grigsby CC, et al. 2007. Age-dependent partition coefficients for a mixture of volatile organic solvents in Sprague-Dawley rats and humans. *J Toxicol Environ Health A* 70(20):1745-1751.
- Makide Y, Tominaga T, Rowland FS. 1979. Gas chromatographic analysis of halogenated hydrocarbons in air over Japan. *Chem Lett* 355-358.

9. REFERENCES

- Maloney EK, Waxman DJ. 1999. Trans-activation of PPAR α and PPAR γ by structurally diverse environmental chemicals. *Toxicol Appl Pharmacol* 161(2):209-218.
- Marco-Urrea E, Gabarrell X, Sarrà M, et al. 2006. Novel aerobic perchloroethylene degradation by the white-rot fungus *Trametes versicolor*. *Environ Sci Technol* 40(24):7796-7802.
- Marth E. 1987. Metabolic changes following oral exposure to tetrachloroethylene in subtoxic concentrations. *Arch Toxicol* 60:293-299.
- Matsushima T, Nagano K, Nishizawa T, et al. 1998. Long term inhalation toxicity studies of five chlorinated hydrocarbons in F344 rats and BDF1 mice. Proceedings of the 1st International Conference of Asian Society of Toxicology, Yokohama, Japan, June 29-July 2, 1997. *J Toxicol Sci* 23(Suppl 2):296.
- Mattei F, Guida F, Matrat M, et al. 2014. Exposure to chlorinated solvents and lung cancer: Results of the ICARE study. *Occup Environ Med* 71(10):681-689. 10.1136/oemed-2014-102182.
- Mattie DR, Bates GD, Jepson GW, et al. 1994. Determination of skin: Air partition coefficients for volatile chemicals: Experimental methods and applications. *Fundam Appl Toxicol* 22:51-57.
- Mattsson JL, Albee RR, Yano BL, et al. 1992. Neurotoxicologic examination of rats exposed to 1,1,2,2-tetrachloroethylene vapor for 13 weeks. The Dow Chemical Company, sponsored by Halogenated Solvents Industry Alliance.
- Mattsson JL, Albee RR, Yano BL, et al. 1998. Neurotoxicologic examination of rats exposed to 1,1,2,2-tetrachloroethylene (perchloroethylene) vapor for 13 weeks. *Neurotoxicol Teratol* 20(1):83-98.
- Matushima T, Hayashi M, Matsuoka A, et al. 1999. Validation study of the *in vitro* micronucleus test in a Chinese hamster lung cell line (CHL/IU). *Mutagenesis* 14(6):569-580.
- Mayr U, Butsch A, Schneider S. 1992. Validation of two *in vitro* test systems for estrogenic activities with zearalenone, phytoestrogens and cereal extracts. *Toxicology* 74(2-3):135-149.
- Mazzullo M, Grilli S, Lattanzi G, et al. 1987. Evidence of DNA binding activity of perchloroethylene. *Res Commun Chem Pathol Pharmacol* 58:215-235.
- McCarthy MC, Hafner HR, Chinkin LR, et al. 2007. Temporal variability of selected air toxics in the United States. *Atmos Environ* 41(34):7180-7194. <http://dx.doi.org/10.1016/j.atmosenv.2007.05.037>. <http://www.sciencedirect.com/science/article/pii/S1352231007004840>.
- McDonald AD, Armstrong B, Cherry NM, et al. 1986. Spontaneous abortion and occupation. *J Occup Med* 28:1232-1238.
- McDonald AD, McDonald JC, Armstrong B, et al. 1987. Occupation and pregnancy outcome. *Br J Ind Med* 44:521-526.
- McDonald GJ, Wertz WE. 2007. PCE, TCE, and TCA vapors in subsurface soil, gas, and indoor air: A case study in upstate New York. *Ground Water Monit Remed* 27(4):86-92.
- McDougal JN, Jepson GW, Clewell HJ III, et al. 1990. Dermal absorption of organic chemical vapors in rats and humans. *Fundam Appl Toxicol* 14:299-308.

9. REFERENCES

- McKernan LT, Ruder AM, Petersen MR, et al. 2008. Biological exposure assessment to tetrachloroethylene for workers in the dry cleaning industry. *Environ Health* 7:12.
- Meckler LC, Phelps DK. 1966. Liver disease secondary to tetrachloroethylene exposure: A case report. *J Am Med Assoc* 197:662-663.
- Mennear J, Maronpot R, Boorman G, et al. 1986. Toxicologic and carcinogenic effects of inhaled tetrachloroethylene in rats and mice. *Dev Toxicol Environ Sci* 12:201-210.
- Meyer BR, Fischbein A, Rosenman K, et al. 1984. Increased urinary enzyme excretion in workers exposed to nephrotoxic chemicals. *Am J Med* 76:989-998.
- Michael LC, Erickson MD, Parks SP, et al. 1980. Volatile environmental pollutants in biological matrices with a headspace purge technique. *Anal Chem* 52:1836-1841.
- Milde G, Nerger M, Mergler R. 1988. Biological degradation of volatile chlorinated hydrocarbons in groundwater. Meeting on Groundwater Microbiology: Problems and Biological Treatment held at the International Association on Water Pollution Research and Control Symposium, Kuopio, Finland, August 4-6, 1987. *Water Sci Technol* 20:67-73.
- *Miligi L, Costantini AS, Benvenuti A, et al. 2006. Occupational exposure to solvents and the risk of lymphomas. *Epidemiology* 17(5):552-561. 10.1097/01.ede.0000231279.30988.4d.
- Monster AC. 1986. Biological monitoring of chlorinated hydrocarbon solvents. *J Occup Med* 28:583-588.
- Monster AC. 1988. Biological markers of solvent exposure. *Arch Environ Health* 43:90-93.
- Monster AC, Smolders JFJ. 1984. Tetrachloroethylene in exhaled air of persons living near pollution sources. *Int Arch Occup Environ Health* 53:331-336.
- Monster AC, Boersma G, Steenweg H. 1979. Kinetics of tetrachloroethylene in volunteers: Influence of exposure concentration and work load. *Int Arch Occup Environ Health* 42:303-309.
- Monster AC, Regouin-Peeters W, Van Schijndel A, et al. 1983. Biological monitoring of occupational exposure to tetrachloroethene. *Scand J Work Environ Health* 9:273-281.
- Morgan B. 1969. Dangers of perchloroethylene (Letter). *Br Med J* 2:513.
- Morselli PL, Franco-Morselli R, Bossi L. 1980. Clinical pharmacokinetics in newborns and infants: Age-related differences and therapeutic implications. *Clin Pharmacokinet* 5(6):485-527.
- Morton LM, Slager SL, Cerhan JR, et al. 2014. Etiologic heterogeneity among non-Hodgkin lymphoma subtypes: The interlymph non-Hodgkin lymphoma subtypes project. *Journal of the National Cancer Institute Monographs* 48:130-144.
- Moser VC, Cheek BM, MacPhail RC. 1995. A multidisciplinary approach to toxicological screening: III. Neurobehavioral toxicity. *J Toxicol Environ Health* 45:173-210.
- Moslen MT, Reynolds ES, Szabo S. 1977. Enhancement of the metabolism and hepatotoxicity of trichloroethylene and perchloroethylene. *Biochem Pharmacol* 26:369-375.

9. REFERENCES

- *Motohashi Y, Miyazaki Y, Takano T. 1993. Assessment of behavioral effects of tetrachloroethylene using a set of time-series analyses. *Neurotoxicol Teratol* 15:3-10.
- Motwani JN, Popp SA, Johnson GM, et al. 1986. Field screening techniques developed under the Superfund program. Management of uncontrolled hazardous waste sites, Hazardous Materials Control Research Institute, Washington, DC, December 1-3, 1986, Silver Spring, MD, 105-109.
- Mumtaz M, Fisher J, Blount B, et al. 2012a. Application of physiologically based pharmacokinetic models in chemical risk assessment. *J Toxicol Article ID* 904603. 10.1155/2012/904603.
- Mumtaz MM, Ray M, Crowell SR, et al. 2012b. Translational research to develop a human PBPK models tool kit—volatile organic compounds (VOCs). *J Toxicol Environ Health Part A* 75(1):6-24.
- Murakami K, Horikawa K. 1995. The induction of micronuclei in mice hepatocytes and reticulocytes by tetrachloroethylene. *Chemosphere* 31(7):3733-3739.
- Murphy CD, Moore RM, White RL. 2000. An isotopic method for determining production of volatile organohalogenes by marine microalgae. *Limnol Oceanogr* 45(8):1868-1871.
- Mutti A, Franchini I. 1987. Toxicity of metabolites to dopaminergic systems and the behavioral effects of organic solvents. *Br J Ind Med* 44:721-723.
- Mutti A, Alinovi R, Bergamaschi E, et al. 1992. Nephropathies and exposure to perchloroethylene in dry-cleaners. *Lancet* 340(8813):189-193.
- Nagano K, Nishizawa T, Yamamoto S, et al. 1998. Inhalation carcinogenesis studies of six halogenated hydrocarbons in rats and mice. In: Chiyoiani K, Hosoda Y, Aizawa Y, eds. *Advances in the prevention of occupational respiratory diseases*. Elsevier Science, 741-746.
- Nakahama T, Fukuhara M, Inouye Y. 1997. Volatile halogenated hydrocarbons in ambient air and the metabolites in human urine in an urban area. *Jpn J Toxicol Environ Health* 43(5):280-284.
- Nakatsuka H, Watanabe T, Takeuchi Y, et al. 1992. Absence of blue-yellow color vision loss among workers exposed to toluene or tetrachloroethylene, mostly at levels below occupational exposure limits. *Int Arch Occup Environ Health* 64:113-117.
- Narotsky MG, Kavlock RJ. 1995. A multidisciplinary approach to toxicological screening: II. Development toxicity. *J Toxicol Environ Health* 45:145-171.
- NAS. 1980. Drinking water and health. Volume 3. Washington, DC: National Academy of Science, Safe Drinking Water Committee, National Academy Press.
- NAS/NRC. 1989. Report of the oversight committee. Biologic markers in reproductive toxicology. Washington, DC: 15-35.
- Nazaroff WW, Weschler CJ. 2004. Cleaning products and air fresheners: Exposure to primary and secondary air pollutants. *Atmos Environ* 38(18):2841-2865.

9. REFERENCES

- NCI. 1977. Bioassay of tetrachloroethylene for possible carcinogenicity. National Cancer Institute. U.S. Department of Health, Education, and Welfare, Public Health Service, National Institutes of Health, DHEW Publ (NIH) 77-813.
- Neely WB, Branson DR, Blau GE. 1974. Partition coefficients to measure bioconcentration potential of organic chemicals in fish. *Environ Sci Technol* 8:1113-1115.
- Nelson BK, Taylor BJ, Setzer JV, et al. 1980. Behavioral teratology of perchloroethylene in rats. *J Environ Pathol Toxicol* 3:233-250.
- Neta G, Stewart PA, Rajaraman P, et al. 2012. Occupational exposure to chlorinated solvents and risks of glioma and meningioma in adults. *Occup Environ Med* 69(11):793-801. 10.1136/oemed-2012-100742.
- Newcombe R. 2000. Perchloroethylene (PCE) in dry cleaning establishments. http://www.webpages.uidaho.edu/etox/resources/case_studies/pce_dry.pdf. July 02, 2013.
- NICNAS. 2001. Tetrachloroethylene. Priority existing chemical assessment report. No. 15. National Industrial Chemicals Notification and Assessment Scheme.
- NIOSH. 1980. Teratogenic-mutagenic risk of workplace contaminants: Trichloroethylene, perchloroethylene, and carbon disulfide. Report to National Institute for Occupational Safety and Health, Cincinnati, OH by Litton Bionetics, Inc., Kensington, MD. Publication no. 82-185075.
- NIOSH. 1994. NIOSH manual of analytical methods. Third edition. Second supplement. Cincinnati, OH: NIOSH, U.S. Department of Health and Human Services, National Institute for Occupational Safety and Health, NIOSH method no. 1003.
- NIOSH. 2013. Tetrachloroethylene. NIOSH pocket guide to chemical hazards. Atlanta, GA: National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention. <http://www.cdc.gov/niosh/npg/>. April 24, 2013.
- NOES. 1990. National Occupational Exposure Survey (1981-1983). Cincinnati, OH: U.S. Department of Health and Human Services, National Institute for Occupational Safety and Health.
- NRC. 1993. Pesticides in the diets of infants and children. Washington, DC. National Research Council.
- NRC. 2010. Review of the Environmental Protection Agency's draft IRIS assessment of tetrachloroethylene. Washington, DC: National Research Council of the National Academies. 68-74, 81-113. http://www.nap.edu/catalog.php?record_id=12863. October 14, 2014.
- NTP. 1986. Toxicology and carcinogenesis studies of tetrachloroethylene (perchloroethylene) (CAS No. 127-18-4) in F344/N rats and B6C3F₁ mice (inhalation studies). National Toxicology Program--technical report series no. 311. Research Triangle Park, NC: U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health. NIH publication no. 86-2567.
- NTP. 2016. Report on carcinogens. 14th edition. Research Triangle Park, NC: U.S. Department of Health and Human Services, Public Health Service, National Toxicology Program. <https://ntp.niehs.nih.gov/pubhealth/roc/index-1.html>. May 30, 2018.

9. REFERENCES

- NYSDH. 2006. Final. Guidance for evaluating soil vapor intrusion in the state of New York. New York State Department of Health.
http://www.health.ny.gov/environmental/investigations/soil_gas/svi_guidance/docs/svi_main.pdf. August 28, 2013.
- *Oddone E, Edefonti V, Scaburri A, et al. 2014. Female breast cancer and electrical manufacturing: Results of a nested case-control study. *J Occup Health* 56(5):369-378.
- Odum J, Green T, Foster JR, et al. 1988. The role of trichloroacetic acid and peroxisome proliferation in the differences in carcinogenicity of perchloroethylene in the mouse and rat. *Toxicol Appl Pharmacol* 92:103-112.
- Ogata M, Takasuka Y, Tomokuni K, et al. 1971. Excretion of organic chlorine compounds in the urine of persons exposed to vapours of trichloroethylene and tetrachloroethylene. *Br J Ind Med* 28:386-390.
- OHM/TADS. 1990. Tetrachloroethylene. Oil and Hazardous Materials/Technical Assistance Data System, Chemical Information System.
- Ohtsuki T, Sato K, Koizumi A, et al. 1983. Limited capacity of humans to metabolize tetrachloroethylene. *Int Arch Occup Environ Health* 51:381-390.
- *Ojajarvi A, Partanen T, Ahlbom A, et al. 2001. Risk of pancreatic cancer in workers exposed to chlorinated hydrocarbon solvents and related compounds: A meta-analysis. *Am J Epidemiol* 153(9):841-850.
- *Ojajarvi A, Partanen T, Ahlbom A, et al. 2007. Estimating the relative risk of pancreatic cancer associated with exposure agents in job title data in a hierarchical Bayesian meta-analysis. *Scand J Work Environ Health* 33(5):325-335.
- Okouchi S. 1986. Volatilization coefficient for stripping trichloroethylene, 1,1,1-trichloroethylene, and tetrachloroethylene from water. *Water Sci Technol* 181:137-138.
- *Olsen GW, Heam S, Cook RR, et al. 1989. Mortality experience of a cohort of Louisiana USA chemical workers. *J Occup Med* 31:32-34.
- Olsen J, Hemminki K, Ahlborg G, et al. 1990. Low birthweight, congenital malformations, and spontaneous abortions among dry cleaning workers in Scandinavia. *Scand J Work Environ Health* 16:163-168.
- Olson MJ, Johnson JT, Reidy CA. 1990. A comparison of male rat and human urinary proteins: Implications for human resistance to hyaline droplet nephropathy. *Toxicol Appl Pharmacol* 102:524-536.
- Onofrij M, Thomas A, Paci C, et al. 1999. Optic neuritis with residual tunnel vision in perchloroethylene toxicity. *Eur Neurol* 41(1):51-53.
- Opdam JJG, Smolders JFJ. 1986. Alveolar sampling and fast kinetics of tetrachloroethene in man. I. Alveolar sampling. *Br J Ind Med* 43:814-824.
- Oppelt ET. 1987. Incineration of hazardous waste. A critical review. *J Air Pollut Control Assoc* 37:558-586.

9. REFERENCES

- OSHA. 2005. Reducing worker exposure to perchloroethylene (PERC) in dry cleaning. U.S. Department of Labor, Occupational Safety and Health Administration.
- OSHA. 2013a. List of highly hazardous chemicals, toxics, and reactives. Occupational safety and health standards. Occupational Safety and Health Administration. Code of Federal Regulations 29 CFR 1910.119, Appendix A. <http://www.osha.gov/law-regs.html>. April 24, 2013.
- OSHA. 2013b. Toxic and hazardous substances. Occupational safety and health standards. Occupational Safety and Health Administration. Code of Federal Regulations 29 CFR 1910.1000, Table Z-2. <http://www.osha.gov/law-regs.html>. April 24, 2013.
- Oshiro WM, Krantz QT, Bushnell PJ. 2008. Characterization of the effects of inhaled perchloroethylene on sustained attention in rats performing a visual signal detection task. *Neurotoxicol Teratol* 30(3):167-174.
- Otson R, Williams DT. 1982. Headspace chromatographic determination of water pollutants. *Anal Chem* 54:942-946.
- Otson R, Williams DT, Bothwell PD. 1982. Volatile organic compounds in water at thirty Canadian potable water treatment facilities. *J Assoc Off Anal Chem* 65:1370-1374.
- Otto D, Hudnell K, Boyes W. 1988. Electrophysiological measures of visual and auditory function as indices of neurotoxicity. *Toxicology* 49:205-218.
- Owen GM, Brozek J. 1966. Influence of age, sex and nutrition on body composition during childhood and adolescence. In: Falkner F, ed. *Human development*. Philadelphia, PA: WB Saunders, 222-238.
- *Palmer RB, Phillips SD. 2007. Chlorinated hydrocarbons. In: Shannon MW, Borron SW, Burns MJ, eds. *Haddad and Winchester's clinical management of poisoning and drug overdose*. 4th ed. Philadelphia, PA: Saunders Elsevier, 1347-1361.
- Pankow JF, Rosen ME. 1988. Determination of volatile compounds in water by purging directly to a capillary column with whole column cryotrapping. *Environ Sci Technol* 22:398-405.
- Park KS, Sorenson DL, Sims JL. 1988. Volatilization of wastewater trace organics in slow rate land treatment systems. *Hazard Waste Hazard Mater* 5:219-229.
- Parsons F, Barrio-Lage G, Rice R. 1985. Biotransformation of chlorinated organic solvents in static microcosms. *Environ Toxicol Chem* 4:739-742.
- Parsons F, Wood PR, Demarco J. 1984. Transformation of tetrachloroethene and trichloroethene in microcosms and groundwater. *J Am Water Works Assoc* 76:56-59.
- Patel R, Janakiraman N, Johnson R, et al. 1973. Pulmonary edema and coma from perchloroethylene. *J Am Med Assoc* 223:1510.
- Paulu C, Aschengrau A, Ozonoff D. 1999. Tetrachloroethylene-contaminated drinking water in Massachusetts and the risk of colon-rectum, lung, and other cancers. *Environ Health Perspect* 107(4):265-271.

9. REFERENCES

- Pearson CR, McConnell G. 1975. Chlorinated Cl and C2 hydrocarbons in the marine environment. *Proc R Soc Lond [Biol]* 189:305-332.
- Peers AM. 1985. Method 5: Determination of trichloroethylene and tetrachloroethylene in air. *IARC Sci Publ* 68:205-211.
- Pegg DG, Zempel JA, Braun WH, et al. 1979. Disposition of (¹⁴C) tetrachloroethylene following oral and inhalation exposure in rats. *Toxicol Appl Pharmacol* 51:465-474.
- Pekari K, Aitio A. 1985a. Method 29: Determination of trichloroacetic acid in urine. *International Agency for Research on Cancer Science Publication* 68:467-472.
- Pekari K, Aitio A. 1985b. Method 30: Determination of 2,2,2-trichloroethanol in urine. *International Agency for Research on Cancer Science Publication* 68:473-478.
- Pellizzari ED, Hartwell TD, Harris BS, et al. 1982. Purgeable organic compounds in mother's milk. *Bull Environ Contam Toxicol* 28:322-328.
- Pennell KD, Jin M, Abriola LM, et al. 1994. Surfactant enhanced remediation of soil columns contaminated by residual tetrachloroethylene. *J Contam Hydrol* 16:35-53.
- Pennell KG, Bozkurt O, Suuberg EM. 2009. Development and application of a three-dimensional finite element vapor intrusion model. *J Air Waste Manage Assoc* 59(4):447-460.
- Pennell KG, Scammell MK, McClean MD, et al. 2013. Sewer gas: An indoor air source of PCE to consider during vapor intrusion investigations. *Ground Water Monit Remed* 33(3):119-126.
- Perrin MC, Opler MG, Harlap S, et al. 2007. Tetrachloroethylene exposure and risk of schizophrenia: Offspring of dry cleaners in a population birth cohort, preliminary findings. *Schizophr Res* 90(1-3):251-254.
- Pesch B, Haerting J, Ranft U, et al. 2000a. Occupational risk factors for renal cell carcinoma: Agent-specific results from a case-control study in Germany. *Int J Epidemiol* 29:1014-1024.
- Pesch B, Haerting J, Ranft U, et al. 2000b. Occupational risk factors for urothelial carcinoma: Agent-specific results from a case-control study in Germany. *Int J Epidemiol* 29:238-247. <http://doi.org/10.1093/ije/29.2.238>.
- *Pezzoli G, Cereda E. 2013. Exposure to pesticides or solvents and risk of Parkinson disease. *Neurology* 80(22):2035-2041. 10.1212/WNL.0b013e318294b3c8.
- Pezzagno G, Imbriani M, Ghittori S, et al. 1988. Urinary concentration, environmental concentration, and respiratory uptake of some solvents: Effect of the work load. *Am Ind Hyg Assoc J* 49:546-552.
- Philip BK, Mumtaz MM, Latendresse JR, et al. 2007. Impact of repeated exposure on toxicity of perchloroethylene in Swiss Webster mice. *Toxicology* 232(1-2):1-14.
- Piet GJ, Morra CHF, DeKruif HAM. 1981. The behavior of organic micropollutants during passage through the soil. *Quality of Groundwater, Proceedings of an International Symposium. Stud Environ Sci* 17:557-564.

9. REFERENCES

- Piper WN, Sparschu GL. 1969. A comparison of the inhalation toxicity of 1,1,1,2-tetrachloroethane and perchloroethylene in the rat. Dow Chemical. Submitted to the U.S. Environmental Protection Agency under TSCA, Section 8d. OTS0515971.
- Pocklington DW. 1992. Determination of tetrachloroethylene. Results of a collaborative study and the standardized method. *J Am Oil Chem Soc* 69(8):789-793.
- Poet TS, Weitz KK, Gies RA, et al. 2002. PBPK modeling of the percutaneous absorption of perchloroethylene from a soil matrix in rats and humans. *Toxicol Sci* 67(1):17-31.
- Polder MD, Hulzebos EM, Jager DT. 1998. Bioconcentration of gaseous organic chemicals in plant leaves: Comparison of experimental data with model predictions. *Environ Toxicol Chem* 17(5):962-968.
- Poli D, Manini P, Andreoli R, et al. 2005. Determination of dichloromethane, trichloroethylene and perchloroethylene in urine samples by headspace solid phase microextraction gas chromatography-mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci* 820(1):95-102.
- Popp W, Muller G, Baltes-Schmitz B, et al. 1992. Concentrations of tetrachloroethene in blood and trichloroacetic acid in urine of workers and neighbors of dry cleaning shops. *Int Arch Occup Environ Health* 63:393-395.
- Potter CL, Chang LW, Deangelo AB, et al. 1996. Effects of four trihalomethanes on DNA strand breaks, renal hyaline droplet formation and serum testosterone in male F-344 rats. *Cancer Lett* 106(2):235-242.
- Prentice, RL. 1986. Comment. *J Am Stat Assoc* 81(395):600-601.
- Price PJ, Hassett CM, Mansfield JI. 1978. Transforming activities of trichloroethylene -and proposed industrial alternatives. *In Vitro* 14:290-293.
- Price RG, Taylor SA, Crutcher E, et al. 1995. The assay of laminin fragments in serum and urine as an indicator of renal damage induced by toxins. *Toxicol Lett* 77:313-318.
- Pukkala E, Martinsen JI, Lynge E, et al. 2009. Occupation and cancer - follow-up of 15 million people in five Nordic countries. *Acta Oncol* 48(5):646-790. 10.1080/02841860902913546.
- Purdue MP, Stewart PA, Friesen MC, et al. 2017. Occupational exposure to chlorinated solvents and kidney cancer: A case-control study. *Occup Environ Med* 74(4):268-274. <http://doi.org/10.1136/oemed-2016-103849>.
- Radican L, Blair A, Stewart P, et al. 2008. Mortality of aircraft maintenance workers exposed to trichloroethylene and other hydrocarbons and chemicals: Extended follow-up. *J Occup Environ Hyg* 50:1306-1319.
- Radican L, Wartenberg D, Rhoads GG, et al. 2006. A retrospective occupational cohort study of end-stage renal disease in aircraft workers exposed to trichloroethylene and other hydrocarbons. *J Occup Environ Med* 48(Jan):1-12.
- Raisanen J, Niemela R, Rosenberg C. 2001. Tetrachloroethylene emissions and exposure in dry cleaning. *J Air Waste Manag Assoc* 51(12):1671-1675.

9. REFERENCES

- Rajamanikandan S, Sindhu T, Durgapriya D, et al. 2012. Protective effect of *Mollugo nudicaulis* Lam. on acute liver injury induced by perchloroethylene in experimental rats. *Asian Pac J Trop Med* 5(11):862-867.
- Rampy LW, Quast JF, Balmer MF, et al. 1978. Results of a long-term inhalation toxicity study on rats of a perchloroethylene (tetrachloroethylene) formulation. Dow Chemical. Submitted to the U.S. Environmental Protection Agency under TSCA, Section 8d. OTS0515968.
- Ramsey JD, Flanagan RJ. 1982. Detection and identification of volatile organic compounds in blood by headspace gas chromatography as an aide to the diagnosis of solvent abuse. *J Chromatogr* 240:423-444.
- Rao HV, Brown DR. 1993. A physiologically based pharmacokinetic assessment of tetrachloroethylene in groundwater for a bathing and showering determination. *Risk Anal* 13(1):37-49.
- Rao VR, Levy K, Lustik M. 1993. Logistic regression of inhalation toxicities of perchloroethylene-Application in noncancer risk assessment. *Regul Toxicol Pharmacol* 18:233-247.
- Rasmussen RA, Harsch DE, Sweany PH, et al. 1977. Determination of atmospheric halocarbons by a temperature-programmed gas chromatographic freezeout concentration method. *J Air Pollut Control Assoc* 27:579-581.
- Rastkari N, Yunesian M, Ahmadkhaniha R. 2011. Exposure assessment to trichloroethylene and perchloroethylene for workers in the dry cleaning industry. *Bull Environ Contam Toxicol* 86(4):363-367.
- *Ratnoff WD, Gress RE. 1980. The familial occurrence of polycythemia vera: Report of a father and a son, with consideration of the possible etiologic role of exposure to organic solvents, including tetrachloroethylene. *Blood* 56:233-236.
- Rea WJ, Pan Y, Laseter JL, et al. 1987. Toxic volatile organic hydrocarbons in chemically sensitive patients. *Clin Ecol* 5:70-74.
- Reinhardt CF, Mullin LS, Maxfield ME. 1973. Epinephrine-induced cardiac arrhythmia potential of some common industrial solvents. *J Occup Med* 15:953-955.
- Reitz, RH, Gargas ML, Mendrala AL, et al. 1996. *In vivo* and *in vitro* studies of perchloroethylene metabolism for physiologically based pharmacokinetic modeling in rats, mice, and humans. *Toxicol Appl Phamacol* 136:289-306.
- RePORTER. 2013. Tetrachloroethylene. National Institutes of Health, Research Portfolio Online Reporting Tools. <http://projectreporter.nih.gov/reporter.cfm>. August 28, 2013.
- RePORTER. 2018. Tetrachloroethylene. National Institutes of Health, Research Portfolio Online Reporting Tools. <http://projectreporter.nih.gov/reporter.cfm>. September 6, 2018.
- Riihimaki V, Pfaffli, P. 1978. Percutaneous absorption of solvent vapors in man. *Scand J Work Environ Health* 4:73-85.

9. REFERENCES

- Robb EV, McGaughey JF, Sykes AL, et al. 1986. Recovery of principal organic hazardous constituents and products of incomplete combustion from a volatile organic sampling train. Research Triangle Park, NC: U.S. Environmental Protection Agency, Air and Energy Engineering Research Lab. EPA/600/7-86/1025.
- Roberts PV, Dandliker PG. 1983. Mass transfer of volatile organic contaminants from aqueous solution to the atmosphere during surface aeration. *Environ Sci Technol* 17:484-489.
- Roberts PV, Goltz MN, Mackay DM. 1986. A natural gradient experiment on solute transport in a sand aquifer. 3. Retardation estimates and mass balance for organic solutes. *Water Resour Res* 22(13):2047-2058.
- Roda C, Kousignian I, Ramond A, et al. 2013. Indoor tetrachloroethylene levels and determinants in Paris dwellings. *Environ Res* 120:1-6.
- Rodriguez CE, Mahle DA, Gearhart JM, et al. 2007. Predicting age-appropriate pharmacokinetics of six volatile organic compounds in the rat utilizing physiologically based pharmacokinetic modeling. *Toxicol Sci* 98(1):43-56.
- Rogan WJ. 1986. Comment. *J Am Stat Assoc* 81(395):602-603.
- Rosengren LE, Kjellstrand P, Haglid KG. 1986. Tetrachloroethylene: Levels of DNA and S-100 in the gerbil CNS after chronic exposure. *Neurobehav Toxicol Teratol* 8:201-206.
- Rowe BL, Toccalino PL, Moran MJ, et al. 2007. Occurrence and potential human-health relevance of volatile organic compounds in drinking water from domestic wells in the United States. *Environ Health Perspect* 115(11):1539-1546.
- Rowe VK, McCollister DD, Spencer HC, et al. 1952. Vapor toxicity of tetrachloroethylene for laboratory animals and human subjects. *AMA Arch Ind Hyg Occup Med* 5:566-579.
- Ruckart PZ, Bove FJ, Maslia M. 2013. Evaluation of exposure to contaminated drinking water and specific birth defects and childhood cancers at Marine Corps Base Camp Lejeune, North Carolina: A case-control study. *Environ Health* 12(104):10.1186. 10.1186/1476-069X-12-104.
- Ruckart PZ, Bove FJ, Maslia M. 2014. Evaluation of contaminated drinking water and preterm birth, small for gestational age, and birth weight at Marine Corps Base Camp Lejeune, North Carolina: A cross-sectional study. *Environ Health* 13:99. 10.1186/1476-069x-13-99.
- Ruckart PZ, Bove FJ, Shanley E, 3rd, et al. 2015. Evaluation of contaminated drinking water and male breast cancer at Marine Corps Base Camp Lejeune, North Carolina: A case control study. *Environ Health* 14:74. 10.1186/s12940-015-0061-4.
- Ruder AM, Ward EM, Brown DP. 1994. Cancer mortality and female and male dry cleaning workers. *J Occup Environ Med* 36(8):867-874.
- Ruder AM, Ward EM, Brown DP. 2001. Mortality in dry-cleaning workers: An update. *Am J Ind Med* 39(2):121-132.

9. REFERENCES

- Ruder AM, Yiin JH, Waters MA, et al. 2013. The Upper Midwest Health Study: Gliomas and occupational exposure to chlorinated solvents. *Occup Environ Med* 70(2):73-80. 10.1136/oemed-2011-100588.
- Ruiz P, Ray M, Fisher J, et al. 2011. Development of a human physiologically based pharmacokinetic (PBPK) toolkit for environmental pollutants. *Int J Mol Sci* 12:7469-7480. 10.3390/ijms12117469.
- Rutkiewicz I, Jakubowska N, Polkowska Z, et al. 2011. Monitoring of Occupational Exposure to volatile organohalogen solvents (VOXs) in human urine samples of dry-cleaner workers by TLHS-DAI-GC-ECD procedure. *Ind Health* 49(1):126-132.
- Ryoo D, Shim H. 2000. Aerobic degradation of tetrachloroethylene by toluene-o-xylene monooxygenase of *Pseudomonas stutzeri* OX1. *Nat Biotechnol* 18(7):775-778.
- Sack TM, Steele DH, Hammerstrom K, et al. 1992. A survey of household products for volatile organic compounds. *Atmos Environ* 26(6):1063-1070.
- Saisho. 1994. Bioaccumulation of volatile chlorinated hydrocarbons in blue mussel, *Mytilus edulus*, and killifish, *Oryzias latipes*. *Jpn J Toxicol Environ Health* 40:274-278.
- Saland G. 1967. Accidental exposure to perchloroethylene. *N Y State J Med* 67:2359-2361.
- Sallmén M, Lindbohm ML, Anttila A, et al. 1998. Time to pregnancy among the wives of men exposed to organic solvents. *Occup Environ Med* 55(1):24-30.
- Sallmén M, Lindbohm ML, Kyyronen P, et al. 1995. Reduced fertility among women exposed to organic solvents. *Am J Ind Med* 27:699-713.
- Sandground JH. 1941. Coma following medication with tetrachloroethylene. *J Am Med Assoc* 117:440-441.
- *Santibanez M, Vioque J, Alguacil J, et al. 2008. Occupational exposures and risk of oesophageal cancer by histological type: A case-control study in eastern Spain. *Occup Environ Med* 65(11):774-781.
- Sarangapani R, Gentry PR, Covington TR, et al. 2003. Evaluation of the potential impact of age- and gender-specific lung morphology and ventilation rate on the dosimetry of vapors. *Inhal Toxicol* 15(10):987-1016.
- Saunders NR, Ek CJ, Habgood MD, et al. 2008. Barriers in the brain: A renaissance? *Trends Neurosci* 31(6):279-286. 10.1016/j.tins.2008.03.003.
- Saunders NR, Liddelow SA, Dziegielewska KM. 2012. Barrier mechanisms in the developing brain. *Front Pharmacol* 3:46. 10.3389/fphar.2012.00046.
- Savolainen H, Pfaffli P, Tengen M, et al. 1977. Biochemical and behavioral effects of inhalation exposure to tetrachloroethylene and dichloromethane. *J Neuropathol Exp Neurol* 36:941-949.
- Scheuplein R, Charnley G, Dourson M. 2002. Differential sensitivity of children and adults to chemical toxicity. I. Biological basis. *Regul Toxicol Pharmacol* 35(3):429-447.

9. REFERENCES

- Schlichting LM, Wright PFA, Stacey NH. 1992. Effects of tetrachloroethylene on hepatic and splenic lymphocytotoxic activities in rodents. *Toxicol Ind Health* 8(5):255-266.
- Schreiber JS. 1993. Predicted infant exposure to tetrachloroethylene in human breast milk. *Risk Anal* 13(5):515-524.
- Schreiber JS, House S, Prohonic E, et al. 1993. An investigation of indoor air contamination in residences above dry cleaners. *Risk Anal* 13(3):335-344.
- Schreiber JS, Hudnell HK, Geller AM, et al. 2002. Apartment residents' and day care workers' exposures to tetrachloroethylene and deficits in visual contrast sensitivity. *Environ Health Perspect* 110(7):655-664.
- Schroers HJ, Jermann E, Begerow J, et al. 1998. Determination of physiological levels of volatile organic compounds in blood using static headspace capillary gas chromatography with serial triple detection. *Analyst* 123(4):715-720.
- Schultz B, Kjeldsen P. 1986. Screening of organic matter in leachates from sanitary landfills using gas chromatography combined with mass spectrometry. *Water Res* 20:965-970.
- Schumann AM, Quast JF, Watanabe PG. 1980. The pharmacokinetics and macromolecular interactions of perchloroethylene in mice and rats as related to oncogenicity. *Toxicol Appl Pharmacol* 55:207-219.
- Schwarzenbach RP, Giger W, Hoehn E, Schneider JK. 1983. Behavior of organic compounds during infiltration of river water to ground water. Field studies. *Environ Sci Technol* 17:472-479.
- Schwarzenbach RP, Molnar-Kubica E, Giger W, et al. 1979. Distribution, residence time, and fluxes of tetrachloroethylene and 1,4-dichlorobenzene in Lake Zurich, Switzerland. *Environ Sci Technol* 13:1367-1373.
- Schwetz BA, Leong BKJ, Gehring PJ. 1975. The effect of maternally inhaled trichloroethylene, perchloroethylene, methyl chloroform, and methylene chloride on embryonal and fetal development in mice and rats. *Toxicol Appl Pharmacol* 32:84-96.
- Seeber A. 1989. Neurobehavioral toxicity of long-term exposure to tetrachloroethylene. *Neurotoxicol Teratol* 11:579-583.
- Seidel HJ, Weber L, Barthel E. 1992. Hematological toxicity of tetrachloroethylene in mice. *Arch Toxicol* 66:228-230.
- *Seidler A, Mohner M, Berger J, et al. 2007. Solvent exposure and malignant lymphoma: A population-based case-control study in Germany. *J Occup Med Toxicol* 2:2. 10.1186/1745-6673-2-2.
- Seiji K, Inoue O, Jin C, et al. 1989. Dose-excretion relationship in tetrachloroethylene-exposed workers and the effect of tetrachloroethylene co-exposure on trichloroethylene metabolism. *Am J Ind Med* 16:675-684.
- Seiji K, Jin C, Watanabe T, et al. 1990. Sister chromatid exchanges in peripheral lymphocytes of workers exposed to benzene, trichloroethylene, or tetrachloroethylene, with references to smoking habits. *Int Arch Occup Environ Health* 62:171-176.

9. REFERENCES

- Seip HM, Alstad J, Carlberg GE, et al. 1986. Measurement of mobility of organic compounds in soils. *Sci Total Environ* 50:87-101.
- Selden AI, Ahlborg G, Jr. 2011. Cancer morbidity in Swedish dry-cleaners and laundry workers: Historically prospective cohort study. *Int Arch Occup Environ Health* 84(4):435-443.
- Seo M, Ikeda K, Okamura T, et al. 2008b. Enhancing effect of chlorinated organic solvents on histamine release and inflammatory mediator production. *Toxicology* 243(1-2):75-83.
- Seo M, Yamagiwa T, Kobayashi R, et al. 2008a. A small amount of tetrachloroethylene ingestions from drinking water accelerates antigen-stimulated allergic responses. *Immunobiology* 213(8):663-669.
- Seo M, Kobayashi R, Okamura T, et al. 2012. Enhancing effects of trichloroethylene and tetrachloroethylene on type I allergic responses in mice. *J Toxicol Sci* 37(2):439-445.
- Sexton K, Adgate JL, Church TR, et al. 2005. Children's exposure to volatile organic compounds as determined by longitudinal measurements in blood. *Environ Health Perspect* 113(3):342-349.
- Shafer TJ, Bushnell PJ, Benignus VA, et al. 2005. Perturbation of voltage-sensitive Ca²⁺ channel function by volatile organic solvents. *J Pharmacol Exp Ther* 315(3):1109-1118.
- Sharanjeet-Kaur (no initial), Mursyid A, Kamaruddin A, et al. 2004. Effect of petroleum derivatives and solvents on colour perception. *Clin Exp Optom* 87(4-5):339-343.
- Sharma PK, McCarty PL. 1996. Isolation and characterization of a facultatively aerobic bacterium that reductively dehalogenates tetrachloroethene to *cis*-1,2-dichloroethene. *Appl Environ Microbiol* 62(3):761-765.
- Shigetoshi Y, Nagahito M, Tomomichi M, et al. 1993. European patent application. Process for the manufacture of tetrachloroethylene. Application number 93109052.6.
- Shimada T, Swanson A, Leber P, et al. 1983. The evaluation for genotoxicity of several halogenated solvents. *Environ Mutagen* 5:447.
- Shusterman D. 2018. Trichloroethane, trichloroethylene, and tetrachloroethylene. In: Olson KR, Anderson IB, Benowitz NL, et al., eds. *Poisoning and drug overdose*. 7th ed. McGraw-Hill Education.
- Silver SR, Pinkerton LE, Fleming DA, et al. 2014. Retrospective cohort study of a microelectronics and business machine facility. *Am J Ind Med* 57(4):412-424. 10.1002/ajim.22288.
- Singh HB, Lillian D, Appleby A, et al. 1975. Atmospheric formation of carbon tetrachloride from tetrachloroethylene. *Environ Lett* 10:253-256.
- Singh HB, Salas LJ, Stiles RE. 1982. Distribution of selected gaseous organic mutagens and suspect carcinogens in ambient air. *Environ Sci Technol* 16:872-880.
- Skender L, Karacic V, Bosner B, et al. 1993. Assessment of exposure to trichloroethylene and tetrachloroethylene in the population of Zagreb, Croatia. *Int Arch Occup Environ Health* 65:S163-S165.

9. REFERENCES

- Skender L, Karacic V, Bosner B, et al. 1994. Assessment of urban population exposure to trichloroethylene and tetrachloroethylene by means of biological monitoring. *Arch Environ Health* 49(6):445-451.
- Skender LJ, Karacic V, Prpic-Majic D. 1991. A comparative study of human levels of trichloroethylene and tetrachloroethylene after occupational exposure. *Arch Environ Health* 46(3):174-178.
- *Smith EM, Miller ER, Woolson RF, et al. 1985. Bladder cancer risk among laundry workers, dry cleaners, and others in chemically related occupations. *J Occup Med* 27:295-297.
- Solet D, Robins TG. 1991. Renal function in dry cleaning workers exposed to perchloroethylene. *Am J Ind Med* 20:601-614.
- Solet D, Robins TG, Sampaio C. 1990. Perchloroethylene exposure assessment among dry cleaning workers. *Am Ind Hyg Assoc J* 51(10):566-574.
- Soni M, Nomiyama H, Nomiyama K. 1990. Chronic inhalation effects of tetrachloroethylene on hepatic and renal microsomal electron transport components and δ -aminolevulinic acid dehydratase in rats. *Toxicol Lett* 54:207-213.
- Sonnenfeld N, Hertz-Picciotto I, Kaye WE. 2001. Tetrachloroethylene in drinking water and birth outcomes at the US Marine Corps Base at Camp Lejeune, North Carolina. *Am J Epidemiol* 154(10):902-908.
- Spearow JL, Gettmann K, Wade M. 2017. Review: Risk assessment implications of variation in susceptibility to perchloroethylene due to genetic diversity, ethnicity, age, gender, diet and pharmaceuticals. *Hum Ecol Risk Assess* 23(6):1466-1492.
- Spencer AB, Earnest GS, Kovein RJ. 1995. In-depth survey report: Exposures during the spotting procedure in a commercial dry cleaner at Widmer's Dry Cleaning, Cincinnati, Ohio. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health. Report No. ECTB 201-13b.
- Spicer CW, Holdren MW, Slivon LE, et al. 1986. Intercomparison of sampling techniques for toxic organic compounds in indoor air. Proceedings of the 1986 EPA/APCA Symposium on Measurement of Toxic Air Pollutants, Raleigh, NC, April 27-30, 1986, Environmental Monitoring Systems Laboratory, U.S. Environmental Protection Agency, EPA600/986013, 45-60.
- Spirtas R, Stewart PA, Lee JS, et al. 1991. Retrospective cohort mortality study of workers at an aircraft maintenance facility. I. Epidemiological results. *Br J Ind Med* 48(8):515-530.
- SRI. 2011. 2011 Directory of chemical producers United States. In: 2011 Directory of chemical producers United States. SRI Consulting, 744-745.
- *Stemhagen A, Slade J, Altman R et al. 1983. Occupational risk factors and liver cancer: A retrospective case-control study of primary liver cancer in New Jersey. *Am J Epidemiol* 117:443-454.
- Stewart RD. 1969. Acute tetrachloroethylene intoxication. *J Am Med Assoc* 208:1490-1492.
- Stewart RD, Dodd HC. 1964. Absorption of carbon tetrachloride, trichloroethylene, tetrachloroethylene, methylene chloride, and 1,1,1-trichloroethane through the human skin. *Am Ind Hyg Assoc J* 25:439-446.

9. REFERENCES

- Stewart RD, Baretta ED, Dodd HC, et al. 1970. Experimental human exposure to tetrachloroethylene. *Arch Environ Health* 20:224-229.
- Stewart RD, Erley DS, Schaffer AW, et al. 1961a. Accidental vapor exposure to anesthetic concentrations of a solvent containing tetrachloroethylene. *Ind Med Surg* 30:327-330.
- Stewart RD, Gay HH, Erley DS, et al. 1961b. Human exposure to tetrachloroethylene vapor: Relationship of expired air and blood concentrations to exposure and toxicity. *Arch Environ Health* 2:516-522.
- Stewart RD, Hake CL, Forster HV, et al. 1981. Tetrachloroethylene: Development of a biologic standard for the industrial worker by breath analysis. Cincinnati, OH: National Institute for Occupational Safety and Health. Contract no. HSM 99-72-84. NIOSH-MCOW-ENVM-PCE-74-6. PB82152166.
- Stewart RD, Hake CL, Wu A, et al. 1977. Effects of perchloroethylene/drug interaction on behavior and neurological function. Final report. Washington, DC: National Institute for Occupational Safety and Health. PB83174607.
- Storm JE, Mazor KA, Aldous KM, et al. 2011. Visual contrast sensitivity in children exposed to tetrachloroethylene. *Arch Environ Occup Health* 66(3):166-177.
- Storm JE, Mazor KA, Shost SJ, et al. 2013. Socioeconomic disparities in indoor air, breath, and blood perchloroethylene level among adult and child residents of buildings with or without a dry cleaner. *Environ Res* 122:88-97.
- Story DL, Meierhenry EF, Tyson CA, et al. 1986. Differences in rat liver enzyme-altered foci produced by chlorinated aliphatics and phenobarbital. *Toxicol Ind Health* 2:351-362.
- Stroo HF, Leeson A, Marqusee JA, et al. 2012. Chlorinated ethene source remediation: Lessons learned. *Environ Sci Technol* 46(12):6438-6447.
- Struwe M, Zeller A, Singer T, et al. 2011. Possibility of methodical bias in analysis of comet assay studies: Re: DNA damage detected by the alkaline comet assay in the liver of mice after oral administration of tetrachloroethylene. *Mutagenesis* 25:133-138.
- *Stutz DR, Ulin S. 1992. Hazardous materials injuries: A handbook for pre-hospital care. Third edition. Beltsville, MD: Bradford Communications Corporations, 286-287.
- Su C, Goldberg ED. 1976. Environmental concentration and fluxes of some halocarbons. In: Windom HL, Duce, KA, eds. *Marine pollutant transfer*. Lexington, MA, 353-374.
- Suber RL. 1989. Clinical pathology for toxicologists. In: Hayes AW, ed. *Principles and methods of toxicology*. 2nd ed. New York, NY: Raven Press, 485-519.
- Sullivan DA, Jones AD, Williams JG. 1985. Results of the U.S. Environmental Protection Agency's air toxics analysis in Philadelphia. In: *Proceedings of the 78th Annual Meeting of the Air Pollution Control Association*. Vol. 2, 1-15.
- Suzaki H, Aoki A, Nomura Y. 1997. Regeneration of olfactory mucosa in mice after inhalation exposure to perchloroethylene. *J Otorhinolaryngol Relat Spec* 59(2):91-96.

9. REFERENCES

- Swan SH, Robins JM. 1986. Comment. *J Am Stat Assoc* 81(395):604-609.
- Swann RL, Laskowski DA, McCall PJ, et al. 1983. A rapid method for the estimation of the environmental parameters octanol/water partition coefficient, soil sorption constant, water to air ratio, and water solubility. *Res Rev* 85:17-28.
- Swenberg JA, Short B, Borghoff S, et al. 1989. The comparative pathobiology of a $\alpha_2\mu$ -globulin nephropathy. *Toxicol Appl Pharmacol* 97:35-46.
- Szakmary E, Ungvary G, Tatrai E. 1997. The offspring-damaging effect of tetrachloroethylene in rats, mice and rabbits. *Cent Eur J Occup Environ Med* 3(1):33-39.
- Tabak HH, Quave SA, Mashni CI, et al. 1981. Biodegradability studies with organic priority pollutant compounds. *J Water Pollut Control Fed* 53:1503-1518.
- Talibov M, Auvinen A, Weiderpass E, et al. 2017. Occupational solvent exposure and adult chronic lymphocytic leukemia: No risk in a population-based case-control study in four Nordic countries. *Int J Cancer* 141(6):1140-1147. <http://doi.org/10.1002/ijc.30814>.
- Tanios MA, El Gamal H, Rosenberg BJ, et al. 2004. Can we still miss tetrachloroethylene-induced lung disease? The emperor returns in new clothes. *Respiration* 71(6):642-645.
- Taskinen H, Anttila A, Lindbohm ML, et al. 1989. Spontaneous abortions and congenital malformations among the wives of men occupationally exposed to organic solvents. *Scand J Work Environ Health* 15:345-352.
- Thomas K, Colborn T. 1992. Organochlorine endocrine disruptors in human tissue. In: Colborn T, Clement C, eds. *Chemically induced alterations in sexual and functional development: The wildlife/human connection*. Princeton, NJ: Princeton Scientific Publishing, 365-394.
- Thomas RG. 1982. Volatilization from water. In: Lyman WJ, Reehl WF, Rosenblatt DH, eds. *Handbook of chemical property estimation methods: Environmental behavior of organic compounds*. New York, NY: McGraw-Hill, 1-2, 13, 25.
- Thompson KM, Evans JS. 1993. Workers' breath as a source of perchloroethylene (pert) in the home. *J Expo Anal Environ Epidemiol* 3(4):417-430.
- Tichenor BA, Sparks LE, Jackson MD, et al. 1990. Emissions of perchloroethylene from dry cleaned fabrics. *Atmos Environ* 24A(5):1219-1229.
- Till C, Rovet JF, Koren G, et al. 2003. Assessment of visual functions following prenatal exposure to organic solvents. *Neurotoxicology* 24(4-5):725-731.
- Tinston DJ. 1995. Perchloroethylene: Multigeneration inhalation study in the rat. Report No: CTL/P/4097. Sponsored by Halogenated Solvents Industry Alliance, HSIA/90/0002.
- 't Mannetje A, De Roos AJ, Boffetta P, et al. 2016. Occupation and risk of non-Hodgkin lymphoma and its subtypes: A pooled analysis from the InterLymph Consortium. *Environ Health Perspect* 124(4):396-405. 10.1289/ehp.1409294.

9. REFERENCES

- Toraason M, Butler MA, Ruder A, et al. 2003. Effect of perchloroethylene, smoking, and race on oxidative DNA damage in female dry cleaners. *Mutat Res* 539(1-2):9-18.
- Toraason M, Clark J, Dankovic D, et al. 1999. Oxidative stress and DNA damage in Fischer rats following acute exposure to trichloroethylene or perchloroethylene. *Toxicology* 138(1):43-53.
- TRI16. 2018. TRI explorer: Providing access to EPA's toxics release inventory data. Washington, DC: U.S. Environmental Protection Agency, Office of Information Analysis and Access. Office of Environmental Information. Toxics Release Inventory. <http://www.epa.gov/triexplorer/>. June 1, 2018.
- Tsulaya VR, Bonashevskaya TI, Zykova VV, et al. 1977. [The toxicological characteristics of some chlorinated hydrocarbons]. *Gig Sanit* 8:50-53. (Russian)
- Tsuruta H. 1975. Percutaneous absorption of organic solvents. (1) Comparative study of the *in vivo* percutaneous absorption of chlorinated solvents in mice. *Ind Health* 13:227-236.
- Tsuruta H. 1977. Percutaneous absorption of organic solvents. (2) A method for measuring the penetration rate of chlorinated solvents through excised rat skin. *Ind Health* 15:131-139.
- Tsuruta H. 1989. Skin absorption of organic solvent vapors in nude mice *in vivo*. *Ind Health* 27:37-47.
- Tu AS, Murray TA, Hatch KM, et al. 1985. *In vitro* transformation of BALB/C-3T3 cell by chlorinated ethanes and ethylenes. *Cancer Lett* 28:85-92.
- Tucker JD, Sorensen KJ, Ruder AM, et al. 2011. Cytogenetic analysis of an exposed-referent study: Perchloroethylene-exposed dry cleaners compared to unexposed laundry workers. *Environ Health* 10:16.
- Union Carbide. 1962. Comparative inhalation toxicities of perchloroethylene. Submitted to the U.S. Environmental Protection Agency under TSCA, Section 8d. OTS0515591.
- Urano K, Murata C. 1985. Adsorption of principal chlorinated organic compounds on soil. *Chemosphere* 14:293-299.
- USGS. 2003. Quality and sources of shallow ground water in areas of recent residential development in Salt Lake Valley, Salt Lake County, Utah. U.S. Geological Survey, U.S. Department of the Interior, National Water-Quality Assessment Program.
- USGS. 2006. Volatile organic compounds in the nation's ground water and drinking-water supply wells. United States Geological Survey, U.S. Department of the Interior, National Water-Quality Assessment Program.
- USGS. 2012. Assessment of soil-gas contamination at three former fuel-dispensing sites, Fort Gordon, Georgia, 2010-2011. United States Geological Survey, U.S. Department of the Interior.
- USITC. 2013. Tetrachloroethylene (Perchloroethylene) import and export data. United States International Trade Commission. <http://www.usitc.gov>. March 11, 2013.
- Valencia R, Mason JM, Woodruff RC, et al. 1985. Chemical mutagenesis testing in *Drosophila* III. Results of 48 coded compounds tested for the National Toxicology Program. *Environ Mutagen* 7:325-348.

9. REFERENCES

- Valic E, Waldhor T, Konnaris C, et al. 1997. Acquired dyschromatopsia in combined exposure to solvents and alcohol. *Int Arch Occup Environ Health* 70(6):403-406.
- Vamvakas S, Herkenhoff M, Dekant W, et al. 1989. Mutagenicity of tetrachloroethene in the Ames test: Metabolic activation by conjugation with glutathione. *J Biochem Toxicol* 4:21-27.
- *van der Mark M, Vermeulen R, Nijssen PC, et al. 2015. Occupational exposure to solvents, metals and welding fumes and risk of Parkinson's disease. *Parkinsonism Relat Disord* 21(6):635-639. 10.1016/j.parkreldis.2015.03.025.
- Van Duuren BL, Goldschmidt BM, Loewengart G, et al. 1979. Carcinogenicity of halogenated olefinic and aliphatic hydrocarbons in mice. *J Natl Cancer Inst* 63:1433-1439.
- Vartiainen T, Pukkala E, Rienoja T, et al. 1993. Population exposure to tri- and tetrachloroethene and cancer risk: Two cases of drinking water pollution. *Chemosphere* 27(7):1171-1181.
- Vaughan TL, Stewart PA, Davis S, et al. 1997. Work in dry cleaning and the incidence of cancer of the oral cavity, larynx, and oesophagus. *Occup Environ Med* 54(9):692-695.
- Veith GD, Macek KJ, Petrocelli SR, et al. 1980. An evaluation of using partition coefficients and water solubility to estimate bioconcentration factors for organic chemicals in fish. In: Eaton JG, Parrish PR, Hendricks AC, eds. *Aquatic Toxicology*. Am Soc Test Mater, ASTM STP 707, 116-129.
- *Vemmer H, Rohleder K, Wieczorek H, et al. 1984. [Perchloroethylene in the tissues of fattening pigs]. *Landwirtsch Forsch* 37:36-43. (German)
- Verplanke AJ, Leummens MH, Herber RF. 1999. Occupational exposure to tetrachloroethene and its effects on the kidneys. *J Occup Environ Med* 41(1):11-16.
- Verschueren K. 1983. *Handbook of environmental data on organic chemicals*. New York, NY: Van Nostrand Reinhold, 1076-1080.
- Viau C, Bernard A, Lauwerys R, et al. 1987. A cross-sectional survey of kidney function in refinery employees. *Am J Ind Med* 11:177-187.
- Vieira I, Sonnier M, Cresteil T. 1996. Developmental expression of CYP2E1 in the human liver: Hypermethylation control of gene expression during the neonatal period. *Eur J Biochem* 238(2):476-483.
- Vieira V, Aschengrau A, Ozonoff D. 2005. Impact of tetrachloroethylene-contaminated drinking water on the risk of breast cancer: Using a dose model to assess exposure in a case-control study. *Environ Health* 4(1):3.
- Vingrys AJ, King-Smith PE. 1988. A quantitative scoring technique for panel tests of color vision. *Invest Ophthalmol Vis Sci* 29:50-63.
- Vizcaya D, Christensen KY, Lavoue J, et al. 2013. Risk of lung cancer associated with six types of chlorinated solvents: Results from two case-control studies in Montreal, Canada. *Occup Environ Med* 70(2):81-85. 10.1136/oemed-2012-101155.

9. REFERENCES

- Vlaanderen J, Straif K, Pukkala E, et al. 2013. Occupational exposure to trichloroethylene and perchloroethylene and the risk of lymphoma, liver, and kidney cancer in four Nordic countries. *Occup Environ Med* 70(6):393-401. 10.1136/oemed-2012-101188.
- Vlaanderen J, Straif K, Ruder A, et al. 2014. Tetrachloroethylene exposure and bladder cancer risk: A meta-analysis of dry-cleaning-worker studies. *Environ Health Perspect* 122(7):661-666. 10.1289/ehp.1307055.
- Vogel TM, McCarty PL. 1985. Biotransformation of tetrachloroethylene to trichloroethylene, dichloroethylene, vinyl chloride and carbon dioxide under methanogenic conditions. *Appl Environ Microbiol* 49:1080-1083.
- Volkel W, Friedewald M, Lederer E, et al. 1998. Biotransformation of perchloroethene: Dose-dependent excretion of trichloroacetic acid, dichloroacetic acid, and N-acetyl-S-(trichlorovinyl)-L-cystein in rats and humans after inhalation. *Toxicol Appl Pharmacol* 153(1):20-27.
- Volkel W, Pahler A, Dekant W. 1999. Gas chromatography-negative ion chemical ionization mass spectrometry as a powerful tool for the detection of mercapturic acids and DNA and protein adducts as biomarkers of exposure to halogenated olefins. *J Chromatogr A* 847(1-2):35-46.
- Voss JU, Roller M, Brinkmann E, et al. 2005. Nephrotoxicity of organic solvents: Biomarkers for early detection. *Int Arch Occup Environ Health* 78(6):475-485.
- Vroblesky DA. 1991. Mapping zones of contaminated ground-water discharge using creek-bottom-sediment vapor samplers, Aberdeen Proving Ground, Maryland. *Ground Water* 29(1):7-12.
- Vyskocil A, Emminger S, Tejral J. 1990. Study on kidney function in female workers exposed to perchloroethylene. *Hum Exp Toxicol* 9:377-380.
- Wakeham SG, Davis AC, Karas JL. 1983. Mesocosm experiments to determine the fate and persistence of volatile organic compounds in coastal seawater. *Environ Sci Technol* 17:611-617.
- Wall JM, Carreon RF. 1984. Perchloroethylene (99.97%) formulation: Acute toxicological properties. Dow Chemical. Submitted to the U.S. Environmental Protection Agency under TSCA, Section 8d. OTS0515966.
- Wallace LA. 1986. Personal exposures, indoor and outdoor air concentrations and exhaled breath concentrations of selected volatile organic compounds measured for 600 residents of New Jersey, North Dakota, North Carolina, and California. *Toxicol Environ Chem* 12:215-236.
- Wallace LA, Pellizzari ED, Hartwell TD, et al. 1986a. Concentrations of 20 volatile organic compounds in the air and drinking water of 350 residents of New Jersey compared with concentrations in their exhaled breath. *J Occup Med* 28:603-608.
- Wallace LA, Pellizzari ED, Hartwell TD, et al. 1986b. Results from the first three seasons of the TEAM study: Personal exposures, indoor-outdoor relationships, and breath levels of toxic air pollutants measured for 355 persons in New Jersey. In: Kopfler PC, Craun G, eds. *Environmental epidemiology*. Chelsea, MI: Lewis Publishers, 181-200.

9. REFERENCES

- Wallace LA, Pellizzari ED, Hartwell TD. 1986c. Total exposure assessment methodology (TEAM) study: Personal exposures, indoor-outdoor relationships, and breath levels of volatile organic compounds in New Jersey. *Environ Int* 12:369-387.
- Wallace LA, Pellizzari ED, Leaderer B, et al. 1987. Emissions of volatile organic compounds from building materials and consumer products. *Atmos Environ* 21:385-393.
- Wallace LA, Pellizzari ED, Sheldon L, et al. 1986d. The total exposure assessment methodology (TEAM) study: Direct measurements of personal exposures through air and water for 600 residents of several U.S. cities. In: Cohen Y, ed. *Pollutants in a multimedia environment*. New York, NY: Plenum Publishing Corp., 289-315.
- Walles SAS. 1986. Induction of single-strand breaks in DNA of mice by trichloroethylene and tetrachloroethylene. *Toxicol Lett* 31:31-35.
- *Wang G, Wang J, Ansari GAS, et al. 2017. Autoimmune potential of perchloroethylene: Role of lipid-derived aldehydes. *Toxicol Appl Pharmacol* 333:76-83. <http://doi.org/10.1016/j.taap.2017.08.009>.
- Wang JL, Chen WL, Tsai SY, et al. 2001. An *in vitro* model for evaluation of vaporous toxicity of trichloroethylene and tetrachloroethylene to CHO-K1 cells. *Chem Biol Interact* 137(2):139-154.
- Wang S, Karlsson JE, Kyrklund T, et al. 1993. Perchloroethylene-induced reduction in glial and neuronal cell marker proteins in rat brain. *Pharmacol Toxicol* 72:273-278.
- Ward RC, Travis CC, Hetrick DM, et al. 1988. Pharmacokinetics of tetrachloroethylene. *Toxicol Appl Pharmacol* 93:108-117.
- Warren DA, Reigle TG, Muralidhara S, et al. 1996. Schedule-controlled operant behavior of rats following oral administration of perchloroethylene: Time course and relationship to blood and brain solvent levels. *J Toxicol Environ Health* 47(4):345-362.
- Watanabe K, Sasaki T, Kawakami K. 1998. Comparisons of chemically-induced mutation among four bacterial strains, *Salmonella typhimurium* TA102 and TA2638, and *Escherichia coli* WP2/pKM101 and WP2 *uvrA*/pKM101: Collaborative study III and evaluation of the usefulness of these strains. *Mutat Res* 416(3):169-181.
- Wayne LG, Orcutt JA. 1960. The relative potentials of common organic solvents as precursors of eye-irritants in urban atmospheres. *J Occup Med* 2:383-388.
- Weant GE, McCormick GS. 1984. Nonindustrial sources of potentially toxic substances and their applicability to source apportionment methods. EPA450/484/003. PB84231232.
- Webler T, Brown HS. 1993. Exposure to tetrachloroethylene via contaminated drinking water pipes in Massachusetts: A predictive model. *Arch Environ Health* 48(5):293-297.
- *Weiderpass E, Pukkala E, Vasama-Neuvonen K, et al. 2001. Occupational exposures and cancers of the endometrium and cervix uteri in Finland. *Am J Ind Med* 39(6):572-580.
- Wenzel DG, Gibson RD. 1951. A study of the toxicity and anthelmintic activity of n-butylidene chloride. *J Pharmacol* 3:169-176.

9. REFERENCES

- Werner M, Birner G, Dekant W. 1996. Sulfoxidation of mercapturic acids derived from tri- and tetrachloroethene by cytochromes P450 3A: A bioactivation reaction in addition to deacetylation and cysteine conjugate β -lyase mediated cleavage. *Chem Res Toxicol* 9(1):41-49.
- West JR, Smith HW, Chasis H. 1948. Glomerular filtration rate, effective renal blood flow, and maximal tubular excretory capacity in infancy. *J Pediatr* 32:10-18.
- White IN, Razvi N, Gibbs AH, et al. 2001. Neoantigen formation and clastogenic action of HCFC-123 and perchloroethylene in human MCL-5 cells. *Toxicol Lett* 124(1-3):129-138.
- WHO. 1987. Tetrachloroethylene health and safety guide. World Health Organization, Distribution and Sales 1211 Geneva 27, Switzerland, 32.
- WHO. 2003. Tetrachloroethene in drinking-water. Background document for development of WHO guidelines for drinking-water quality. Geneva, Switzerland: World Health Organization.
- WHO. 2010. Guidelines for indoor air quality: Selected pollutants. Geneva, Switzerland: World Health Organization. http://www.euro.who.int/__data/assets/pdf_file/0009/128169/e94535.pdf. April 24, 2013.
- WHO. 2011. Guidelines for drinking-water quality. 4th ed. Geneva, Switzerland: World Health Organization. http://www.who.int/water_sanitation_health/publications/2011/dwq_guidelines/en/index.html. April 24, 2013.
- Widdowson EM, Dickerson JWT. 1964. Chemical composition of the body. In: Comar CL, Bronner F, eds. *Mineral metabolism: An advance treatise. Volume II: The elements Part A*. New York, NY: Academic Press, 1-247.
- Williams P, Benton L, Warmerdam J, et al. 2002. Comparative risk analysis of six volatile organic compounds in California drinking water. *Environ Sci Technol* 36(22):4721-4728.
- Wilson JT, Enfield CG, Dunlap WJ, et al. 1981. Transport and fate of selected organic pollutants in a sandy soil. *J Environ Qual* 10:501-506.
- Wilson JT, McNabb JF, Balkwill DL, et al. 1983. Enumeration and characterization of bacteria indigenous to a shallow water-table aquifer. *Ground Water* 21:134-142.
- Windham GC, Shusterman D, Swan S, et al. 1991. Exposure to organic solvents and adverse pregnancy outcome. *Am J Ind Med* 20:241-259.
- Windham GC, Zhang L, Gunier R, et al. 2006. Autism spectrum disorders in relation to distribution of hazardous air pollutants in the San Francisco Bay area. *Environ Health Perspect* 114(9):1438-1444.
- Won D, Corsi RL, Rynes M. 2000. New indoor carpet as an adsorptive reservoir for volatile organic compounds. *Environ Sci Technol* 34(19):4193-4198.
- Wright WH, Bozicevick J, Gordon LS. 1937. Studies on oxyuriasis. V. Therapy with-single doses of tetrachloroethylene. *J Am Med Assoc* 109:570-573.
- Yllner S. 1961. Urinary metabolites of ^{14}C -tetrachloroethylene in mice. *Nature* 191:820.

9. REFERENCES

- Yokley KA, Evans MV. 2008. Physiological changes associated with aging result in lower internal doses of toluene and perchloroethylene in simulations using pharmacokinetic modeling. *Toxicol Environ Chem* 90(3):475-492.
- Zapór L, Skowron J, Golofit-Szymczak M. 2002. The cytotoxicity of some organic solvents on isolated hepatocytes in monolayer culture. *Int J Occup Saf Ergon* 8(1):121-129.
- Zhou YH, Cichocki JA, Soldatow VY, et al. 2017. Editor's highlight: Comparative dose-response analysis of liver and kidney transcriptomic effects of trichloroethylene and tetrachloroethylene in B6C3F1 mouse. *Toxicol Sci* 160(1):95-110. <http://doi.org/10.1093/toxsci/kfx165>.
- Ziegler EE, Edwards BB, Jensen RL, et al. 1978. Absorption and retention of lead by infants. *Pediatr Res* 12(1):29-34.
- Zielhuis GA, Gijzen R, Van Der Gulden JWJ. 1989. Menstrual disorders among dry cleaning workers. *Scand J Work Environ Health* 15:238.
- Ziener CE, Braunsdorf PP. 2014. Trace analysis in end-exhaled air using direct solvent extraction in gas sampling tubes: Tetrachloroethene in workers as an example. *Int J Anal Chem* 2014:904512. 10.1155/2014/904512.
- Ziglio G, Beltramelli G, Pegliaso F. 1984. A procedure for determining plasmatic trichloroacetic acid in human subjects exposed to chlorinated solvents at environmental levels. *Arch Environ Contam Toxicol* 13:129-134.
- Zoccolillo L, Amendola L, Insogna S. 2009. Comparison of atmosphere/aquatic environment concentration ratio of volatile chlorinated hydrocarbons between temperate regions and Antarctica. *Chemosphere* 76:1525-1532.
- Zytner RG, Biswas N, Bewtra JK. 1989. Volatilization of perchloroethylene from stagnant water and soil. In: Bell JM, ed. *Proceedings of the 43rd Industrial Waste Conference*, Purdue University, May 10-12, 1988. Chelsea, MI: Lewis Publishers, Inc., 101-108.