

9. REFERENCES

- Abdallah MA, Harrad S. 2010. Modification and calibration of a passive air sampler for monitoring vapor and particulate phase brominated flame retardants in indoor air: Application to car interiors. *Environ Sci Technol* 44(8):3059-3065.
- Abdallah MA, Harrad S. 2014. Polybrominated diphenyl ethers in UK human milk: Implications for infant exposure and relationship to external exposure. *Environ Int* 63:130-136.
- Abdallah MA, Harrad S, Covaci A. 2009. Isotope dilution method for determination of polybrominated diphenyl ethers using liquid chromatography coupled to negative ionization atmospheric pressure photoionization tandem mass spectrometry: Validation and application to house dust. *Anal Chem* 81(17):7460-7467.
- Abdallah MA, Tilston E, Harrad S, et al. 2012. *In vitro* assessment of the bioaccessibility of brominated flame retardants in indoor dust using a colon extended model of the human gastrointestinal tract. *J Environ Monit* 14(12):3276-3283.
- Abdelouahab N, Ainmelk Y, Takser L. 2011. Polybrominated diphenyl ethers and sperm quality. *Reprod Toxicol* 31(4):546-550.
- Abdelouahab N, Langlois MF, Lavoie L, et al. 2013. Maternal and cord-blood thyroid hormone levels and exposure to polybrominated diphenyl ethers and polychlorinated biphenyls during early pregnancy. *Am J Epidemiol* 178(5):701-713. 10.1093/aje/kwt141.
- ACGIH. 2014. Threshold limit values for chemical substances and physical agents and biological exposure indices. Cincinnati, OH: American Conference of Governmental Industrial Hygienists.
- Adgent MA, Hoffman K, Goldman BD, et al. 2014. Brominated flame retardants in breast milk and behavioural and cognitive development at 36 months. *Paediatr Perinat Epidemiol* 28(1):48-57.
- Adinolfi M. 1985. The development of the human blood-CSF-brain barrier. *Dev Med Child Neurol* 27:532-537.
- Adlercreutz H. 1995. Phytoestrogens: Epidemiology and a possible role in cancer protection. *Environ Health Perspect Suppl* 103(7):103-112.
- Agay-Shay K, Martinez D, Valvi D, et al. 2015. Exposure to endocrine-disrupting chemicals during pregnancy and weight at 7 years of age: A multi-pollutant approach. *Environ Health Perspect* 123(10):1030-1037. 10.1289/ehp.1409049.
- Ahn KC, Gee SJ, Tsai HJ, et al. 2009. Immunoassay for monitoring environmental and human exposure to the polybrominated diphenyl ether BDE-47. *Environ Sci Technol* 43(20):7784-7790.
- AIHA. 2014. Current ERPG Values (2014). Fairfax, VA: American Industrial Hygiene Association. <https://www.aiha.org/get-involved/AIHAGuidelineFoundation/EmergencyResponsePlanningGuidelines/Documents/2014%20ERP%20Values.pdf>. September 9, 2014.

9. REFERENCES

- Airaksinen R, Rantakokko P, Eriksson JG, et al. 2011. Association between type 2 diabetes and exposure to persistent organic pollutants. *Diabetes Care* 34(9):1972-1979. 10.2337/dc10-2303.
- Akutsu K, Obana H, Okihashi M, et al. 2001. GC/MS analysis of polybrominated diphenyl in fish collected from the inland sea of Seto, Japan. *Chemosphere* 44:1325-1333.
- Akutsu K, Takatori S, Nozawa S, et al. 2008. Polybrominated diphenyl ethers in human serum and sperm quality. *Bull Environ Contam Toxicol* 80(4):345-350.
- Alaee M, Sergeant DB, Ikonomou MG, et al. 2001. A gas chromatography/high-resolution mass spectrometry (GC/HRMS) method for determination of polybrominated diphenyl ethers in fish. *Chemosphere* 44(6):1489-1895.
- Alaee M, Sergeant DB, Muir DCG, et al. 1999. Distribution of polybrominated diphenyl ethers in the Canadian environment. *Organohalogen Compounds* 40:347-350.
- Albina ML, Alonso V, Linares V, et al. 2010. Effects of exposure to BDE-99 on oxidative status of liver and kidney in adult rats. *Toxicology* 271(1-2):51-56.
- Aleksa K, Carnevale A, Goodyer C, et al. 2012a. Detection of polybrominated biphenyl ethers (PBDEs) in pediatric hair as a tool for determining in utero exposure. *Forensic Sci Int* 218(1-3):37-43.
- Aleksa K, Liesivuori J, Koren G. 2012b. Hair as a biomarker of polybrominated diethyl ethers' exposure in infants, children and adults. *Toxicol Lett* 210(2):198-202.
- Allchin CR, Law RJ, Morris S. 1999. Polybrominated diphenylethers in sediments and biota downstream of potential sources in the UK. *Environ Pollut* 105:197-207.
- Allchin CR, Morris S, Law RJ, et al. 2000. Polybrominated diphenyl ether residues in cormorant (*Phalacrocorax L.*) livers from England, UK. *Organohalogen Compounds* 47:190-193.
- Allen JG, Sumner AL, Nishioka MG, et al. 2013. Air concentrations of PBDEs on in-flight airplanes and assessment of flight crew inhalation exposure. *J Expo Sci Environ Epidemiol* 23(4):337-342.
- Alm H, Kultima K, Scholz B, et al. 2008. Exposure to brominated flame retardant PBDE-99 affects cytoskeletal protein expression in the neonatal mouse cerebral cortex. *Neurotoxicology* 29(4):628-637.
- Alm H, Scholz B, Kultima K, et al. 2010. *In vitro* neurotoxicity of PBDE-99: Immediate and concentration-dependent effects on protein expression in cerebral cortex cells. *J Proteome Res* 9(3):1226-1235.
- Alonso V, Linares V, Belles M, et al. 2010. Effects of BDE-99 on hormone homeostasis and biochemical parameters in adult male rats. *Food Chem Toxicol* 48(8-9):2206-2211.
- Altman PL, Dittmer DS. 1974. In: *Biological handbooks: Biology data book*. Vol. III. 2nd ed. Bethesda, MD: Federation of American Societies for Experimental Biology, 1987-2008, 2041.
- Andersen ME, Krishnan K. 1994. Relating *in vitro* to *in vivo* exposures with physiologically based tissue dosimetry and tissue response models. In: Salem H, ed. *Animal test alternatives: Refinement, reduction, replacement*. New York: Marcel Dekker, Inc., 9-25.

9. REFERENCES

- Andersen ME, Clewell HJ III, Gargas ML, et al. 1987. Physiologically based pharmacokinetics and the risk assessment process for methylene chloride. *Toxicol Appl Pharmacol* 87:185-205.
- Anderson HA, Wolff MS, Fischbein A, et al. 1978. Investigation of the health status of Michigan Chemical Corporation employees. *Environ Health Perspect* 23:187-191.
- Andersson O, Blomkvist G. 1981. Polybrominated aromatic pollutants found in fish in Sweden. *Chemosphere* 10(9):1051-1060.
- Andersson O, Wartanian A. 1992. Levels of polybrominated camphenes (toxaphene), chlordane compounds and polybrominated diphenyl ethers in seals from Swedish waters. *Ambio* 21(8):550-552.
- Andres J, Lambert I, Robertson L, et al. 1983. The comparative biologic and toxic potencies of polychlorinated biphenyls and polybrominated biphenyls. *Toxicol Appl Pharmacol* 70:204-215.
- Ankarberg E, Fredriksson A, Jakobsson E, et al. 2001. Increased susceptibility to adult flame retardants exposure (PBDE 99) in mice neonatally exposed to nicotine. The second international workshop on brominated flame retardants. BFR 2001. Stockholm, Sweden, 233-235.
- Antignac JP, Cariou R, Maume D, et al. 2008. Exposure assessment of fetus and newborn to brominated flame retardants in France: Preliminary data. *Mol Nutr Food Res* 52(2):258-265.
- Antignac JP, Cariou R, Zalko D, et al. 2009. Exposure assessment of French women and their newborn to brominated flame retardants: Determination of tri- to deca- polybromodiphenylethers (PBDE) in maternal adipose tissue, serum, breast milk and cord serum. *Environ Pollut* 157(1):164-173.
- Argus Research Laboratories. 1985a. Initial submission: Embryo/fetal toxicity and teratogenic potential study of Saytex 115 administered orally via gavage to CrI:COBS CD(SD)BR presumed pregnant rats. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8E. OTS0000973.
- Argus Research Laboratories. 1985b. Embryo/fetal toxicity and teratogenic potential study of Saytex 111 administered orally via gavage to pregnant rats (final report-draft) with cover letter dated 050785. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8E. OTS0509725.
- Aschebrook-Kilfoy B, DellaValle CT, Purdue M, et al. 2015. Polybrominated diphenyl ethers and thyroid cancer risk in the Prostate, Colorectal, Lung, and Ovarian Cancer Screening Trial cohort. *Am J Epidemiol* 181(11):883-888. 10.1093/aje/kwu358.
- Asplund L, Athanasiadou M, Sjodin A, et al. 1999a. Organohalogen substances in muscle, egg and blood from healthy Baltic salmon (*Salmo salar*) and Baltic salmon that produced offspring with the M74 syndrome. *Ambio* 28(1):67-76.
- Asplund L, Hornung M, Peterson RE, et al. 1999b. Levels of polybrominated diphenyl ethers (PBDEs) in fish from the Great Lakes and Baltic Sea. *Organohalogen Compounds* 40:351-354.
- Athanasiadou M, Cuadra SN, Marsh G, et al. 2008. Polybrominated diphenyl ethers (PBDEs) and bioaccumulative hydroxylated PBDE metabolites in young humans from Managua, Nicaragua. *Environ Health Perspect* 116(3):400-408.
- Atlas E, Giam CS. 1987. Ambient concentration and precipitation scavenging of atmospheric organic pollutants. *Water Air Soil Pollut* 38:19-36.

9. REFERENCES

- ATSDR. 1989. Decision guide for identifying substance-specific data needs related to toxicological profiles; Notice. Agency for Toxic Substances and Disease Registry, Division of Toxicology. Fed Regist 54(174):37618-37634.
- ATSDR. 1990. Biomarkers of organ damage or dysfunction for the renal, hepatobiliary, and immune systems. Subcommittee on Biomarkers of Organ Damage and Dysfunction. Atlanta, GA: Agency for Toxic Substances and Disease Registry.
- ATSDR. 1994. Toxicological profile for chlorodibenzofurans. Atlanta, GA: U.S. Department of Health and Human Services.
- ATSDR. 1998. Toxicological profile for chlorinated dibenzo-p-dioxins. Atlanta, GA: U.S. Department of Health and Human Services.
- ATSDR. 2000. Toxicological profile for polychlorinated biphenyls. Atlanta, GA: U.S. Department of Health and Human Services.
- ATSDR. 2004. Toxicological profile for polybrominated biphenyls and polybrominated diphenyl ethers. Atlanta, GA: U.S. Department of Health and Human Services.
- ATSDR. 2015. PBDE's. Full SPL data. Substance priority list (SPL) resource page. Agency for Toxic Substances and Disease Registry, Centers for Disease Control and Prevention. <http://www.atsdr.cdc.gov/SPL/resources/index.html>. July 6, 2016.
- Bahn AK, Mills JL, Snyder PJ, et al. 1980. Hypothyroidism in workers exposed to polybrominated biphenyls. N Engl J Med 302(1):31-33.
- Ballschmiter K, Zell M. 1980. Baseline studies of the global pollution: Occurrence of organohalogenes in pristine European and Antarctic aquatic environments. Int J Environ Anal Chem 8:15-35.
- Bannister R, Biegel L, Davis D, et al. 1989. 6-Methyl-1,3,8-trichlorodibenzofuran (MCDF) as a 2,3,7,8-tetrachlorodibenzo-p-dioxin antagonist in C57BL/6 mice. Toxicology 54:139-150.
- Bansal R, Tighe D, Danai A, et al. 2014. Polybrominated diphenyl ether (DE-71) interferes with thyroid hormone action independent of effects on circulating levels of thyroid hormone in male rats. Endocrinology 155(10):4104-4112. 10.1210/en.2014-1154.
- Barnes DG, Dourson M. 1988. Reference dose (RfD): Description and use in health risk assessments. Regul Toxicol Pharmacol 8:471-486.
- Batterman S, Godwin C, Chernyak S, et al. 2010. Brominated flame retardants in offices in Michigan, USA. Environ Int 36(6):548-556.
- Becker RA, Bergfelt DR, Borghoff S, et al. 2012. Interlaboratory study comparison of the 15-day intact adult male rat screening assay: Evaluation of an antithyroid chemical and a negative control chemical. Birth Defects Res B Dev Reprod Toxicol 95(1):63-78.
- Behnisch PA, Hosoe K, Sakai S. 2003. Brominated dioxin-like compounds: *In vitro* assessment in comparison to classical dioxin-like compounds and other polyaromatic compounds. Environ Int 29(6):861-877.

9. REFERENCES

- Belles M, Alonso V, Linares V, et al. 2010. Behavioral effects and oxidative status in brain regions of adult rats exposed to BDE-99. *Toxicol Lett* 194(1-2):1-7.
- Bellinger DC. 2013. Prenatal exposures to environmental chemicals and children's neurodevelopment: An update. *Saf Health Work* 4(1):1-11.
- Berger GS, ed. 1994. Epidemiology of endometriosis. In: *Endometriosis: Advanced management and surgical techniques*. New York, NY: Springer-Verlag, 3-7.
- Bergman A, Athanasiadou M, Wehler EK, et al. 1999. Polybrominated environmental pollutants. Human and wildlife exposures. *Organohalogen Compounds* 43:89-92.
- Beser MI, Beltran J, Yusa V. 2014. Design of experiment approach for the optimization of polybrominated diphenyl ethers determination in fine airborne particulate matter by microwave-assisted extraction and gas chromatography coupled to tandem mass spectrometry. *J Chromatogr A* 1323:1-10.
- Betts K. 2006. PBDEs and PCBs in computers, cars, and homes. *Environ Sci Technol* 40(24):7452.
- BFRIP. 2002. Decabromodiphenyl ether (a.k.a. decabromodiphenyl oxide, DBDPO). Voluntary children's chemical evaluation program (VCCEP). Data Summary. Arlington, VA: American Chemistry Council's Brominated Flame Retardant Industry Panel (BFRIP).
- Bidleman TF. 1988. Atmospheric processes. Wet and dry deposition of organic compounds are controlled by their vapor-particle partitioning. *Environ Sci Technol* 22:361-367.
- Bieniek D, Bahadir M, Korte F. 1989. Formation of heterocyclic hazardous compounds by thermal degradation of organic compounds. *Heterocycles* 28(2):719-722.
- Biesemeier JA, Ariano JM, Banasik M, et al. 2011. Sample characterization: *A priori* to evaluating absorption, distribution, and metabolism. *Toxicology* 287(1-3):160-161; author reply 162-163.
- Biesemeier JA, Beck MJ, Silberberg H, et al. 2010. Effects of dose, administration route, and/or vehicle on decabromodiphenyl ether concentrations in plasma of maternal, fetal, and neonatal rats and in milk of maternal rats. *Drug Metab Dispos* 38(10):1648-1654.
- Bizkarguenaga E, Iparragirre A, Zabaleta I, et al. 2014. Focused ultrasound assisted extraction for the determination of PBDEs in vegetables and amended soil. *Talanta* 119:53-59.
- Blanco J, Mulero M, Domingo JL, et al. 2012. Gestational exposure to BDE-99 produces toxicity through upregulation of CYP isoforms and ROS production in the fetal rat liver. *Toxicol Sci* 127(1):296-302.
- Blanco J, Mulero M, Domingo JL, et al. 2014. Perinatal exposure to BDE-99 causes decreased protein levels of cyclin D1 via GSK3 β activation and increased ROS production in rat pup livers. *Toxicol Sci* 137(2):491-498.
- Blanco J, Mulero M, Heredia L, et al. 2013. Perinatal exposure to BDE-99 causes learning disorders and decreases serum thyroid hormone levels and BDNF gene expression in hippocampus in rat offspring. *Toxicology* 308:122-128.

9. REFERENCES

- Blanco J, Mulero M, Lopez M, et al. 2011. BDE-99 deregulates BDNF, Bcl-2 and the mRNA expression of thyroid receptor isoforms in rat cerebellar granular neurons. *Toxicology* 290:305-311.
- Blay P, Nilsson C, Owman C, et al. 1993. Transthyretin expression in the rat brain: Effect of thyroid functional state and role in thyroxine transport. *Brain Res* 632(1-2):114-120.
- Bloom M, Spliethoff H, Vena J, et al. 2008. Environmental exposure to PBDEs and thyroid function among New York anglers. *Environ Toxicol Pharmacol* 25(3):386-392.
- Bloom MS, Jansing RL, Kannan K, et al. 2014. Thyroid hormones are associated with exposure to persistent organic pollutants in aging residents of upper Hudson River communities. *Int J Hyg Environ Health* 217(4-5):473-482.
- Bocio A, Llobet JM, Domingo JL, et al. 2003. Polybrominated diphenyl ethers (PBDEs) in foodstuffs: Human exposure through the diet. *J Agric Food Chem* 7(51):3191-3195.
- Bondy GS, Gaertner D, Cherry W, et al. 2011. Brominated diphenyl ether (BDE) levels in liver, adipose, and milk from adult and juvenile rats exposed by gavage to the DE-71 technical mixture. *Environ Toxicol* 26(6):677-690.
- Bondy GS, Lefebvre DE, Aziz S, et al. 2013. Toxicologic and immunologic effects of perinatal exposure to the brominated diphenyl ether (BDE) mixture DE-71 in the Sprague-Dawley rat. *Environ Toxicol* 28(4):215-228.
- Booij K, Zegers BN, Boon JP. 2000. Levels of some polybrominated diphenyl ether (PBDE) flame retardants along the Dutch coast as derived from their accumulation in SPMDs and blue mussels (*Mytilus edulis*). *Organohalogen Compounds* 47:89-92.
- Bowers WJ, Wall PM, Nakai JS, et al. 2015. Behavioral and thyroid effects of in utero and lactational exposure of Sprague-Dawley rats to the polybrominated diphenyl ether mixture DE71. *Neurotoxicol Teratol* 52(Pt B):127-142. 10.1016/j.ntt.2015.08.002.
- Boyages SC. 2000. The neuromuscular system and brain in hypothyroidism. In: Braverman LE, Utiger RD, eds. *Werner and Ingbar's the thyroid*. Philadelphia, PA: Lippincott Williams & Wilkins, 803-810.
- Bradman A, Castorina R, Gaspar F, et al. 2014. Flame retardant exposures in California early childhood education environments. *Chemosphere* 116:61-66. 10.1016/j.chemosphere.2014.02.072.
- Bradner JM, Suragh TA, Caudle WM. 2013. Alterations to the circuitry of the frontal cortex following exposure to the polybrominated diphenyl ether mixture, DE-71. *Toxicology* 312:48-55.
- Braekvelt E, Tittlemier SA, Tomy GT. 2003. Direct measurement of octanol-water partition coefficients of some environmentally relevant brominated diphenyl ether congeners. *Chemosphere* 51(7):563-567.
- Branchi I, Alleva E, Costa LG. 2001. A preliminary characterization of behavioural alterations following perinatal exposure to a polybrominated diphenylether (PBDE 99). The second international workshop on brominated flame retardants. BFR 2001. Stockholm, Sweden, 75.
- Branchi I, Alleva E, Costa LG. 2002. Effects of perinatal exposure to a polybrominated diphenyl ether (PBDE 99) on mouse neurobehavioural development. *Neurotoxicology* 23(3):375-384.

9. REFERENCES

- Branchi I, Capone F, Vitalone A, et al. 2005. Early developmental exposure to BDE 99 or Aroclor 1254 affects neurobehavioural profile: Interference from the administration route. *Neurotoxicology* 26(2):183-192.
- Braun JM, Kalkbrenner AE, Just AC, et al. 2014. Gestational exposure to endocrine-disrupting chemicals and reciprocal social, repetitive, and stereotypic behaviors in 4- and 5-year-old children: The HOME study. *Environ Health Perspect* 122(5):513-520. 10.1289/ehp.1307261.
- Breivik K, Sweetman A, Pacyna JM, et al. 2002. Towards a global historical emission inventory for selected PCB congeners— a mass balance approach. 2. Emissions. *Sci Total Environ* 290:199-224.
- Brenner KS, Knies H. 1990. Formation of polybrominated dibenzofurans (PBDF's) and dioxins (PBDD's) during extrusion production of a polybutyleneterephthalate (PBTP)/glass fiber resin blended with decabromodiphenylether (DBDPE)/Sb₂O₃: Product and workplace analysis. *Organohalogen Compounds* 2:319-324.
- Brenner KS, Knies H. 1993. Workplace of PBDFs and PBDDs during extrusion production and injection modeling of a polybutyleneterephthalate (PBTP) glass fiber/tetrabromobisphenol A carbonate oligomer (BC52*)/Sb₂O₃-resin; part II. *Chemosphere* 26(11):1953-1963.
- Breslin WJ, Kirk HD, Zimmer MA. 1989. Teratogenic evaluation of a polybromodiphenyl oxide mixture in New Zealand white rabbits following oral exposure. *Fundam Appl Toxicol* 12(1):151-157.
- Brilliant LB, Van Amburg G, Isbister J, et al. 1978. Breast-milk monitoring to measure Michigan's contamination with polybrominated biphenyls. *Lancet* Sept:643-646.
- British Industrial Biological Research Association. 1977. The acute oral toxicity of pentabromodiphenylether to rats. Submitted to U.S. Environmental Protection Agency under TSCA Section 8D. OTS0522287.
- Brown GG, Preisman RC, Anderson MD, et al. 1981. Memory performance of chemical workers exposed to polybrominated biphenyls. *Science* 212:1413-1415.
- Bruchajzer E. 2011. Porphyrigenic effect of pentabromodiphenyl ether after repeated administration to rats. *Arch Toxicol* 85(8):965-974.
- Bruchajzer E, Frydrych B, Sporny S, et al. 2010. Toxicity of penta- and decabromodiphenyl ethers after repeated administration to rats: A comparative study. *Arch Toxicol* 84(4):287-299.
- Bruchajzer E, Frydrych B, Sporny S, et al. 2011. The effect of short-term intoxication of rats with pentabromodiphenyl ether (in mixture mimic commercial products). *Hum Exp Toxicol* 30(5):363-378.
- Bruchajzer E, Frydrych B, Szymanska JA. 2012. Octabromodiphenyl ether - porphyrigenicity after repeated administration to rats. *Int J Occup Med Environ Health* 25(4):392-403.
- Buck Louis GM, Sundaram R, Schisterman EF, et al. 2013. Persistent environmental pollutants and couple fecundity: The LIFE study. *Environ Health Perspect* 121(2):231-236.

9. REFERENCES

- Bull K, Basu N, Zhang S, et al. 2007. Dietary and *in utero* exposure to a pentabrominated diphenyl ether mixture did not affect cholinergic parameters in the cerebral cortex of ranch mink (*Mustela vison*). *Toxicol Sci* 96(1):115-122.
- Buratovic S, Viberg H, Fredriksson A, et al. 2014. Developmental exposure to the polybrominated diphenyl ether PBDE 209: Neurobehavioural and neuroprotein analysis in adult male and female mice. *Environ Toxicol Pharmacol* 38(2):570-585. 10.1016/j.etap.2014.08.010.
- Burreau S, Axelman J, Brogman D, et al. 1997. Dietary uptake in pike (*Esox incius*) of some polychlorinated biphenyls, polychlorinated naphthalenes and polybrominated diphenyl ethers administered in natural diet. *Environ Toxicol Chem* 16:2508-2513.
- Burreau S, Broman D, Zebuhr Y. 1999. Biomagnification quantification of PBDEs in fish using stable nitrogen isotopes. *Organohalogen Compounds* 40:363-366.
- Burreau S, Zeb HR, Ishaq R, et al. 2000. Comparison of biomagnification of PBDEs in food chains from the Baltic Sea and the North Atlantic Sea. *Organohalogen Compounds* 47:253-255.
- Burse VW, Needham LL, Liddle JA, et al. 1980. Interlaboratory comparison for polybrominated biphenyls in human serum. *J Anal Toxicol* 4:22-26.
- Buser HR. 1986. Polybrominated dibenzofurans and dibenzo-p-dioxins: Thermal reaction products of polybrominated diphenyl ether flame retardants. *Environ Sci Technol* 20:404-408.
- Butt CM, Miranda ML, Stapleton HM. 2016. Development of an analytical method to quantify PBDEs, OH-BDEs, HBCDs, 2,4,6-TBP, EH-TBB, and BEH-TEBP in human serum. *Anal Bioanal Chem* 408(10):2449-2459. 10.1007/s00216-016-9340-3.
- Byun HM, Benachour N, Zalko D, et al. 2015. Epigenetic effects of low perinatal doses of flame retardant BDE-47 on mitochondrial and nuclear genes in rat offspring. *Toxicology* 328:152-159. 10.1016/j.tox.2014.12.019.
- Cai Y, Zhang W, Hu J, et al. 2011. Characterization of maternal transfer of decabromodiphenyl ether (BDE-209) administered to pregnant Sprague-Dawley rats. *Reprod Toxicol* 31(1):106-110.
- Canton RF, Letcher R, Sanderson T, et al. 2003. Effects of brominated flame retardants on activity of the steroidogenic enzyme aromatase (CYP19) in H295R human adrenocortical carcinoma cells in culture. *Organohalogen Compounds* 61:104-106.
- Canton RF, Sanderson JT, Letcher RJ, et al. 2005. Inhibition and induction of aromatase (CYP19) activity by brominated flame retardants in H295R human adrenocortical carcinoma cells. *Toxicol Sci* 88(2):447-455.
- Canton RF, Scholten DE, Marsh G, et al. 2008. Inhibition of human placental aromatase activity by hydroxylated polybrominated diphenyl ethers (OH-PBDEs). *Toxicol Appl Pharmacol* 227(1):68-75.
- Capen CC. 1997. Mechanistic data and risk assessment of selected toxic points of the thyroid gland. *Toxicol Pathol* 25:39-48.
- Caravati EM, Mcguigan MA, Whyte IM, et al. 2004. Polyhalogenated biphenyls. In: Dart RC, ed. *Medical toxicology*. 3rd ed. Philadelphia, PA: Lippincott Williams & Wilkins, 1342-1343.

9. REFERENCES

- Carlson GP. 1980a. Induction of xenobiotic metabolism in rats by brominated diphenyl ethers administered for 90 days. *Toxicol Lett* 6:207-212.
- Carlson GP. 1980b. Induction of xenobiotic metabolism in rats by short-term administration of brominated phenyl ethers. *Toxicol Lett* 5(1):19-26.
- Carmichael SL, Herring AH, Sjodin A, et al. 2010. Hypospadias and halogenated organic pollutant levels in maternal mid-pregnancy serum samples. *Chemosphere* 80(6):641-646.
- Carrizo D, Grimalt JO, Ribas-Fito N, et al. 2007. Influence of breastfeeding in the accumulation of polybromodiphenyl ethers during the first years of child growth. *Environ Sci Technol* 41(14):4907-4912.
- CDC. 2015. Fourth national report on human exposure to environmental chemical. Updated tables, February 2015. Atlanta, GA: Centers for Disease Control and Prevention, U.S. Department of Health and Human Services.
http://www.cdc.gov/biomonitoring/pdf/FourthReport_UpdatedTables_Feb2015.pdf. March 10, 2015
- Ceccatelli R, Faass O, Schlumpf M, et al. 2006. Gene expression and estrogen sensitivity in rat uterus after developmental exposure to the polybrominated diphenylether PBDE 99 and PCB. *Toxicology* 220(2-3):104-116.
- Chanda JJ, Anderson HA, Glamb RW, et al. 1982. Cutaneous effects of exposure to polybrominated biphenyls (PBBs): The Michigan PBB incident. *Environ Res* 29:97-108.
- Chao HR, Shy CG, Wang SL, et al. 2010. Impact of non-occupational exposure to polybrominated diphenyl ethers on menstruation characteristics of reproductive-age females. *Environ Int* 36(7):728-735.
- Chao HR, Tsou TC, Huang HL, et al. 2011. Levels of breast milk PBDEs from southern Taiwan and their potential impact on neurodevelopment. *Pediatr Res* 70(6):596-600.
- Chao HR, Wang SL, Lee WJ, et al. 2007. Levels of polybrominated diphenyl ethers (PBDEs) in breast milk from central Taiwan and their relation to infant birth outcome and maternal menstruation effects. *Environ Int* 33(2):239-245.
- Charles River Laboratories. 1998. Baseline hematology and clinical chemistry values for Charles River Wistar rats- (CLR:(WI)BR) as a function of sex and age. Wilmington, MA: Charles River Laboratories.
- Chen G, Bunce NJ. 2001. PBDE congeners as Ah receptor agonist and antagonists. *Organohalogen Compounds* 53:353-356.
- Chen A, Chung E, DeFranco EA, et al. 2011. Serum PBDEs and age at menarche in adolescent girls: Analysis of the National Health and Nutrition Examination Survey 2003-2004. *Environ Res* 111(6):831-837.
- Chen A, Park JS, Linderholm L, et al. 2013. Hydroxylated polybrominated diphenyl ethers in paired maternal and cord sera. *Environ Sci Technol* 47(8):3902-3908.
- Chen A, Yolton K, Rauch SA, et al. 2014. Prenatal polybrominated diphenyl ether exposures and neurodevelopment in U.S. children through 5 years of age: The HOME study. *Environ Health Perspect* 122(8):856-862. 10.1289/ehp.1307562.

9. REFERENCES

- Chen G, Konstantinov AD, Chittim BG, et al. 2001. Synthesis of polybrominated diphenyl ethers and their capacity to induce CYPIA by the Ah receptor mediated pathway. *Environ Sci Technol* 35(18):3749-3756.
- Chen J, Liufu C, Sun W, et al. 2010. Assessment of the neurotoxic mechanisms of decabrominated diphenyl ether (PBDE-209) in primary cultured neonatal rat hippocampal neurons includes alterations in second messenger signaling and oxidative stress. *Toxicol Lett* 192(3):431-439.
- Chen L, Wang C, Cui C, et al. 2015. Prenatal exposure to polybrominated diphenyl ethers and birth outcomes. *Environ Pollut* 206:32-37. 10.1016/j.envpol.2015.06.019.
- Chen LJ, Lebetkin EH, Sanders JM, et al. 2006. Metabolism and disposition of 2,2',4,4',5-pentabromodiphenyl ether (BDE99) following a single or repeated administration to rats or mice. *Xenobiotica* 36(6):515-534.
- Chen Q, Yu L, Yang L, et al. 2012. Bioconcentration and metabolism of decabromodiphenyl ether (BDE-209) result in thyroid endocrine disruption in zebrafish larvae. *Aquat Toxicol* 110-111:141-148.
- Chen SJ, Ma YJ, Wang J, et al. 2009. Brominated flame retardants in children's toys: Concentration, composition, and children's exposure and risk assessment. *Environ Sci Technol* 43(11):4200-4206.
- Cheng J, Gu J, Ma J, et al. 2009. Neurobehavioural effects, redox responses and tissue distribution in rat offspring developmental exposure to BDE-99. *Chemosphere* 75(7):963-968.
- Cheng SW, Randall K, Kotchevar AT. 2008. *In vitro* metabolism studies of polybrominated diphenyl ethers using rat and human liver microsomes. *Am J Biochem Biotechnol* 4(3):295-303.
- Chessells M, Hawker DW, Connell DW. 1992. Influence of solubility in lipid on bioconcentration of hydrophobic compounds. *Ecotoxicol Environ Saf* 23:260-273.
- Chevrier C, Warembourg C, Le Maner-Idrissi G, et al. 2016. Childhood exposure to polybrominated diphenyl ethers and neurodevelopment at six years of age. *Neurotoxicology* 54:81-88. 10.1016/j.neuro.2016.03.002.
- Chevrier J, Harley KG, Bradman A, et al. 2010. Polybrominated diphenyl ether (PBDE) flame retardants and thyroid hormone during pregnancy. *Environ Health Perspect* 118(10):1444-1449.
- Chevrier J, Harley KG, Bradman A, et al. 2011. Prenatal exposure to polybrominated diphenyl ether flame retardants and neonatal thyroid-stimulating hormone levels in the CHAMACOS study. *Am J Epidemiol* 174(10):1166-1174.
- Chi Y, Xia H, Su M, et al. 2011. Metabonomic phenotyping reveals an embryotoxicity of decabrominated diphenyl ether in mice. *Chem Res Toxicol* 24(11):1976-1983.
- Choi J-W, Fujimaki S, Kitamura K, et al. 2003. Historical trends of PBDD/Fs, PBDEs, PCDD/Fs and dioxin-like PCBs in sediment cores from Tokyo Bay. *Organohalogen Compounds* 61:119-122.
- Christen V, Crettaz P, Oberli-Schrammli A, et al. 2010. Some flame retardants and the antimicrobials triclosan and triclocarban enhance the androgenic activity *in vitro*. *Chemosphere* 81(10):1245-1252.

9. REFERENCES

- Christensen JH, Platz J. 2001. Screening of polybrominated diphenyl ethers in blue mussels, marine and freshwater sediments in Denmark. *J Environ Monit* 3(5):543-547.
- Clewell HJ III, Andersen ME. 1985. Risk assessment extrapolations and physiological modeling. *Toxicol Ind Health* 1(4):111-131.
- Coburn CG, Curras-Collazo MC, Kodavanti PR. 2007. Polybrominated diphenyl ethers and *ortho*-substituted polychlorinated biphenyls as neuroendocrine disruptors of vasopressin release: Effects during physiological activation *in vitro* and structure-activity relationships. *Toxicol Sci* 98(1):178-186.
- Colborn T, vom Saal F, Soto A. 1993. Developmental effects of endocrine-disrupting chemicals in wildlife and humans. *Environ Health Perspect* 101(5):378-384.
- Costa LG, Aschner M, Vitalone A, et al. 2004. Developmental neuropathology of environmental agents. *Annu Rev Pharmacol Toxicol* 44:87-110.
- Costa LG, Pellacani C, Dao K, et al. 2015. The brominated flame retardant BDE-47 causes oxidative stress and apoptotic cell death *in vitro* and *in vivo* in mice. *Neurotoxicology* 48:68-76. 10.1016/j.neuro.2015.03.008.
- Costa LG, Tagliaferri S, Roque PJ, et al. 2016. Role of glutamate receptors in tetrabrominated diphenyl ether (BDE-47) neurotoxicity in mouse cerebellar granule neurons. *Toxicol Lett* 241:159-166. 10.1016/j.toxlet.2015.11.026.
- Covaci A, de Boer J, Ryan JJ, et al. 2002. Determination of polybrominated diphenyl ethers and polychlorinated biphenyls in human adipose tissues by large-volume injection-narrow-bore capillary gas chromatography/electron impact low-resolution mass spectrometry. *Anal Chem* 74(4):790-798.
- Covaci A, Voorspoels S, de Boer J. 2003. Determination of brominated flame retardants with emphasis on polybrominated diphenyl ethers (PBDEs) in environmental and human samples- a review. *Environ Int* 29:735-756.
- Cowell WJ, Lederman SA, Sjodin A, et al. 2015. Prenatal exposure to polybrominated diphenyl ethers and child attention problems at 3-7 years. *Neurotoxicol Teratol* 52(Pt B):143-150. 10.1016/j.ntt.2015.08.009.
- Cramer PH, Ayling RE, Thornburg KR, et al. 1990. Evaluation of an analytical method for the determination of polybrominated dibenzo-p-dioxins/dibenzofurans (PBDD/PBDF) in human adipose. *Chemosphere* 20(7-9):821-827.
- Crisp TM, Clegg ED, Cooper RL, et al. 1998. Environmental endocrine disruption: An effects assessment and analysis. *Environ Health Perspect* 106(1):11-56.
- Curcic M, Durgo K, Kopjar N, et al. 2014. Cadmium and decabrominated diphenyl ether mixture: *In vitro* evaluation of cytotoxic, prooxidative and genotoxic effects. *Environ Toxicol Pharmacol* 38(2):663-671. 10.1016/j.etap.2014.07.021.
- Dang VH, Choi KC, Jeung EB. 2007. Tetrabromodiphenyl ether (BDE 47) evokes estrogenicity and calbindin-D9k expression through an estrogen receptor-mediated pathway in the uterus of immature rats. *Toxicol Sci* 97(2):504-511.

9. REFERENCES

- Daniels JL, Pan IJ, Jones R, et al. 2010. Individual characteristics associated with PBDE levels in U.S. human milk samples. *Environ Health Perspect* 118(1):155-160.
- Danish EPA. 1999. Brominated flame retardants. Danish Environmental Protection Agency. <http://www.statensnet.dk/pligtarkiv/fremvis.pl?vaerkid=3548&reprid=0&filid=32&iarkiv=1>. October 15, 2014.
- Danon-Schaffer MN, Mahecha-Botero A, Grace JR, et al. 2013. Mass balance evaluation of polybrominated diphenyl ethers in landfill leachate and potential for transfer from e-waste. *Sci Total Environ* 461-462:290-301.
- Darnerud PO, Sinjari T. 1996. Effects of polybrominated diphenyl ethers (PBDEs) and polybrominated biphenyls (PCBs) on thyroxine and TSH blood levels in rats and mice. *Organohalogen Compounds* 29:316-319.
- Darnerud PO, Thuvander A. 1998. Studies on immunological effects of polybrominated diphenyl ether (PBDE) and polychlorinated biphenyl (PCB) exposure in rats and mice. *Organohalogen Compounds* 35:415-418.
- Darnerud PO, Atuma S, Aune M, et al. 1998. Polybrominated diphenyl ethers (PBDEs) in breast milk from primiparous women in Uppsala County, Sweden. *Organohalogen Compounds* 35:411-414.
- Darnerud PO, Eriksen GS, Johannesson T, et al. 2001. Polybrominated diphenyl ethers: Occurrence, dietary exposure, and toxicology. *Environ Health Perspect* 109:49-64.
- Daso AP, Fatoki OS, Odendaal JP, et al. 2013. Polybrominated diphenyl ethers (PBDEs) and 2,2',4,4',5,5'-hexabromobiphenyl (BB-153) in landfill leachate in Cape Town, South Africa. *Environ Monit Assess* 185(1):431-439.
- Daston GP, Gooch JW, Breslin WJ, et al. 1997. Environmental estrogens and reproductive health: A discussion of the human and environmental data. *Reprod Toxicol* 11(4):465-481.
- Daubie S, Bisson JF, Lalonde R, et al. 2011. Neurobehavioral and physiological effects of low doses of polybrominated diphenyl ether (PBDE)-99 in male adult rats. *Toxicol Lett* 204(1):57-63.
- de Boer J. 1989. Organochlorine compounds and bromodiphenylethers in livers of Atlantic cod (*Gadus morhua*) from the North Sea, 1977-1987. *Chemosphere* 18(11/12):2131-2140.
- de Boer J. 1990. Brominated diphenyl ethers in Dutch freshwater and marine fish. *Organohalogen Compounds* 2:315-318.
- de Boer J. 2000. First worldwide interlaboratory study on polybrominated diphenyl ethers (PBDEs). *Organohalogen Compounds* 45:118-121.
- de Boer J, Cofino WP. 2002. First world-wide interlaboratory study on polybrominated diphenyl ethers (PBDEs). *Chemosphere* 46(5):625-633.
- de Boer J, de Boer K, Boon JP. 2000a. Polybrominated biphenyls and diphenylethers. In: Paasivirta J, ed. *Handbook of environmental chemistry Vol. 3 part K*. Berlin Heidelberg: Springer-Verlag, 61-95.

9. REFERENCES

- de Boer J, van der Horst A, Wester PG. 2000b. PBDEs and PBBs in suspended particulate matter, sediments, sewage treatment plant in- and effluent and biota from The Netherlands. *Organohalogen Compounds* 47:85-88.
- de Boer J, Wester PG, Klamer HJ, et al. 1998a. Do flame retardants threaten ocean life? *Nature* 394(6688):28-29.
- de Boer J, Wester PG, Pastor RD, et al. 1998b. Polybrominated biphenyls and diphenylethers in sperm whales and other marine mammals. A new threat to ocean life? *Organohalogen Compounds* 35:383-386.
- de Cock M, Maas YG, van de Bor M. 2012. Does perinatal exposure to endocrine disruptors induce autism spectrum and attention deficit hyperactivity disorders? Review. *Acta Paediatrica* 101(8):811-818.
- de Kok JJ, De Kok A, Brinkman UA, et al. 1977. Analysis of polybrominated biphenyls. *J Chromatogr* 142:367-383.
- de Wit CA. 2000. Levels and trends of BFRs in the European environment. The second international workshop on brominated flame retardants. BFR 2001. Stockholm, Sweden, 135-138.
- de Wit CA. 2002. An overview of brominated flame retardants in the environment. *Chemosphere* 46(5):583-624.
- Dickhut RM, Cincinelli A, Cochran M, et al. 2012. Aerosol-mediated transport and deposition of brominated diphenyl ethers to Antarctica. *Environ Sci Technol* 46(6):3135-3140.
- Ding G, Yu J, Cui C, et al. 2015. Association between prenatal exposure to polybrominated diphenyl ethers and young children's neurodevelopment in China. *Environ Res* 142:104-111. 10.1016/j.envres.2015.06.008.
- Dingemans MM, de Groot A, van Kleef RG, et al. 2008. Hydroxylation increases the neurotoxic potential of BDE-47 to affect exocytosis and calcium homeostasis in PC12 cells. *Environ Health Perspect* 116(5):637-643.
- Dingemans MM, Heusinkveld HJ, Bergman A, et al. 2010a. Bromination pattern of hydroxylated metabolites of BDE-47 affects their potency to release calcium from intracellular stores in PC12 cells. *Environ Health Perspect* 118(4):519-525.
- Dingemans MM, Ramakers GM, Gardoni F, et al. 2007. Neonatal exposure to brominated flame retardant BDE-47 reduces long-term potentiation and postsynaptic protein levels in mouse hippocampus. *Environ Health Perspect* 115(6):865-870.
- Dingemans MM, van den Berg M, Bergman A, et al. 2010b. Calcium-related processes involved in the inhibition of depolarization-evoked calcium increase by hydroxylated PBDEs in PC12 cells. *Toxicol Sci* 114(2):302-309.
- Dingemans MM, van den Berg M, Westerink RH. 2011. Neurotoxicity of brominated flame retardants: (In)direct effects of parent and hydroxylated polybrominated diphenyl ethers on the (developing) nervous system. *Environ Health Perspect* 119(7):900-907.

9. REFERENCES

- Dishaw LV, Powers CM, Ryde IT, et al. 2011. Is the pentaBDE replacement, tris (1,3-dichloro-2-propyl) phosphate (TDCPP), a developmental neurotoxicant? Studies in PC12 cells. *Toxicol Appl Pharmacol* 256(3):281-289.
- Dodder NG, Strandberg B, Hites RA. 2000. Concentrations and spatial variations of polybrominated diphenyl ethers in fish and air from the northeastern United States. *Organohalogen Compounds* 47:69-72.
- Dodder NG, Strandberg B, Hites RA. 2002. Concentrations and spatial variations of polybrominated diphenyl ethers and several organochlorine compounds in fishes from the northeastern United States. *Environ Sci Technol* 36(2):146-151.
- DOE. 2012. Table 3: Protective Action Criteria (PAC) Rev 27 based on applicable 60-minute AEGLs, ERPGs, or TEELs. The chemicals are listed by CASRN. February 2012. Oak Ridge, TN: U.S. Department of Energy and Subcommittee on Consequence Assessment and Protective Actions (SCAPA). <http://www.atlant.com/DOE/teels/teel/Table3.pdf>. September 9, 2014.
- Domino EF, Wright DD, Domino SE. 1980. GC-EC analysis of polybrominated biphenyl constituents of Firemaster FF-1 using tetrabromobiphenyl as an internal standard. *J Anal Toxicol* 4:299-304.
- Donauer S, Chen A, Xu Y, et al. 2015. Prenatal exposure to polybrominated diphenyl ethers and polyfluoroalkyl chemicals and infant neurobehavior. *J Pediatr* 166(3):736-742. 10.1016/j.jpeds.2014.11.021.
- Dong Y, Li L, Bie P, et al. 2014. Polybrominated diphenyl ethers in farmland soils: Source characterization, deposition contribution and apportionment. *Sci Total Environ* 466-467:524-532.
- Donnelly J, Grange AH, Nunn NJ, et al. 1987. Analysis of thermoplastic resins for brominated dibenzofurans. *Biomed Environ Mass Spectrom* 18(10):884-896.
- Dow Chemical Co. 1975. Results of a reproduction study in rats maintained on diets containing decabromodiphenyl oxide. Dow Chemical Company. Submitted to U.S. Environmental Protection Agency under TSCA Section 8D. OTS0522252.
- Dow Chemical Co. 1985. Decabromodiphenyloxide: A summary of an oral teratology study in Sprague-Dawley rats with cover letter dated 030890. Dow Chemical Company. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. OTS0522284.
- Dreiem A, Okoniewski RJ, Brosch KO, et al. 2010. Polychlorinated biphenyls and polybrominated diphenyl ethers alter striatal dopamine neurochemistry in synaptosomes from developing rats in an additive manner. *Toxicol Sci* 118(1):150-159.
- Driscoll LL, Gibson AM, Hieb A. 2009. Chronic postnatal DE-71 exposure: Effects on learning, attention and thyroxine levels. *Neurotoxicol Teratol* 31(2):76-84.
- Driscoll LL, Kaplan J, Bucuvalas E, et al. 2012. Acute postnatal exposure to the pentaBDE commercial mixture DE-71 at 5 or 15 mg/kg/day does not produce learning or attention deficits in rats. *Neurotoxicol Teratol* 34(1):20-26.
- Dufault C, Poles G, Driscoll LL. 2005. Brief postnatal PBDE exposure alters learning and the cholinergic modulation of attention in rats. *Toxicol Sci* 88(1):172-180.

9. REFERENCES

- Dumler R, Lenoir D, Thoma H, et al. 1990. Thermal formation of polybrominated dibenzofurans and dioxins from decabromodiphenyl ether flame retardants influence of antimony-III oxide and the polymer matrix. *Chemosphere* 20(10-12):1867-1874.
- Dumler R, Thoma H, Lenoir D, et al. 1989. PBDF and PBDD from the combustion of bromine containing flame retarded polymers: A survey. *Chemosphere* 19(12):2023-2031.
- Dunnick JK, Brix A, Cunny H, et al. 2012. Characterization of polybrominated diphenyl ether toxicity in Wistar Han rats and use of liver microarray data for predicting disease susceptibilities. *Toxicol Pathol* 40(1):93-106.
- Easton MDL, Luszniak D, Von der Geest E. 2002. Preliminary examination of contaminant loadings in farmed salmon, wild salmon and commercial salmon feed. *Chemosphere* 46(7):1053-1074.
- Ebert J, Lorenze W, Bahadir M. 1999. Optimization of the analytical performance of polybrominated dibenzo-p-dioxins and dibenzofurans (PBDD-F). *Chemosphere* 39(6):977-986.
- EC. 2014. Report from the commission on the working of committees during 2008. Brussels: European Commission.
- Eggesbo M, Thomsen C, Jorgensen JV, et al. 2011. Associations between brominated flame retardants in human milk and thyroid-stimulating hormone (TSH) in neonates. *Environ Res* 111(6):737-743.
- Eguchi A, Nomiya K, Minh Tue N, et al. 2015. Residue profiles of organohalogen compounds in human serum from e-waste recycling sites in North Vietnam: Association with thyroid hormone levels. *Environ Res* 137:440-449. 10.1016/j.envres.2015.01.007.
- Eisenreich SJ, Looney BB, Thornton JD. 1981. Airborne organic contaminants in the Great Lakes ecosystem. *Environ Sci Technol* 15(1):30-38.
- Ek CJ, Dziegielewska KM, Habgood MD, et al. 2012. Barriers in the developing brain and neurotoxicology. *Neurotoxicology* 33(3):586-604.
- el Dareer SM, Kalin JR, Tillery KF, et al. 1987. Disposition of decabromobiphenyl ether in rats dosed intravenously or by feeding. *J Toxicol Environ Health* 22(4):405-415.
- Eljarrat E, Lacorte S, Barcelo D. 2002. Optimization of congener-specific analysis of 40 polybrominated diphenyl ethers by gas chromatography/mass spectrometry. *J Mass Spectrom* 37(1):76-84.
- Ellis-Hutchings RG, Cherr GN, Hanna LA, et al. 2006. Polybrominated diphenyl ether (PBDE)-induced alterations in vitamin A and thyroid hormone concentrations in the rat during lactation and early postnatal development. *Toxicol Appl Pharmacol* 215(2):135-145.
- Ellis-Hutchings RG, Cherr GN, Hanna LA, et al. 2009. The effects of marginal maternal vitamin A status on penta-brominated diphenyl ether mixture-induced alterations in maternal and conceptual vitamin A and fetal development in the Sprague Dawley rat. *Birth Defects Res B Dev Reprod Toxicol* 86(1):48-57.
- Emond C, Raymer JH, Studabaker WB, et al. 2010. A physiologically based pharmacokinetic model for developmental exposure to BDE-47 in rats. *Toxicol Appl Pharmacol* 242(3):290-298.

9. REFERENCES

Emond C, Sanders JM, Wikoff D, et al. 2013. Proposed mechanistic description of dose-dependent BDE-47 urinary elimination in mice using a physiologically based pharmacokinetic model. *Toxicol Appl Pharmacol* 273(2):335-344.

ENVIRON. 2003a. Voluntary children's chemical evaluation program pilot. Tier I assessment of the potential health risks to children associated with exposure to the commercial octabromodiphenyl ether product. CAS No. 32536-52-0. Emeryville, CA: ENVIRON Int. Corp.

ENVIRON. 2003b. Voluntary children's chemical evaluation program pilot. Tier I assessment of the potential health risks to children associated with exposure to the commercial pentabromodiphenyl ether product. CAS No. 32534-81-9. Emeryville, CA: ENVIRON Int. Corp.

EPA. 1990. Interim methods for development of inhalation reference concentrations. Washington, DC: U.S. Environmental Protection Agency, Office of Health and Environmental Assessment, Office of Research and Development, Environmental Criteria and Assessment Office. EPA600/8-90066A.

EPA. 1995. Toxic chemical release inventory. Reporting form R and instructions. Washington, DC: U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics. EPA745K95051.

EPA. 1997. Special report on environmental endocrine disruption: An effects assessment and analysis. Washington, DC: U.S. Environmental Protection Agency, Risk Assessment Forum. EPA630R96012.

EPA. 1998a. Appendix A. Chemical substances subject to proposed or final TSCA rules or orders. In: Instructions for reporting for the 1998 partial updating of the TSCA chemical inventory data base. Washington, DC: U.S. Environmental Protection Agency, Office of Prevention Pesticides and Toxic Substances, A-1 to A-10. <http://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=700004YF.txt>. September 24, 2014.

EPA. 1998b. Notice of availability of draft RCRA waste minimization PBT chemical list. U.S. Environmental Protection Agency. Federal Register 63 FR 60332. <http://www.gpo.gov/fdsys/pkg/FR-1998-11-09/pdf/98-29952.pdf>. September 10, 2014.

EPA. 2005. Toxic chemical release inventory reporting forms and instructions: Revised 2004 version. Section 313 of the Emergency Planning and Community Right-to-Know Act (Title III of the Superfund Amendments and Reauthorization Act of 1986). U.S. Environmental Protection Agency, Office of Environmental Information. EPA260B05001.

EPA. 2006. List of lists. Consolidated list of chemicals subject to the Emergency Planning and Community Right-to-Know Act (EPCRA) and Section 112(r) of the Clean Air Act. Washington, D.C.: U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response, 1-38. EPA550B01003. http://www.epa.gov/emergencies/docs/chem/title3_Oct_2006.pdf. September 24, 2014.

EPA. 2008a. Toxicological review of 2,2',4,4',5-pentabromodiphenyl ether (BDE-99) (CAS No. 60348-60-9). In support of summary information on the Integrated Risk Information System (IRIS). Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/iris/toxreviews/1008tr.pdf>. December 19, 2014.

EPA. 2008b. Toxicological review of decabromodiphenyl ether (BDE-209) (CAS No. 1163-19-5). In support of summary information on the Integrated Risk Information System (IRIS). Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/iris/toxreviews/0035tr.pdf>. December 19, 2014.

9. REFERENCES

- EPA. 2008c. Toxicological review of 2,2',4,4',5,5'-hexabromodiphenyl ether (BDE-153) (CAS No. 68631-49-2). In support of summary information on the Integrated Risk Information System (IRIS). Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/iris/toxreviews/1009tr.pdf>. December 18, 2014
- EPA. 2008d. Toxicological review of 2,2',4,4'-tetrabromodiphenyl ether (BDE-47) (CAS No. 5436-43-1). In support of summary information on the Integrated Risk Information System (IRIS). Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/iris/toxreviews/1010tr.pdf>. December 19, 2014
- EPA. 2009a. Drinking water contaminant candidate list 3-final. U.S. Environmental Protection Agency. Federal Register 74 FR 51850:51850-51862. <http://www.gpo.gov/fdsys/pkg/FR-2009-10-08/pdf/E9-24287.pdf>. September 9, 2014.
- EPA. 2009b. National primary drinking water regulations. Washington, DC: U.S. Environmental Protection Agency, Office of Ground Water and Drinking Water. EPA816F090004. <http://water.epa.gov/drink/contaminants/upload/mcl-2.pdf>. September 10, 2014.
- EPA. 2009c. The national study of chemical residues in lake fish tissue. Washington, DC: U.S. Environmental Protection Agency, Office of Water. EPA823R09006.
- EPA. 2010. An exposure assessment of polybrominated diphenyl ethers. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development. EPA600R08086F.
- EPA. 2012. 2012 Edition of the drinking water standards and health advisories. Washington, DC: U.S. Environmental Protection Agency, Office of Water. EPA822S12001. <http://water.epa.gov/action/advisories/drinking/upload/dwstandards2012.pdf>. September 9, 2014.
- EPA. 2013a. AEGL chemical data. Final AEGLs (131). Washington, DC: U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics. http://www.epa.gov/oppt/aegl/pubs/compiled_aegls_update_nov2013.pdf. September 9, 2014.
- EPA. 2013b. List of substances. Table 1 to 68.130-List of regulated toxic substances and threshold quantities for accidental release prevention. U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 68.130, Subpart F. <http://www.gpo.gov/fdsys/pkg/CFR-2013-title40-vol16/pdf/CFR-2013-title40-vol16-sec68-130.pdf>. September 24, 2014.
- EPA. 2013c. Designation of hazardous substances. U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 116.4, Subpart D. <http://www.gpo.gov/fdsys/pkg/CFR-2013-title40-vol23/pdf/CFR-2013-title40-vol23-sec116-4.pdf>. September 9, 2014.
- EPA. 2013d. Determination of reportable quantities. Table 117.3. Reportable quantities of hazardous substances designated pursuant to section 311 of the Clean Water Act. U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 117.3, Subpart A. <http://www.gpo.gov/fdsys/pkg/CFR-2013-title40-vol23/pdf/CFR-2013-title40-vol23-part117-subpartA.pdf>. September 9, 2014.

9. REFERENCES

- EPA. 2013e. Identification and listing of hazardous waste. Appendix VIII to Part 261-Hazardous constituents. U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 261, Appendix VIII to Part 261. <http://www.gpo.gov/fdsys/pkg/CFR-2013-title40-vol27/pdf/CFR-2013-title40-vol27-part261-appVIII.pdf>. September 10, 2014.
- EPA. 2013f. Groundwater monitoring list. U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 264, Appendix IX to Part 264. <http://www.gpo.gov/fdsys/pkg/CFR-2013-title40-vol27/pdf/CFR-2013-title40-vol27-part264-appIX.pdf>. September 10, 2014.
- EPA. 2013g. Designation of hazardous substances. Table 302.4-List of hazardous substances and reportable quantities. U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 302.4. <http://www.gpo.gov/fdsys/pkg/CFR-2013-title40-vol29/pdf/CFR-2013-title40-vol29-sec302-4.pdf>. September 10, 2014.
- EPA. 2013h. Chemicals and chemical categories to which this part applies. U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 372.65, Subpart D. <http://www.gpo.gov/fdsys/pkg/CFR-2013-title40-vol29/pdf/CFR-2013-title40-vol29-part372.pdf>. September 10, 2014.
- EPA. 2013i. The list of extremely hazardous substances and their threshold planning quantities. U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 355, Appendix A to Part 355. <http://www.gpo.gov/fdsys/pkg/CFR-2013-title40-vol29/pdf/CFR-2013-title40-vol29-part355-appA.pdf>. September 10, 2014.
- EPA. 2013j. Semiannual regulatory flexibility agenda and semiannual regulatory agenda. U.S. Environmental Protection Agency. <http://www.reginfo.gov>. August 12, 2014.
- EPA. 2013k. Contaminants of Emerging Concern (CECs) in fish: Polybrominated diphenyl ethers (PBDEs). Washington, DC: U.S. Environmental Protection Agency, Office of Water. EPA820F13003.
- EPA. 2014a. The Clean Air Act amendments of 1990 list of hazardous air pollutants. Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/ttn/atw/orig189.html>. September 9, 2014.
- EPA. 2014b. National ambient air quality standards (NAAQS). Washington, DC: U.S. Environmental Protection Agency, Office of Air and Radiation. <http://www.epa.gov/air/criteria.html>. January 08, 2014.
- EPA. 2014c. National recommended water quality criteria. Washington, DC: U.S. Environmental Protection Agency, Office of Water, Office of Science and Technology. <http://water.epa.gov/scitech/swguidance/standards/criteria/current/index.cfm>. January 08, 2014.
- EPA. 2014d. Master testing list. Washington, DC: U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics. <http://www.epa.gov/opptintr/chemtest/pubs/index1.pdf>. September 10, 2014.
- EPA. 2014e. Estimation Program Interface Suite™ for Microsoft Windows®, v4.10. U.S. Environmental Protection Agency. <http://www.epa.gov/opptintr/exposure/pubs/episuitedl.htm>. October 1, 2014.
- Eriksson P, Fischer C, Fredriksson A. 2006. Polybrominated diphenyl ethers, a group of brominated flame retardants, can interact with polychlorinated biphenyls in enhancing developmental neurobehavioral defects. *Toxicol Sci* 94(2):302-309.

9. REFERENCES

- Eriksson P, Jakobsson E, Fredriksson A. 1998. Developmental neurotoxicity of brominated flame-retardants, polybrominated diphenyl ethers, and tetrabromo-bis-phenol A. *Organohalogen Compounds* 35:375-377.
- Eriksson P, Jakobsson E, Fredriksson A. 2001. Brominated flame retardants: A novel class of developmental neurotoxicants in our environment. *Environ Health Perspect* 109(9):903-908.
- Eriksson P, Viberg H, Fischer C, et al. 2002a. A comparison on developmental neurotoxic effects of hexabromocyclododecane, 2,2',4,4',5,5'-hexabromodiphenyl ether (PBDE 153) and 2,2',4,4',5,5'-hexachlorobiphenyl (PCB 153). *Organohalogen Compounds* 57:389-390.
- Eriksson P, Viberg H, Jakobsson E, et al. 1999. PBDE 2,2', 4,4', 5-pentabromodiphenyl ether, causes permanent neurotoxic effects during a defined period of neonatal brain development. *Organohalogen Compounds* 40:333-336.
- Eriksson P, Viberg H, Jakobsson E, et al. 2002b. A brominated retardant, 2,2',4,4',5-pentabromodiphenyl ether: Uptake, retention, and induction of neurobehavioral alterations in mice during a critical phase of neonatal brain development. *Toxicol Sci* 67:98-103.
- Erkin-Cakmak A, Harley KG, Chevrier J, et al. 2015. In utero and childhood polybrominated diphenyl ether exposures and body mass at age 7 years: The CHAMACOS study. *Environ Health Perspect* 123(6):636-642. 10.1289/ehp.1408417.
- Ernest SR, Wade MG, Lalancette C, et al. 2012. Effects of chronic exposure to an environmentally relevant mixture of brominated flame retardants on the reproductive and thyroid system in adult male rats. *Toxicol Sci* 127(2):496-507.
- Erratico CA, Moffatt SC, Bandiera SM. 2011. Comparative oxidative metabolism of BDE-47 and BDE-99 by rat hepatic microsomes. *Toxicol Sci* 123(1):37-47.
- Erratico CA, Szeitz A, Bandiera SM. 2012. Oxidative metabolism of BDE-99 by human liver microsomes: Predominant role of CYP2B6. *Toxicol Sci* 129(2):280-292.
- Erratico CA, Szeitz A, Bandiera SM. 2013. Biotransformation of 2,2',4,4'-tetrabromodiphenyl ether (BDE-47) by human liver microsomes: Identification of cytochrome P450 2B6 as the major enzyme involved. *Chem Res Toxicol* 26(5):721-731.
- Eskenazi B, Chevrier J, Rauch SA, et al. 2013. *In utero* and childhood polybrominated diphenyl ether (PBDE) exposures and neurodevelopment in the CHAMACOS study. *Environ Health Perspect* 121(2):257-262. 10.1289/ehp.1205597.
- EU. 2001. Diphenyl ether, pentabromo derivative (pentabromodiphenyl ether). European Union Risk Assessment Report. Luxembourg: Office for Official Publications of the European Committees, 1-124.
- EU. 2002. Bis(pentabromophenyl) ether. European Union Risk Assessment Report. Luxembourg: Office for Official Publications of the European Committees. EINECS No. 214-604-9, 1-279.
- EU. 2003a. Diphenyl ether, octabromo derivative. European Union Risk Assessment Report. Luxembourg: Office for Official Publications of the European Committees, 1-279.

9. REFERENCES

- EU. 2003b. Consolidated TEXT. CONSLEG system. Office for Official Publications of the European Communities. European Union. http://europa.eu.int/erullex/en/consleg/pdf/2003/en_2003L0011_do_001.pdf. September 01, 2004.
- Evandri MG, Mastrangelo S, Costa LG, et al. 2003. *In vitro* assessment of mutagenicity and clastogenicity of BDE-99, a pentabrominated diphenyl ether flame retardant. *Environ Mol Mutagen* 42(2):85-90.
- Eyster JT, Humphrey HEB, Kimbrough RD. 1983. Partitioning of polybrominated biphenyls (PBBs) in serum, adipose tissue, breast milk, placenta, cord blood, and biliary fluid, and feces. *Arch Environ Health* 38:47-53.
- Fan CY, Besas J, Kodavanti PR. 2010. Changes in mitogen-activated protein kinase in cerebellar granule neurons by polybrominated diphenyl ethers and polychlorinated biphenyls. *Toxicol Appl Pharmacol* 245(1):1-8.
- Fattore E, Filipsson AF, Hanberg A, et al. 2001. Toxicity of a technical mixture of polybrominated diphenyl ethers following 28 days of oral exposure in male and female rats. *Organohalogen Compounds* 53:357-361
- Fawkes J, Albro PW, Walters DB, et al. 1982. Comparison of extraction methods for determination of polybrominated biphenyl residues in animal tissue. *Anal Chem* 54:1866-1871.
- FDA. 2013. Bottled water. U.S. Food and Drug Administration. Code of Federal Regulations 21 CFR 165.110, Subpart B.
- FDA. 2014. Everything added to food in the United States (EAFUS). Washington, DC: U.S. Food and Drug Administration. <http://www.accessdata.fda.gov/scripts/fcn/fcnavigation.cfm?rpt=eafuslisting>. January 08, 2014.
- Fehring NV. 1975. Determination of polybrominated biphenyl residues in dairy products. *J Assoc Off Anal Chem* 58(5):978-982.
- Feng C, Xu Q, Jin Y, et al. 2016a. Determination of urinary bromophenols (BrPs) as potential biomarkers for human exposure to polybrominated diphenyl ethers (PBDEs) using gas chromatography-tandem mass spectrometry (GC-MS/MS). *J Chromatogr B Analyt Technol Biomed Life Sci* 1022:70-74. 10.1016/j.jchromb.2016.03.041.
- Feng Y, Hu Q, Meng G, et al. 2015. Simulating long-term occupational exposure to decabrominated diphenyl ether using C57BL/6 mice: Biodistribution and pathology. *Chemosphere* 128:118-124. 10.1016/j.chemosphere.2015.01.012.
- Feng Y, Zeng W, Wang Y, et al. 2016b. Long-term exposure to high levels of decabrominated diphenyl ether inhibits CD4 T-cell functions in C57Bl/6 mice. *J Appl Toxicol* 36(9):1112-1119. 10.1002/jat.3270.
- Feo ML, Gross MS, McGarrigle BP, et al. 2013. Biotransformation of BDE-47 to potentially toxic metabolites is predominantly mediated by human CYP2B6. *Environ Health Perspect* 121(4):440-446.
- Fernlof G, Gadhasson I, Podra K, et al. 1997. Lack of effects of some individual polybrominated diphenyl ether (PBDE) and polychlorinated biphenyl congeners on human lymphocyte functions *in vitro*. *Toxicol Lett* 90(2-3):189-197.

9. REFERENCES

- Fischer C, Fredriksson A, Eriksson P. 2008. Coexposure of neonatal mice to a flame retardant PBDE 99 (2,2',4,4',5-pentabromodiphenyl ether) and methyl mercury enhances developmental neurotoxic defects. *Toxicol Sci* 101(2):275-285.
- Fisher DA, Brown RS. 2000. Thyroid physiology in the prenatal period and during childhood. In: Braverman LE, Utiger RD, eds. *Werner and Ingbar's the thyroid*. Philadelphia, PA: Lippincott Williams & Wilkins, 959-972.
- Fitzgerald EF, Shrestha S, Gomez MI, et al. 2012. Polybrominated diphenyl ethers (PBDEs), polychlorinated biphenyls (PCBs) and neuropsychological status among older adults in New York. *Neurotoxicology* 33(1):8-15.
- Fomon SJ. 1966. Body composition of the infant: Part I: The male "reference infant". In: Falkner F, ed. *Human development*. Philadelphia, PA: WB Saunders, 239-246.
- Fomon SJ, Haschke F, Ziegler EE, et al. 1982. Body composition of reference children from birth to age 10 years. *Am J Clin Nutr* 35(Suppl 5):1169-1175.
- Foster WG, Gregorovich S, Morrison KM, et al. 2011. Human maternal and umbilical cord blood concentrations of polybrominated diphenyl ethers. *Chemosphere* 84(10):1301-1309. 10.1016/j.chemosphere.2011.05.028.
- Fowles JR, Fairbrother A, Baecher-Steppan L, et al. 1994. Immunologic and endocrine effects of the flame-retardant pentabromodiphenyl ether (DE-71) in C57BL/6J mice. *Toxicology* 86(1-2):49-61.
- Frederiksen M, Vorkamp K, Mathiesen L, et al. 2010. Placental transfer of the polybrominated diphenyl ethers BDE-47, BDE-99 and BDE-209 in a human placenta perfusion system: An experimental study. *Environ Health* 9:32.
- Frederiksen M, Vorkamp K, Thomsen M, et al. 2009. Human internal and external exposure to PBDEs—a review of levels and sources. *Int J Hyg Environ Health* 212(2):109-134.
- Fries GF. 1985. The PBB episode in Michigan: An overall appraisal. *CRC Crit Rev Toxicol* 16:105-156.
- Frouin H, Lebeof M, Hammill M, et al. 2011. PBDEs in serum and blubber of harbor, grey and harp seal pups from Eastern Canada. *Chemosphere* 82(5):663-669.
- Fujii Y, Nishimura E, Kato Y, et al. 2014. Dietary exposure to phenolic and methoxylated organohalogen contaminants in relation to their concentrations in breast milk and serum in Japan. *Environ Int* 63:19-25
- Fujimoto H, Woo GH, Inoue K, et al. 2011. Impaired oligodendroglial development by decabromodiphenyl ether in rat offspring after maternal exposure from mid-gestation through lactation. *Reprod Toxicol* 31(1):86-94.
- Furl C, Meredith C. 2010. *PBT monitoring: PBDE flame retardants in Spokane River fish, 2009*. Olympia, Washington: Washington State Department of Ecology, Toxics Studies Unit.

9. REFERENCES

- Gallego E, Grimalt JO, Bartrons M, et al. 2007. Altitudinal gradients of PBDEs and PCBs in fish from European high mountain lakes. *Environ Sci Technol* 41(7):2196-2202.
- Garcia-Reyero N, Escalon BL, Prats E, et al. 2014. Effects of BDE-209 contaminated sediments on zebrafish development and potential implications to human health. *Environ Int* 63:216-223.
- Gascon M, Fort M, Martinez D, et al. 2012. Polybrominated diphenyl ethers (PBDEs) in breast milk and neuropsychological development in infants. *Environ Health Perspect* 120(12):1760-1765.
- Gascon M, Vrijheid M, Martinez D, et al. 2011. Effects of pre and postnatal exposure to low levels of polybromodiphenyl ethers on neurodevelopment and thyroid hormone levels at 4 years of age. *Environ Int* 37(3):605-611.
- Gassmann K, Schreiber T, Dingemans MM, et al. 2014. BDE-47 and 6-OH-BDE-47 modulate calcium homeostasis in primary fetal human neural progenitor cells via ryanodine receptor-independent mechanisms. *Arch Toxicol* 88(8):1537-1548. 10.1007/s00204-014-1217-7.
- Gauthier MS, Rabasa-Lhoret R, Prud'homme D, et al. 2014. The metabolically healthy but obese phenotype is associated with lower plasma levels of persistent organic pollutants as compared to the metabolically abnormal obese phenotype. *J Clin Endocrinol Metab* 99(6):E1061-1066. 10.1210/jc.2013-3935.
- Gaylor MO, Harvey E, Hale RC. 2013. Polybrominated diphenyl ether (PBDE) accumulation by earthworms (*Eisenia fetida*) exposed to biosolids-, polyurethane foam microparticle-, and penta-BDE-amended soils. *Environ Sci Technol* 47(23):13831-13839. 10.1021/es403750a.
- Gee JR, Moser VC. 2008. Acute postnatal exposure to brominated diphenylether 47 delays neuromotor ontogeny and alters motor activity in mice. *Neurotoxicol Teratol* 30(2):79-87.
- Gee JR, Hedge JM, Moser VC. 2008. Lack of alterations in thyroid hormones following exposure to polybrominated diphenyl ether 47 during a period of rapid brain development in mice. *Drug Chem Toxicol* 31(2):245-254.
- Gill U, Chu I, Ryan JJ, et al. 2004. Polybrominated diphenyl ethers: Human tissue levels and toxicology. *Rev Environ Contam Toxicol* 183:55-97.
- Giwercman A, Carlsen E, Keiding N, et al. 1993. Evidence for increasing incidence of abnormalities of the human testis: A review. *Environ Health Perspect Suppl* 101(2):65-71.
- Glinoyer D, De Nayer P, Bourdoux P, et al. 1990. Regulation of maternal thyroid during pregnancy. *J Clin Endocrinol Metab* 71:276-287.
- Great Lakes Chemical Corporation. 1978. Octabromodiphenyl ether. Subacute inhalation toxicity study in rats. International Research and Development Corporation. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. OTS0522293.
- Great Lakes Chemical Corporation. 2000. A 90-day inhalation toxicity study of octabromodiphenyl oxide in albino rats, dated 04/04/02. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8E. OTS0574171-1.

9. REFERENCES

- Gregoraszczyk EL, Dobrzanska G, Karpeta A. 2015. Effects of 2,2',4,4'-tetrabromodiphenyl ether (BDE47) on the enzymes of phase I (CYP2B1/2) and phase II (SULT1A and COMT) metabolism, and differences in the action of parent BDE-47 and its hydroxylated metabolites, 5-OH-BDE-47 and 6-OH-BDE47, on steroid secretion by luteal cells. *Environ Toxicol Pharmacol* 40(2):498-507. 10.1016/j.etap.2015.07.011.
- Gregoraszczyk EL, Siembida M, Grzyb D, et al. 2012. Polybrominated diphenylethers (PBDEs) act as apoptotic factors in the corpus luteum in addition to having a short-term stimulatory effect on progesterone secretion by luteal cells. *Toxicol Mech Methods* 22(2):131-138. 10.3109/15376516.2011.606433.
- Gross MS, Butryn DM, McCarrigle BP, et al. 2015. Primary role of cytochrome P450 2B6 in the oxidative metabolism of 2,2',4,4',6-pentabromodiphenyl ether (BDE-100) to hydroxylated BDEs. *Chem Res Toxicol* [pub ahead of print].
- Gump BB, Yun S, Kannan K. 2014. Polybrominated diphenyl ether (PBDE) exposure in children: Possible associations with cardiovascular and psychological functions. *Environ Res* 132:244-250. 10.1016/j.envres.2014.04.009.
- Guo W, Holden A, Smith SC, et al. 2016. PBDE levels in breast milk are decreasing in California. *Chemosphere* 150:505-513. 10.1016/j.chemosphere.2015.11.032.
- Gustafsson K, Bjork M, Burreau S, et al. 1999. Bioaccumulation kinetic of brominated flame retardants (polybrominated diphenyl ethers) in blue mussels (*Mytilus edulis*). *Environ Toxicol Chem* 18(6):1218-1224.
- Guyot R, Chatonnet F, Gillet B, et al. 2014. Toxicogenomic analysis of the ability of brominated flame retardants TBBPA and BDE-209 to disrupt thyroid hormone signaling in neural cells. *Toxicology* 325:125-132. 10.1016/j.tox.2014.08.007.
- Guzelian PS, Henry CJ, Olin SS, eds. 1992. Similarities and differences between children and adults: Implications for risk assessment. Washington, DC: International Life Sciences Institute Press.
- Haddad S, Poulin P, Krishnan K. 2000. Relative lipid content as the sole mechanistic determinant of the adipose tissue:blood partition coefficients of highly lipophilic organic chemicals. *Chemosphere* 40(8):839-843.
- Hagenmaier H, She J, Benz T, et al. 1992. Analysis of sewage sludge for polyhalogenated dibenzo-p-dioxins, dibenzofurans, and diphenylethers. *Chemosphere* 25(1-10):1457-1462.
- Haglund PS, Zook DR, Buser H-R, et al. 1997. Identification and quantification of polybrominated ethers and methoxy-polybrominated diphenyl ethers in Baltic biota. *Environ Sci Technol* 31:3281-3287.
- Hagmar L, Bjork J, Sjodin A, et al. 2001. Plasma levels of persistent organohalogenes and hormone levels in adult male humans. *Arch Environ Health* 56(2):138-143.
- Hagmar L, Jakobsson K, Thuresson K, et al. 2000. Computer technicians are occupationally exposed to polybrominated diphenyl ethers and tetrabromobisphenol A. *Organohalogen Compounds* 47:202-205.
- Hakk H, Huwe J, Lorentzen M. 2001. A mass balance study of a commercial pentabromodiphenyl ether mixture in male Sprague-Dawley rats. *Organohalogen Compounds* 52:5-8.

9. REFERENCES

- Hakk H, Huwe J, Low M, et al. 2006. Tissue disposition, excretion and metabolism of 2,2',4,4',6-pentabromodiphenyl ether (BDE-100) in male Sprague-Dawley rats. *Xenobiotica* 36(1):79-94.
- Hakk H, Huwe JK, Larsen GL. 2009. Absorption, distribution, metabolism and excretion (ADME) study with 2,2',4,4',5,6'-hexabromodiphenyl ether (BDE-154) in male Sprague-Dawley rats. *Xenobiotica* 39(1):46-56.
- Hakk H, Larsen G, Bergman A, et al. 2002b. Binding of brominated diphenyl ethers to male rat carrier proteins. *Xenobiotica* 32(12):1079-1091.
- Hakk H, Larsen G, Klasson-Wehler E. 2002a. Tissue disposition, excretion and metabolism of 2,2',4,4',5-pentabromodiphenyl ether (BDE-99) in the male Sprague-Dawley rat. *Xenobiotica* 32(5):369-382.
- Hale RC, Alaee M, Manchester-Neesvig JB, et al. 2003. Polybrominated diphenyl ether flame retardants in the North American environment. *Environ Int* 29:771-779.
- Hale RC, La Guardia MJ, Harvey EP, et al. 2000. Comparison of brominated diphenyl ether fire retardant and organochlorine burdens in fish from Virginia (USA). *Organohalogen Compounds* 47:65-68.
- Hale RC, La Guardia MJ, Harvey EP, et al. 2001b. Polybrominated diphenyl ether flame retardants in Virginia freshwater fishes (USA). *Environ Sci Technol* 35(23):4585-4591.
- Hale RC, La Guardia MJ, Harvey EP, et al. 2001a. Persistent pollutants in land-applied sludges. *Nature* 412(6843):140-141.
- Hale RC, La Guardia MJ, Harvey EP, et al. 2002. Potential role of fire retardant-treated polyurethane foam as a source of brominated diphenyl ethers to the US environment. *Chemosphere* 46(5):729-735.
- Hale RC, La Guardia MJ, Harvey E, et al. 2012. Polybrominated diphenyl ethers in U.S. sewage sludges and biosolids: Temporal and geographical trends and uptake by corn following land application. *Environ Sci Technol* 46(4):2055-2063.
- Hallgren S, Darnerud P. 1998. Effects of polybrominated diphenyl ethers (PBDEs), polychlorinated biphenyls (PCBs) and chlorinated paraffins (CPs) on thyroid hormone levels and enzyme activities in rats. *Organohalogen Compounds* 35:391-394.
- Hallgren S, Darnerud PO. 2002. Polybrominated diphenyl ethers (PBDEs), polychlorinated biphenyls (PCBs) and chlorinated paraffins (CPs) in rats-testing interactions and mechanisms for thyroid hormone effects. *Toxicology* 177(2-3):227-243.
- Hallgren S, Viberg H. 2016. Postnatal exposure to PFOS, but not PBDE 99, disturb dopaminergic gene transcription in the mouse CNS. *Environ Toxicol Pharmacol* 41:121-126. 10.1016/j.etap.2015.11.016.
- Hallgren S, Fredriksson A, Viberg H. 2015. More signs of neurotoxicity of surfactants and flame retardants - Neonatal PFOS and PBDE 99 cause transcriptional alterations in cholinergic genes in the mouse CNS. *Environ Toxicol Pharmacol* 40(2):409-416. 10.1016/j.etap.2015.06.014.

9. REFERENCES

- Hallgren S, Sinjari T, Hakansson H, et al. 2001. Effects of polybrominated diphenyl ethers (PBDEs) and polychlorinated biphenyls (PCBs) on thyroid hormone and vitamin A levels in rats and mice. *Arch Toxicol* 75(4):200-208.
- Hamers T, Kamstra JH, Sonneveld E, et al. 2006. *In vitro* profiling of the endocrine-disrupting potency of brominated flame retardants. *Toxicol Sci* 92(1):157-173.
- Hamm S, Strickeling M, Ranken PF, et al. 2001. Determination of polybrominated diphenyl ethers and PBDD/Fs during the recycling of high impact polystyrene containing decabromodiphenyl ether and antimony oxide. *Chemosphere* 44(6):1353-1360.
- Han G, Ding G, Lou X, et al. 2011. Correlations of PCBs, dioxin, and PBDE with TSH in children's blood in areas of computer E-waste recycling. *Biomed Environ Sci* 24(2):112-116.
- Hardell L, Bavel B, Lindstrom G, et al. 2006. *In utero* exposure to persistent organic pollutants in relation to testicular cancer risk. *Int J Androl* 29(1):228-234.
- Hardell L, Lindstrom G, van Bavel B, et al. 1998. Concentrations of the flame retardant 2,2',4,4'-tetrabrominated diphenyl ether in human adipose tissue in Swedish persons and the risk for non-Hodgkin's lymphoma. *Oncol Res* 10(8):429-432.
- Hardy ML. 1999. Regulatory status and environmental properties of brominated flame retardants undergoing risk assessment in the EU: DBDPO, OBDPO, PEBDPO and HBCD. *Polym Degrad Stab* 64(3):545-556.
- Hardy ML. 2000a. Properties of the major commercial PBDPO flame retardant, DBDPO, in comparison to PBB and PCB. *Organohalogen Compounds* 47:233-236.
- Hardy ML. 2000b. The toxicity of the commercial polybrominated diphenyl oxide flame retardants: DBDPO, OBDPO, PeBDPO. *Organohalogen Compounds* 47:41-44.
- Hardy ML. 2001. Assessment of reported decabromodiphenyl oxide blood and air levels in Swedish workers and their workplace. The second international workshop on brominated flame retardants. BFR 2001. Stockholm, Sweden, 121-124.
- Hardy ML. 2002a. A comparison of the properties of the major commercial PBDPO/PBDE product to those of major PBB and PCB products. *Chemosphere* 45(5):717-728.
- Hardy ML. 2002b. The toxicology of the three commercial polybrominated diphenyl oxide (ether) flame retardants. *Chemosphere* 46(5):757-777.
- Hardy M, Biesemeier J, Manor O. 2001. Results of a prenatal developmental toxicity study of decabromodiphenyl oxide in rats. The second international workshop on brominated flame retardants. BFR 2001. Stockholm, Sweden, 253-257.
- Hardy ML, Banasik M, Stedeford T. 2009. Toxicology and human health assessment of decabromodiphenyl ether. *Crit Rev Toxicol* 39 Suppl 3:1-44.
- Hardy ML, Schroeder R, Biesemeier J, et al. 2002. Prenatal oral (gavage) developmental toxicity study of decabromodiphenyl oxide in rats. *Int J Toxicol* 21:83-91.

9. REFERENCES

- Harley KG, Chevrier J, Aguilar Schall R, et al. 2011. Association of prenatal exposure to polybrominated diphenyl ethers and infant birth weight. *Am J Epidemiol* 174(8):885-892.
- Harley KG, Marks AR, Chevrier J, et al. 2010. PBDE concentrations in women's serum and fecundability. *Environ Health Perspect* 118(5):699-704. 10.1093/aje/kwr212.
- Harner T. 2001. Measurements of octanol-air partition coefficients (KOA) for polybrominated diphenyl ethers (PBDEs): Predicting partitioning in the environment. The second international workshop on brominated flame retardants. BFR 2001. Stockholm, Sweden, 55-58.
- Harrad S, Hunter S. 2006. Concentrations of polybrominated diphenyl ethers in air and soil on a rural-urban transect across a major UK conurbation. *Environ Sci Technol* 40(15):4548-4553.
- Harrad S, Hazrati S, Ibarra C. 2006. Concentrations of polychlorinated biphenyls in indoor air and polybrominated diphenyl ethers in indoor air and dust in Birmingham, United Kingdom: Implications for human exposure. *Environ Sci Technol* 40(15):4633-4638.
- Harrad S, Ibarra C, Diamond M, et al. 2008. Polybrominated diphenyl ethers in domestic indoor dust from Canada, New Zealand, United Kingdom and United States. *Environ Int* 34(2):232-238.
- Harrad S, Wijesekera R, Hunter S, et al. 2004. Preliminary assessment of U.K. human dietary and inhalation exposure to polybrominated diphenyl ethers. *Environ Sci Technol* 38:2345-2350.
- Hazrati S, Harrad S. 2006. Causes of variability in concentrations of polychlorinated biphenyls and polybrominated diphenyl ethers in indoor air. *Environ Sci Technol* 40(24):7584-7589.
- Hazrati S, Harrad S. 2007. Calibration of polyurethane foam (PUF) disk passive air samplers for quantitative measurement of polychlorinated biphenyls (PCBs) and polybrominated diphenyl ethers (PBDEs): Factors influencing sampling rates. *Chemosphere* 67:448-455.
- He P, He W, Wang A, et al. 2008b. PBDE-47-induced oxidative stress, DNA damage and apoptosis in primary cultured rat hippocampal neurons. *Neurotoxicology* 29(1):124-129.
- He P, Wang A, Niu Q, et al. 2011. Toxic effect of PBDE-47 on thyroid development, learning, and memory, and the interaction between PBDE-47 and PCB153 that enhances toxicity in rats. *Toxicol Ind Health* 27(3):279-288.
- He P, Wang AG, Xia T, et al. 2009. Mechanisms underlying the developmental neurotoxic effect of PBDE-47 and the enhanced toxicity associated with its combination with PCB153 in rats. *Neurotoxicology* 30(6):1088-1095.
- He W, He P, Wang A, et al. 2008a. Effects of PBDE-47 on cytotoxicity and genotoxicity in human neuroblastoma cells *in vitro*. *Mutat Res* 649(1-2):62-70.
- He Y, Murphy MB, Yu RM, et al. 2008c. Effects of 20 PBDE metabolites on steroidogenesis in the H295R cell line. *Toxicol Lett* 176(3):230-238.
- Hearn LK, Hawker DW, Toms LM, et al. 2013. Assessing exposure to polybrominated diphenyl ethers (PBDEs) for workers in the vicinity of a large recycling facility. *Ecotoxicol Environ Saf* 92:222-228.

9. REFERENCES

- Helleday T, Tuominen KL, Bergman A, et al. 1999. Brominated flame retardants induce intragenic recombination in mammalian cells. *Mutat Res* 439(2):137-147.
- Hendriks HS, Antunes Fernandes EC, Bergman A, et al. 2010. PCB-47, PBDE-47, and 6-OH-PBDE-47 differentially modulate human GABA_A and $\alpha_4\beta_2$ nicotinic acetylcholine receptors. *Toxicol Sci* 118(2):635-642.
- Hendriks HS, Meijer M, Muilwijk M, et al. 2014. A comparison of the in vitro cyto- and neurotoxicity of brominated and halogen-free flame retardants: Prioritization in search for safe(r) alternatives. *Arch Toxicol* 88(4):857-869. 10.1007/s00204-013-1187-1.
- Herbstman JB, Sjodin A, Apelberg BJ, et al. 2008. Birth delivery mode modifies the associations between prenatal polychlorinated biphenyl (PCB) and polybrominated diphenyl ether (PBDE) and neonatal thyroid hormone levels. *Environ Health Perspect* 116(10):1376-1382.
- Herbstman JB, Sjodin A, Kurzon M, et al. 2010. Prenatal exposure to PBDEs and neurodevelopment. *Environ Health Perspect* 118(5):712-719.
- Heredia L, Torrente M, Colomina MT, et al. 2012. Behavioral effects of oral subacute exposure to BDE-209 in young adult mice: A preliminary study. *Food Chem Toxicol* 50(3-4):707-712.
- Hertz-Picciotto I, Bergman A, Fangstrom B, et al. 2011. Polybrominated diphenyl ethers in relation to autism and developmental delay: A case-control study. *Environ Health* 10(1):1.
- Hesse JL, Powers RA. 1978. Polybrominated biphenyl (PBB) contamination of the Pine River, Gratiot, and Midland counties, Michigan. *Environ Health Perspect* 23:19-25.
- Hillery BR, Basu I, Sweet CW, et al. 1997. Temporal and spatial trends in a long-term study of gas-phase PCB concentrations near the Great Lakes. *Environ Sci Technol* 31:1811-1816.
- Hites RA. 2004. Polybrominated diphenyl ethers in the environment and in people: A meta-analysis of concentrations. *Environ Sci Technol* 38(4):945-956.
- Ho KL, Yau MS, Murphy MB, et al. 2015. Urinary bromophenol glucuronide and sulfate conjugates: Potential human exposure molecular markers for polybrominated diphenyl ethers. *Chemosphere* 133:6-12. 10.1016/j.chemosphere.2015.03.003.
- Hoel DG, Davis DL, Miller AB, et al. 1992. Trends in cancer mortality in 15 industrialized countries, 1969-1986. *J Natl Cancer Inst* 84(5):313-320.
- Hoffman K, Adgent M, Goldman BD, et al. 2012. Lactational exposure to polybrominated diphenyl ethers and its relation to social and emotional development among toddlers. *Environ Health Perspect* 120(10):1438-1442.
- Hoffman K, Webster TF, Sjodin A, et al. 2016. Toddler's behavior and its impacts on exposure to polybrominated diphenyl ethers. *J Expo Sci Environ Epidemiol*:1-5. 10.1038/jes.2016.11.
- Hoh E, Hites RA. 2005. Brominated flame retardants in the atmosphere of the east-central United States. *Environ Sci Technol* 39(20):7794-7802.

9. REFERENCES

- Hollowell G, Norman W, Staehling W, et al. 1998. Iodine nutrition in the United States. Trends and public health implications: Iodine excretion data from national health and nutrition examination surveys I and III (1971-1974 and 1988-1994). *J Clin Endocrinol Metab* 83(10):3401-3408.
- Holmes AK, Koller KR, Kieszak SM, et al. 2014. Case-control study of breast cancer and exposure to synthetic environmental chemicals among Alaska Native women. *Int J Circumpolar Health* 73:25760. 10.3402/ijch.v73.25760.
- Hong SK, Sohn KH, Kim IY, et al. 2010. Polybrominated diphenyl ethers orally administration to mice were transferred to offspring during gestation and lactation with disruptions on the immune system. *Immune Network* 10(2):64-74.
- Hooper H, McDonald TA. 2000. The PBDEs: An emerging environmental challenge and another reason for breast-milk monitoring programs. *Environ Health Perspect* 108(5):387-392.
- Hooper K, She J, Sharp M, et al. 2007. Depuration of polybrominated diphenyl ethers (PBDEs) and polychlorinated biphenyls (PCBs) in breast milk from California first-time mothers (primiparae). *Environ Health Perspect* 115(9):1271-1275.
- Hoppe AA, Carey GB. 2007. Polybrominated diphenyl ethers as endocrine disruptors of adipocyte metabolism. *Obesity* 15(12):2942-2950.
- Hori S, Akutsu K, Kitagawa M, et al. 2000. Development of analysis for polybrominated diphenyl ether in seafood and actual contamination of seafood. *Organohalogen Compounds* 47:214-217.
- Hovander L, Malmberg T, Athanasiadou M, et al. 2002. Identification of hydroxylated PCB metabolites and other phenolic halogenated pollutants in human blood plasma. *Arch Environ Contam Toxicol* 42:105-117.
- Howie L, Dickerson R, Davis D, et al. 1990. Immunosuppressive and monooxygenase induction activities of polychlorinated diphenyl ether congeners in C57BL/6N mice: Quantitative structure-activity relationships. *Toxicol Appl Pharmacol* 105:254-263.
- HSDB. 2012. Polybrominated diphenyl ethers. Hazardous Substances Data Bank. National Library of Medicine. <http://toxnet.nlm.nih.gov>. October 9, 2014.
- Hua I, Kang N, Jafvert CT, et al. 2003. Heterogeneous photochemical reactions of decabromodiphenyl ether. *Environ Toxicol Chem* 22:798-804.
- Huang F, Wen S, Li J, et al. 2014. The human body burden of polybrominated diphenyl ethers and their relationships with thyroid hormones in the general population in Northern China. *Sci Total Environ* 466-467:609-615.
- Huang S, Cui Y, Guo X, et al. 2015. 2,2',4,4'-Tetrabromodiphenyl ether disrupts spermatogenesis, impairs mitochondrial function and induces apoptosis of early leptotene spermatocytes in rats. *Reprod Toxicol* 51:114-124. 10.1016/j.reprotox.2015.01.009.
- Huang SC, Giordano G, Costa LG. 2010. Comparative cytotoxicity and intracellular accumulation of five polybrominated diphenyl ether congeners in mouse cerebellar granule neurons. *Toxicol Sci* 114(1):124-132.

9. REFERENCES

- Hughes MF, Edwards BC, Mitchell CT, et al. 2001. *In vitro* dermal absorption of flame retardant chemicals. *Food Chem Toxicol* 39(12):1263-1270.
- Hung H, Kallenborn R, Breivik K, et al. 2010. Atmospheric monitoring of organic pollutants in the Arctic under the Arctic Monitoring and Assessment Programme (AMAP), 1993-2006. *Sci Total Environ* 408(15):2854-2873.
- Hurley S, Reynolds P, Goldberg D, et al. 2011. Adipose levels of polybrominated diphenyl ethers and risk of breast cancer. *Breast Cancer Res Treat* 129(2):505-511.
- Huwe JK, Hakk H, Birnbaum LS. 2008. Tissue distribution of polybrominated diphenyl ethers in male rats and implications for biomonitoring. *Environ Sci Technol* 42(18):7018-7024.
- Huwe JK, Hakk H, Lorentzsen M. 2002b. A mass balance feeding study of a commercial octabromodiphenyl ether mixture in rats. *Organohalogen Compounds* 58:229-223.
- Huwe JK, Lorentzsen M, Thuresson K, et al. 2002a. Analysis of mono- to deca-brominated diphenyl ethers in chickens at the part per billion level. *Chemosphere* 46:635-640.
- IARC. 2014. Agents classified by the IARC monographs. Volumes 1-110. Lyon, France: International Agency for Research on Cancer.
<http://monographs.iarc.fr/ENG/Classification/ClassificationsCASOrder.pdf>. September 9, 2014.
- Ibhazehiebo K, Iwasaki T, Kimura-Kuroda J, et al. 2011. Disruption of thyroid hormone receptor-mediated transcription and thyroid hormone-induced Purkinje cell dendrite arborization by polybrominated diphenyl ethers. *Environ Health Perspect* 119(2):168-175.
- Ikonomou MG, Rayne S. 2002. Chromatographic and ionization properties of polybrominated diphenyl ethers using GC/high-resolution MS with metastable atom bombardment and electron impact ionization. *Anal Chem* 74:5263-5272.
- Ikonomou MG, Crewe N, He T, et al. 1999. Polybrominated-diphenyl-ether in biota samples from coastal British Columbia, Canada. *Organohalogen Compounds* 40:341-345.
- Ikonomou MG, Fisher M, He T, et al. 2000. Congener patterns, spatial and temporal trends of polybrominated diphenyl ethers in biota samples from the Canadian west coast and the northwest territories. *Organohalogen Compounds* 47:77-80.
- Ikonomou MG, Rayne S, Fischer M, et al. 2002. Occurrence and congener profiles of polybrominated diphenyl ethers (PBDEs) in environmental samples from coastal British Columbia, Canada. *Chemosphere* 46(5):649-663.
- Imamura, M, Tung, TC. 1984. A trial of fasting cure for PCB-poisoned patients in Taiwan. *Am J Ind Med* 5:147-153.
- IRDC. 1974. Acute toxicity studies in rats and rabbits with test data and cover letter. International Research and Development Corporation. Submitted to U.S. EPA under TSCA Section 8D. OTS0523319.

9. REFERENCES

- IRDC. 1975a. Octabromodiphenyl ether. Acute toxicity studies in rats and rabbits with cover sheet and letter dated 030890. International Research and Development Corporation. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. OTS052222968.
- IRDC. 1975b. Pentabromodiphenyl ether. Acute toxicity studies in rats and rabbits. In: The acute oral toxicity of pentabromodiphenyl ether to rats with cover sheets and letter dated 030890. International Research and Development Corporation. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. OTS05222286.
- IRDC. 1976. Decabromodiphenyl ether and octabromodiphenyl ether. In: A twenty-eight day toxicity study in rats with test data and cover sheet. International Research and Development Corporation. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. OTS0523322.
- IRDC. 1977. Octabromodiphenyl ether. Thirteen week feeding study in rats. In: Thirteen week feeding study in rats with attachments, cover sheet and letter dated 030890. International Research and Development Corporation. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. OTS0522297.
- IRIS. 2003a. Hexabromodiphenyl ether (CASRN 36483-60-0). Integrated Risk Information System. Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/iris/subst/0494.htm>. September 10, 2014.
- IRIS. 2003b. Nonabromodiphenyl ether (CASRN 63936-56-1). Integrated Risk Information System. Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/iris/subst/0495.htm>. September 10, 2014.
- IRIS. 2003c. Octabromodiphenyl ether (CASRN 32536-52-0). Integrated Risk Information System. Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/iris/subst/0180.htm>. September 10, 2014.
- IRIS. 2003d. Tetrabromodiphenyl ether (CASRN 40088-47-9). Integrated Risk Information System. Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/iris/subst/0493.htm>. September 10, 2014.
- IRIS. 2003e. Tribromodiphenyl ether (CASRN 49690-94-0). Integrated Risk Information System. Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/iris/subst/0492.htm>. September 10, 2014.
- IRIS. 2004. Pentabromodiphenyl ether (CASRN 32534-81-9). Integrated Risk Information System. Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/iris/subst/0184.htm>. September 10, 2014.
- IRIS. 2005. p,p'-Dibromodiphenyl ether (CASRN 2050-47-7). Integrated Risk Information System. Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/iris/subst/0491.htm>. September 10, 2014.
- IRIS. 2006. p-Bromodiphenyl ether (CASRN 101-55-3). Integrated Risk Information System. Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/iris/subst/0490.htm>. September 9, 2014.

9. REFERENCES

- IRIS. 2008a. 2,2',3,3',4,4',5,5',6,6'-Decabromodiphenyl ether (BDE-209) (CASRN 1163-19-5). Integrated Risk Information System. Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/iris/subst/0035.htm>. September 10, 2014.
- IRIS. 2008b. 2,2',4,4'-Tetrabromodiphenyl ether (BDE-47) (CASRN 5436-43-1). Integrated Risk Information System. Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/iris/subst/1010.htm>. September 10, 2014.
- IRIS. 2008c. 2,2',4,4',5-Pentabromodiphenyl ether (BDE-99) (CASRN 60348-60-9). Integrated Risk Information System. Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/iris/subst/1008.htm>. September 10, 2014.
- IRIS. 2008d. 2,2',4,4',5,5'-Hexabromodiphenyl ether (BDE-153) (CASRN 68631-49-2). Integrated Risk Information System. Washington, DC: U.S. Environmental Protection Agency. <http://www.epa.gov/iris/subst/1009.htm>. September 10, 2014.
- Ismail N, Gewurtz SB, Pleskach K, et al. 2009. Brominated and chlorinated flame retardants in Lake Ontario, Canada, lake trout (*Salvelinus namaycush*) between 1979 and 2004 and possible influences of food-web changes. *Environ Toxicol Chem* 28(5):910-920.
- Jakobsson K, Fang J, Athanasiadou M, et al. 2012. Polybrominated diphenyl ethers in maternal serum, umbilical cord serum, colostrum and mature breast milk. Insights from a pilot study and the literature. *Environ Int* 47:121-130.
- Jandacek RJ, Heubi JE, Buckley DD, et al. 2014. Reduction of the body burden of PCBs and DDE by dietary intervention in a randomized trial. *J Nutr Biochem* 25:483-488.
- Jansson B, Asplund L. 1987. Brominated flame retardants-ubiquitous environmental pollutants? *Chemosphere* 16(10-12):2343-2349.
- Jansson B, Andersson R, Asplund L, et al. 1991. Multiresidue method for the gas-chromatographic analysis of some polychlorinated and polybrominated pollutants in biological samples. *Fresenius Z Anal Chem* 340:439-445.
- Jansson B, Andersson R, Asplund L, et al. 1993. Chlorinated and brominated persistent organic compounds in biological samples from the environment. *Environ Toxicol Chem* 12(7):1163-1174.
- Jaret P. 2000. Health concerns: Defense systems under fire. *Natl Wildl* 38(6):36-41.
- Ji K, Choi K, Giesy JP, et al. 2011. Genotoxicity of several polybrominated diphenyl ethers (PBDEs) and hydroxylated PBDEs, and their mechanisms of toxicity. *Environ Sci Technol* 45(11):5003-5008.
- Johanson CE. 1980. Permeability and vascularity of the developing brain: Cerebellum vs cerebral cortex. *Brain Res* 190(1):3-16.
- Johansson N, Viberg H, Fredriksson A, et al. 2008. Neonatal exposure to deca-brominated diphenyl ether (PBDE 209) causes dose-response changes in spontaneous behaviour and cholinergic susceptibility in adult mice. *Neurotoxicology* 29(6):911-919. 10.1016/j.neuro.2008.09.008.
- Johnson A, Olson N. 2001. Analysis and occurrence of polybrominated diphenyl ethers in Washington state freshwater fish. *Arch Environ Contam Toxicol* 41(3):339-344.

9. REFERENCES

- Johnson PI, Altshul L, Cramer DW, et al. 2012. Serum and follicular fluid concentrations of polybrominated diphenyl ethers and *in vitro* fertilization outcome. *Environ Int* 45:9-14.
- Johnson PI, Stapleton HM, Mukherjee B, et al. 2013. Associations between brominated flame retardants in house dust and hormone levels in men. *Sci Total Environ* 445-446:177-184.
- Julander A, Karlsson M, Hagstrom K, et al. 2005. Polybrominated diphenyl ethers--plasma levels and thyroid status of workers at an electronic recycling facility. *Int Arch Occup Environ Health* 78(7):584-592.
- Karmaus W, Osuch JR, Landgraf J, et al. 2011. Prenatal and concurrent exposure to halogenated organic compounds and gene expression of CYP17A1, CYP19A1, and oestrogen receptor α and β genes. *Occup Environ Med* 68(6):430-437.
- Karpeta A, Barc J, Ptak A, et al. 2013. The 2,2',4,4'-tetrabromodiphenyl ether hydroxylated metabolites 5-OH-BDE-47 and 6-OH-BDE-47 stimulate estradiol secretion in the ovary by activating aromatase expression. *Toxicology* 305:65-70.
- Karpeta A, Rak-Mardyla A, Jerzak J, et al. 2011. Congener-specific action of PBDEs on steroid secretion, CYP17, 17 β -HSD and CYP19 activity and protein expression in porcine ovarian follicles. *Toxicol Lett* 206(3):258-263.
- Karpeta A, Ptak A, Gregoraszczyk EL. 2014. Different action of 2,2',4,4'-tetrabromodiphenyl ether (BDE-47) and its hydroxylated metabolites on ER α and ER β gene and protein expression. *Toxicol Lett* 229(1):250-256. 10.1016/j.toxlet.2014.05.022.
- Kawashiro Y, Fukata H, Omori-Inoue M, et al. 2008. Perinatal exposure to brominated flame retardants and polychlorinated biphenyls in Japan. *Endocr J* 55(6):1071-1084.
- Kearns GL, Abdel-Rahman SM, Alander SW, et al. 2003. Developmental pharmacology--drug disposition, action, and therapy in infants and children. *N Engl J Med* 349(12):1157-1167.
- Kicinski M, Viaene MK, Den Hond E, et al. 2012. Neurobehavioral function and low-level exposure to brominated flame retardants in adolescents: A cross-sectional study. *Environ Health* 11:86. <http://www.ehjournal.net/content/11/1/86>. October 9, 2014.
- Kierkegaard A, Bignert A, Sellstrom, et al. 2004. Polybrominated diphenyl ethers (PBDEs) and their methoxylated derivatives in pike from Swedish waters with emphasis on temporal trends, 1967-2000. *Environ Pollut* 130(2):187-198.
- Kim KH, Bose DD, Ghogha A, et al. 2011c. Para- and ortho-substitutions are key determinants of polybrominated diphenyl ether activity toward ryanodine receptors and neurotoxicity. *Environ Health Perspect* 119(4):519-526.
- Kim M, Guerra P, Theocharides M, et al. 2013b. Polybrominated diphenyl ethers in sewage sludge and treated biosolids: Effect factors and mass balance. *Water Res* 47:6496-6505.
- Kim S, Park J, Kim HJ, et al. 2013a. Association between several persistent organic pollutants and thyroid hormone levels in serum among the pregnant women of Korea. *Environ Int* 59:442-448.

9. REFERENCES

- Kim S, Park J, Kim HJ, et al. 2015. Association between several persistent organic pollutants and thyroid hormone levels in cord blood serum and bloodspot of the newborn infants of Korea. *PLoS ONE* 10(5):e0125213. 10.1371/journal.pone.0125213.
- Kim TH, Bang du Y, Lim HJ, et al. 2012a. Comparisons of polybrominated diphenyl ethers levels in paired South Korean cord blood, maternal blood, and breast milk samples. *Chemosphere* 87(1):97-104.
- Kim TS, Kim CY, Lee HK, et al. 2011b. Estrogenic activity of persistent organic pollutants and parabens based on the stably transfected human estrogen receptor- α transcriptional activation assay (OECD TG 455). *Toxicol Res* 27(3):181-184.
- Kim UJ, Kim MY, Hong YH, et al. 2012b. Assessment of impact of internal exposure to PBDEs on human thyroid function-comparison between congenital hypothyroidism and normal paired blood. *Environ Sci Technol* 46(11):6261-6268.
- Kim UJ, Lee IS, Kim HS, et al. 2011a. Monitoring of PBDEs concentration in umbilical cord blood and breast milk from Korean population and estimating the effects of various parameters on accumulation in humans. *Chemosphere* 85(3):487-493.
- Kim UJ, Sim WJ, Hong Y-H, et al. 2011d. Monitoring of brominated flame retardants in blood pair serum, Korea - focused to compare between normal and metabolic diseased group. *Organohalogen Compounds* 73:587-590.
- Kimbrough RD, Korver MP, Burse VW, et al. 1980. The effect of different diets or mineral oil on liver pathology and polybrominated biphenyl concentration in tissues. *Toxicol Appl Pharmacol* 52:442-453.
- Klasson-Wehler EK, Hovander L, Bergman A. 1997. New organohalogen in human plasma-Identification and quantification. *Organohalogen Compounds* 33:420-425.
- Klasson Wehler E, Morck A, Hakk H. 2001. Metabolism of polybrominated diphenyl ethers in the rat. The second international workshop on brominated flame retardants. *BFR 2001*. Stockholm, Sweden, 93-97.
- Klosterhaus SL, Stapleton HM, La Guardia MJ, et al. 2012. Brominated and chlorinated flame retardants in San Francisco Bay sediments and wildlife. *Environ Int* 47:56-65.
- Kociba RJ, Frauson LO, Humiston CG, et al. 1975. Results of a two-year dietary feeding study with decabromodiphenyl oxide (DBDPO) in rats. *J Combust Toxicol* 2(4):267-285.
- Kodavanti PRS. 2003. Differential effects of polybrominated diphenyl ethers and polychlorinated biphenyls on intracellular signaling in rat neuronal cultures. *Organohalogen Compounds* 65:1-4.
- Kodavanti PRS, Derr-Yellin E. 2001. Differential effects of polybrominated diphenyl ethers and polychlorinated biphenyls on [3 H]arachidonic acid release in rat neural cells. *Organohalogen Compounds* 53:185-189.
- Kodavanti PRS, Derr-Yellin EC. 2002. Differential effects of polybrominated diphenyl ethers and polychlorinated biphenyls on [3 H]arachidonic acid release in rat cerebellar granule neurons. *Toxicol Sci* 68:452-457.

9. REFERENCES

- Kodavanti PR, Coburn CG, Moser VC, et al. 2010. Developmental exposure to a commercial PBDE mixture, DE-71: Neurobehavioral, hormonal, and reproductive effects. *Toxicol Sci* 116(1):297-312.
- Kodavanti PR, Royland JE, Osorio C, et al. 2015. Developmental exposure to a commercial PBDE mixture: Effects on protein networks in the cerebellum and hippocampus of rats. *Environ Health Perspect* 123(5):428-436. 10.1289/ehp.1408504.
- Koenig CM, Lango J, Pessah IN, et al. 2012. Maternal transfer of BDE-47 to offspring and neurobehavioral development in C57BL/6J mice. *Neurotoxicol Teratol* 34(6):571-580. 10.1016/j.ntt.2012.09.005.
- Kohler M, Zennegg M, Gerecke AC, et al. 2003. Increasing concentrations of decabromodiphenyl ether (decaBDE) in Swiss sewage sludge since 1993. *Organohalogen Compounds* 61:123-126.
- Kojima H, Takeuchi S, Uramaru N, et al. 2009. Nuclear hormone receptor activity of polybrominated diphenyl ethers and their hydroxylated and methoxylated metabolites in transactivation assays using Chinese hamster ovary cells. *Environ Health Perspect* 117(8):1210-1218.
- Komori M, Nishio K, Kitada M, et al. 1990. Fetus-specific expression of a form of cytochrome P-450 in human livers. *Biochemistry* 29(18):4430-4433.
- Koskenniemi JJ, Virtanen HE, Kiviranta H, et al. 2015. Association between levels of persistent organic pollutants in adipose tissue and cryptorchidism in early childhood: A case-control study. *Environ Health* 14:78. 10.1186/s12940-015-0065-0.
- Krishnan K, Andersen ME. 1994. Physiologically based pharmacokinetic modeling in toxicology. In: Hayes AW, ed. *Principles and methods of toxicology*. 3rd ed. New York, NY: Raven Press, Ltd., 149-188.
- Krishnan K, Andersen ME, Clewell HJ III, et al. 1994. Physiologically based pharmacokinetic modeling of chemical mixtures. In: Yang RSH, ed. *Toxicology of chemical mixtures: Case studies, mechanisms, and novel approaches*. San Diego, CA: Academic Press, 399-437.
- Kucharska A, Covaci A, Vanermen G, et al. 2015. Non-invasive biomonitoring for PFRs and PBDEs: New insights in analysis of human hair externally exposed to selected flame retardants. *Sci Total Environ* 505:1062-1071. 10.1016/j.scitotenv.2014.10.043.
- Kuehl DW, Haebler R. 1995. Organochlorine, organobromine, metal and selenium residues in bottlenose dolphins (*Tursiops truncatus*) collected during an unusual mortality event in the Gulf of Mexico. *Arch Environ Contam Toxicol* 28(4):494-499.
- Kuehl DW, Haebler R, Potter C. 1991. Chemical residues in dolphins from the U.S. Atlantic coast including Atlantic bottlenose obtained during the 1987/88 mass mortality. *Chemosphere* 22(11):1071-1084.
- Kumar J, Lind PM, Salihovic S, et al. 2014a. Influence of persistent organic pollutants on the complement system in a population-based human sample. *Environ Int* 71:94-100. 10.1016/j.envint.2014.06.009.

9. REFERENCES

- Kumar J, Lind PM, Salihovic S, et al. 2014b. Persistent organic pollutants and inflammatory markers in a cross-sectional study of elderly Swedish people: The PIVUS cohort. *Environ Health Perspect* 122(9):977-983. 10.1289/ehp.1307613.
- Kuosmanen K, Hyotylainen T, Hartonen K, et al. 2002. Pressurized hot water extraction coupled on-line with liquid chromatography-gas chromatography for the determination of brominated flame retardants in sediment samples. *J Chromatogr A* 943(1):113-122.
- Kuriyama S, Chahoud I. 2003. Maternal exposure to low dose 2,2',4,4',5 pentabromo diphenyl ether (PBDE 99) impairs male reproductive performance in adult rat offspring. *Organohalogen Compounds* 61:92-95.
- Kuriyama SN, Talsness CE, Grote K, et al. 2005. Developmental exposure to low dose PBDE 99: Effects on male fertility and neurobehavior in rat offspring. *Environ Health Perspect* 113(2):149-154.
- Kuriyama SN, Wanner A, Fidalgo-Neto AA, et al. 2007. Developmental exposure to low-dose PBDE-99: Tissue distribution and thyroid hormone levels. *Toxicology* 242(1-3):80-90.
- Kwan CS, Takada H, Mizukawa K, et al. 2013. PBDEs in leachates from municipal solid waste dumping sites in tropical Asian countries: Phase distribution and debromination. *Environ Sci Pollut Res Int* 20(6):4188-4204.
- Kwiecińska P, Wrobel A, Gregoraszczyk EL. 2011. Combinatory effects of PBDEs and 17 β -estradiol on MCF-7 cell proliferation and apoptosis. *Pharmacol Rep* 63(1):189-194.
- Lacorte S, Ikononou MG. 2009. Occurrence and congener specific profiles of polybrominated diphenyl ethers and their hydroxylated and methoxylated derivatives in breast milk from Catalonia. *Chemosphere* 74(3):412-420.
- Ladenson P, Singer P, Ain K, et al. 2000. American Thyroid Association guidelines for detection of thyroid dysfunction. *Arch Intern Med* 160:1573-1575.
- LaFranchi S. 1999. Congenital hypothyroidism: Etiologies, diagnosis, and management. *Thyroid* 9(7):735-740.
- La Guardia MJ, Hale RC, Harvey E, et al. 2000. Endocrine disruptors (octylphenol, nonylphenol, nonyl phenol ethoxylates and polybrominated diphenyl ethers) in land applied sewage sludge biosolids. In: Preprints of extended abstracts. American Chemical Society, Division of Environmental Chemistry. 220th ACS National Meeting, Washington, DC. Vol. 40(2):97-99.
- La Guardia MJ, Hale RC, Harvey E. 2006. Detailed polybrominated diphenyl ether (PBDE) congener composition of the widely used penta-, octa-, and deca-PBDE technical flame-retardant mixtures. *Environ Sci Technol* 40(20):6247-6254.
- Lai Y, Cai Z. 2012. *In vitro* metabolism of hydroxylated polybrominated diphenyl ethers and their inhibitory effects on 17 β -estradiol metabolism in rat liver microsomes. *Environ Sci Pollut Res Int* 19(8):3219-3227.
- Lai Y, Lu M, Gao X, et al. 2011. New evidence for toxicity of polybrominated diphenyl ethers: DNA adduct formation from quinone metabolites. *Environ Sci Technol* 45(24):10720-10727.

9. REFERENCES

- Laird BD, Goncharov AB, Chan HM. 2013. Body burden of metals and persistent organic pollutants among Inuit in the Canadian Arctic. *Environ Int* 59:33-40.
- LaKind JS, Berlin CM. 2000. PBDEs in breast milk: Where do we go from here? *Organohalogen Compounds* 47:241-244.
- LaKind JS, Berlin CM, Jr., Sjodin A, et al. 2009. Do human milk concentrations of persistent organic chemicals really decline during lactation? *Chemical concentrations during lactation and milk/serum partitioning. Environ Health Perspect* 117(10):1625-1631.
- Landrigan PJ, Wilcox KR, Silva J. 1979. Cohort study of Michigan residents exposed to polybrominated biphenyls: Epidemiologic and immunologic findings. *Ann NY Acad Sci* 320:284-294.
- Law RJ, Allchin CR, Bennett ME, et al. 2000. Polybrominated diphenyl ethers in the blubber of harbour porpoises (*Phocoena phocoena L.*) stranded on the coasts of England and Wales. *Organohalogen Compounds* 47:249-252.
- Law RJ, Herzke D, Harrad S, et al. 2008. Levels and trends of HBCD and BDEs in the European and Asian environments, with some information for other BFRs. *Chemosphere* 73(2):223-241.
- Lee DH, Lind PM, Jacobs DR, Jr., et al. 2011. Polychlorinated biphenyls and organochlorine pesticides in plasma predict development of type 2 diabetes in the elderly. The Prospective Investigation of the Vasculature in Uppsala Seniors (PIVUS) study. *Diabetes Care* 34(8):1778-1784. 10.2337/dc10-2116.
- Lee DH, Lind PM, Jacobs DR, Jr., et al. 2012. Background exposure to persistent organic pollutants predicts stroke in the elderly. *Environ Int* 47:115-120.
- Lee E, Kim TH, Choi JS, et al. 2010. Evaluation of liver and thyroid toxicity in Sprague-Dawley rats after exposure to polybrominated diphenyl ether BDE-209. *J Toxicol Sci* 35(4):535-545.
- Leeder JS, Kearns GL. 1997. Pharmacogenetics in pediatrics: Implications for practice. *Pediatr Clin North Am* 44(1):55-77.
- Leijds MM, Koppe JG, Olie K, et al. 2009. Effects of dioxins, PCBs, and PBDEs on immunology and hematology in adolescents. *Environ Sci Technol* 43(20):7946-7951.
- Leijds MM, ten Tusscher GW, Olie K, et al. 2012. Thyroid hormone metabolism and environmental chemical exposure. *Environ Health* 11(Suppl 1):S10. <http://www.ehjournal.net/content/11/S1/S10>. October 9, 2014.
- Leijds MM, van Teunenbroek T, Olie K, et al. 2008. Assessment of current serum levels of PCDD/Fs, dl-PCBs and PBDEs in a Dutch cohort with known perinatal PCDD/F exposure. *Chemosphere* 73(2):176-181.
- Leikin JB, Pauloucek FP. 2008. Polychlorinated biphenyls. In: Leikin JB, Pauloucek FP, eds. *Poisoning and toxicology handbook*. 4th ed. Boca Raton, FL: CRC Press, Taylor & Francis Group, 840.
- Lemesh KA. 1992. Polychlorinated biphenyls: An overview of metabolic toxicologic and health consequences. *Vet Hum Toxicol* 34:256-268.

9. REFERENCES

- Lenoir D, Zier B, Bieniek D, et al. 1994. The influence of water and metals on PBDD/F concentration in incineration of decabromobiphenyl ether in polymeric matrices. *Chemosphere* 28(11):1921-1928.
- Leonetti C, Butt CM, Hoffman K, et al. 2016. Concentrations of polybrominated diphenyl ethers (PBDEs) and 2,4,6-tribromophenol in human placental tissues. *Environ Int* 88:23-29. 10.1016/j.envint.2015.12.002.
- Lepom P, Berndt M, Duffek A, et al. 2010. Oral bioavailability of PBDEs in dust using and *in vitro* gastrointestinal model. 5th International Symposium on brominated flame retardants (BFR2010), Kyoto, 7-9 April 2010. http://www.researchgate.net/publication/239593679_Oral_Bioaccessibility_of_PBDEs_in_Dust_Using_a_n_In_Vitro_Gastrointestinal_Model. March 25, 2015.
- Leung HW. 1993. Physiologically-based pharmacokinetic modeling. In: Ballantine B, Marrs T, Turner P, eds. *General and applied toxicology*. Vol. 1. New York, NY: Stockton Press, 153-164.
- Li A, Rockne KJ, Sturchio N, et al. 2006. Polybrominated diphenyl ethers in the sediments of the Great Lakes. 4. Influencing factors, trends, and implications. *Environ Sci Technol* 40(24):7528-7534.
- Li LX, Chen L, Meng XZ, et al. 2013a. Exposure levels of environmental endocrine disruptors in mother-newborn pairs in China and their placental transfer characteristics. *PLoS ONE* 8(5):e62526. <http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0062526>. October 9, 2014.
- Li X, Gao Y, Guo LH, et al. 2013b. Structure-dependent activities of hydroxylated polybrominated diphenyl ethers on human estrogen receptor. *Toxicology* 309:15-22.
- Liang SX, Gao HX, Zhao YY, et al. 2010. Effects of repeated exposure to decabrominated diphenyl ether (BDE-209) on mice nervous system and its self repair. *Environ Toxicol Pharmacol* 29(3):297-301.
- Life Sciences Research Israel Ltd. 1987. FR-1208: Teratology study in the rat. Submitted to the U.S. Environmental Protection Agency under TSCA Section 4/8. OTS0513908.
- Lignell S, Aune M, Darnerud PO, et al. 2016. Maternal body burdens of PCDD/Fs and PBDEs are associated with maternal serum levels of thyroid hormones in early pregnancy: A cross-sectional study. *Environ Health* 15(1):55. 10.1186/s12940-016-0139-7.
- Lignell S, Aune M, Darnerud PO, et al. 2013. Prenatal exposure to polychlorinated biphenyls (PCBs) and polychlorinated diphenyl ethers (PBDEs) may influence birth weight among infants in a Swedish cohort with background exposure: A cross-sectional study. *Environ Health* 12:44.
- Lim JS, Lee DH, Jacobs DR, Jr. 2008. Association of brominated flame retardants with diabetes and metabolic syndrome in the U.S. population, 2003-2004. *Diabetes Care* 31(9):1802-1807.
- Lin SM, Chen FA, Huang YF, et al. 2011. Negative associations between PBDE levels and thyroid hormones in cord blood. *Int J Hyg Environ Health* 214(2):115-120.
- Lind PM, van Bavel B, Salihovic S, et al. 2012. Circulating levels of persistent organic pollutants (POPs) and carotid atherosclerosis in the elderly. *Environ Health Perspect* 120(1):38-43.
- Lind Y, Darnerud PO, Atuma S, et al. 2003. Polybrominated diphenyl ethers in breast milk from Uppsala County, Sweden. *Environ Res* 93(2):186-194.

9. REFERENCES

- Lindstöm GUM. 1999. Aspects of polybrominated diphenyl ethers as indoor, occupational, and environmental pollutants. *Organohalogen Compounds* 43:445-446.
- Lindström G, Hardell L, Van Bavel B, et al. 1998. Current level of 2,2',4,4'-tetrabrominated diphenyl ether in human adipose tissue in Sweden. A risk factor for non-Hodgkin's lymphoma? *Organohalogen Compounds* 35:431-434.
- Lindström G, Wingfors H, Dam M, et al. 1999. Identification of 19 polybrominated diphenyl ethers (PBDEs) in long-finned pilot whale (*Globicephala melas*) from the Atlantic. *Arch Environ Contam Toxicol* 36(3):355-363.
- Lioy PJ. 2002. Residues from the world trade center disaster in lower Manhattan and potential human exposures. *Int J Toxicol* 21(6):540.
- Liu L, Salamova A, Hites RA. 2014. Halogenated flame retardants in baby food from the United States and from China and the estimated dietary intakes by infants. *Environ Sci Technol* 48(16):9812-9818.
- Liu LY, He K, Hites RA, et al. 2016. Hair and nails as noninvasive biomarkers of human exposure to brominated and organophosphate flame retardants. *Environ Sci Technol* 50(6):3065-3073. 10.1021/acs.est.5b05073.
- Liu X, Wang J, Lu C, et al. 2015. The role of lysosomes in BDE 47-mediated activation of mitochondrial apoptotic pathway in HepG2 cells. *Chemosphere* 124:10-21. 10.1016/j.chemosphere.2014.10.054.
- Liu X, Zhan H, Zeng X, et al. 2012. The PBDE-209 exposure during pregnancy and lactation impairs immune function in rats. *Mediators Inflamm* 2012:692467.
- Livingston AL. 1978. Forage plant estrogens. *J Toxicol Environ Health* 4(2-3):301-324.
- Loganathan BG, Kannn K, Watanabe, et al. 1995. Isomer-specific determination and toxic evaluation of polychlorinated biphenyls, polychlorinated/brominated dibenzo-p-dioxins and dibenzofurans, polybrominated biphenyl ethers, and extractable organic halogen in carp from the Buffalo River, New York. *Environ Sci Technol* 29(7):1832-1838.
- Lopez-Espinosa MJ, Costa O, Vizcaino E, et al. 2015. Prenatal exposure to polybrominated flame retardants and fetal growth in the INMA Cohort (Spain). *Environ Sci Technol* 49(16):10108-10116. 10.1021/acs.est.5b01793.
- Lorber M. 2008. Exposure of Americans to polybrominated diphenyl ethers. *J Expo Sci Environ Epidemiol* 18(1):2-19.
- Luckey F, Fowler B, Litten S. 2001. Establishing baseline levels of polybrominated diphenyl ethers in Lake Ontario surface waters. The second international workshop on brominated flame retardants. *BFR* 2001. Stockholm, Sweden, 337-339.
- Lundgren M, Darnerud PO, Blomberg J, et al. 2009. Polybrominated diphenyl ether exposure suppresses cytokines important in the defence to coxsackievirus B3 infection in mice. *Toxicol Lett* 184(2):107-113.

9. REFERENCES

- Lupton SJ, McGarrigle BP, Olson JR, et al. 2009. Human liver microsome-mediated metabolism of brominated diphenyl ethers 47, 99, and 153 and identification of their major metabolites. *Chem Res Toxicol* 22(11):1802-1809.
- Luross JM, Aalee M, Sergeant DB, et al. 2000. Spatial and temporal distribution of polybrominated diphenyl ethers in lake trout from the Great Lakes. *Organohalogen Compounds* 47:73-76.
- Luross JM, Alae M, Sergeant DB, et al. 2002. Spatial distribution of polybrominated diphenyl ethers and polybrominated biphenyls in lake trout from the Laurentian Great Lakes. *Chemosphere* 46(5):665-672.
- Lyman WJ, Reehl WF, Rosenblatt, DH. 1990. Handbook of chemical properties estimation methods. American Chemical Society: Washington, DC, 4-9, 15-1 to 15-29.
- Ma J, Qiu X, Zhang J, et al. 2012a. State of polybrominated diphenyl ethers in China: An overview. *Chemosphere* 88(7):769-778.
- Ma J, Qiu X, Ren A, et al. 2012b. Using placenta to evaluate the polychlorinated biphenyls (PCBs) and polybrominated diphenyl ethers (PBDEs) exposure of fetus in a region with high prevalence of neural tube defects. *Ecotoxicol Environ Saf* 86:141-146. 10.1016/j.ecoenv.2012.09.005.
- MacPhail R, Farmer JD, Padnos BK, et al. 2003. Lack of effect of perinatal exposure to a polybrominated diphenyl ether mixture (DE-71) on the habituation of motor activity in adult rats. *Toxicol Sci* 72(S-1):123.
- Main KM, Kiviranta H, Virtanen HE, et al. 2007. Flame retardants in placenta and breast milk and cryptorchidism in newborn boys. *Environ Health Perspect* 115(10):1519-1526.
- Makey CM, McClean MD, Braverman LE, et al. 2016. Polybrominated diphenyl ether exposure and thyroid function tests in North American adults. *Environ Health Perspect* 124(4):420-425. 10.1289/ehp.1509755.
- Malarvannan G, Isobe T, Covaci A, et al. 2013. Accumulation of brominated flame retardants and polychlorinated biphenyls in human breast milk and scalp hair from the Philippines: Levels, distribution and profiles. *Sci Total Environ* 442:366-379.
- Malmberg T, Athanasiadou M, Marsh G, et al. 2005. Identification of hydroxylated polybrominated diphenyl ether metabolites in blood plasma from polybrominated diphenyl ether exposed rats. *Environ Sci Technol* 39(14):5342-5348.
- Manchester-Neesvig JB, Valters K, Sonzogni WC. 2001. Comparison of polybrominated diphenyl ethers (PBDEs) and polychlorinated biphenyls (PCBs) in Lake Michigan salmonids. *Environ Sci Technol* 35(6):1072-1077.
- Maranghi F, Tassinari R, Moracci G, et al. 2013. Dietary exposure of juvenile female mice to polyhalogenated seafood contaminants (HBCD, BDE-47, PCB-153, TCDD): Comparative assessment of effects in potential target tissues. *Food Chem Toxicol* 56:443-449.
- Marchesini GR, Meimaridou A, Haasnoot W, et al. 2008. Biosensor discovery of thyroxine transport disrupting chemicals. *Toxicol Appl Pharmacol* 232(1):150-160. 10.1016/j.taap.2008.06.014.

9. REFERENCES

- Marchitti SA, LaKind JS, Naiman DQ, et al. 2013. Improving infant exposure and health risk estimates: Using serum data to predict polybrominated diphenyl ether concentrations in breast milk. *Environ Sci Technol* 47(9):4787-4795.
- Mariussen E, Fonnum F. 2002. The effect of pentabromodiphenyl ether, hexabromocyclododecane and tetrabromobisphenol-A on dopamine uptake into rat brain synaptosomes. *Organohalogen Compounds* 57:395-398.
- Mariussen E, Fonnum F. 2003. The effect of brominated flame retardants on neurotransmitter uptake into rat brain synaptosomes and vesicles. *Neurochem Int* 43(4-5):533-542.
- Mariussen E, Andersson PL, Olstrom H, et al. 2003. The effect of various substituents in ortho position of biphenyls on respiratory burst, intracellular calcium elevation in human granulocytes, and uptake of dopamine into rat synaptic vesicles and synaptosomes. *Environ Toxicol Pharmacol* 14(1-2):43-50.
- Markowski VP. 2007. Thyroid disruption and behavioral impairments following developmental exposure to PBDEs: Correlative or causative relationship? *Neurotoxicol Teratol* 29(3):413.
- Marsh G, Athanasiadou M, Athanassiadis I, et al. 2006. Identification of hydroxylated metabolites in 2,2',4,4'-tetrabromodiphenyl ether exposed rats. *Chemosphere* 63(4):690-697.
- Marsh G, Bergman A, Bladh L-G, et al. 1998. Synthesis of p-hydroxybromodiphenyl ethers and binding to the thyroid receptor. *Organohalogen Compounds* 37:305-308.
- Martin PA, Mayne GJ, Bursian FS, et al. 2007. Immunotoxicity of the commercial polybrominated diphenyl ether mixture DE-71 in ranch mink (*Mustela vison*). *Environ Toxicol Chem* 26(5):988-997.
- Mayr U, Butsch A, Schneider S. 1992. Validation of two *in vitro* test systems for estrogenic activities with zearalenone, phytoestrogens and cereal extracts. *Toxicology* 74(2-3):135-149.
- Mazdai A, Dodder NG, Abernathy MP, et al. 2003. Polybrominated diphenyl ethers in maternal and fetal blood samples. *Environ Health Perspect* 111:1249-1252.
- McAllister D, Mazac CJ, Gorsich R, et al. 1990. Analysis of polymers containing brominated diphenyl ethers as flame retardants after molding under various conditions. *Chemosphere* 10-12:1537-1541.
- McConnell EE, Harris MW, Moore JA. 1980. Studies on the use of activated charcoal and cholestyramine for reducing the body burden of polybrominated biphenyls. *Drug Chem Toxicol* 3:277-292.
- McDonald TA. 2002. A perspective on the potential health risks of PBDEs. *Chemosphere* 46(5):745-755.
- McIntyre RL, Kenerson HL, Subramanian S, et al. 2015. Polybrominated diphenyl ether congener, BDE-47, impairs insulin sensitivity in mice with liver-specific Pten deficiency. *BMC obesity* 2:3. 10.1186/s40608-014-0031-3.
- McKinney RF, Chaw SJK, Rickenbacher U, et al. 1987. Polychlorinated biphenyls and related compound interactions with specific binding sites for thyroxine in rat liver nuclear extracts. *J Med Chem* 30:79-86.

9. REFERENCES

- Meeker JD, Johnson PI, Camann D, et al. 2009. Polybrominated diphenyl ether (PBDE) concentrations in house dust are related to hormone levels in men. *Sci Total Environ* 407(10):3425-3429.
- Meerts IA, Marsh G, van Leeuwen-Bol I, et al. 1998. Interaction of polybrominated diphenyl ether metabolites (PBDE-OH) with human transthyretin *in vitro*. *Organohalogen Compounds* 37:309-312.
- Meerts IA, Letcher RJ, Hoving S, et al. 2001. *In vitro* estrogenicity of polybrominated diphenyl ethers, hydroxylated PBDEs and polybrominated bisphenol A compounds. *Environ Health Perspect* 109(4):399-407.
- Meerts IA, van Zanden JJ, Luijks EA, et al. 2000. Potent competitive interactions of some brominated flame retardants and related compounds with human transthyretin *in vitro*. *Toxicol Sci* 56(1):95-104.
- Meijer L, Martijn A, Melessen J, et al. 2012. Influence of prenatal organohalogen levels on infant male sexual development: Sex hormone levels, testes volume and penile length. *Hum Reprod* 27(3):867-872.
- Meijer L, Weiss J, Van Velzen M, et al. 2008. Serum concentrations of neutral and phenolic organohalogenes in pregnant women and some of their infants in The Netherlands. *Environ Sci Technol* 42(9):3428-3433.
- Meironyté Guvenius D, Noren K. 1999. Polybrominated diphenyl ethers in human liver and adipose tissues. A pilot study. *Organohalogen Compounds* 40:372-382.
- Meironyté Guvenius D, Noren K. 2001. Polybrominated diphenyl ethers in Swedish human milk. The follow-up study. The second international workshop on brominated flame retardants. BFR 2001. Stockholm, Sweden, 303-305.
- Meironyté Guvenius D, Bergman A, Noren K. 2001. Polybrominated diphenyl ethers in Swedish human liver and adipose tissue. *Arch Environ Contam Toxicol* 40:564-570.
- Meironyté D, Noren K, Bergman A. 1999. Analysis of polybrominated diphenyl ethers in Swedish human milk. A time related trend study, 1972-1997. *J Toxicol Environ Health A* 58(6):329-341.
- Meneses M, Wingfors H, Schuhmacher M, et al. 1999. Polybrominated diphenyl ethers detected in human adipose tissue from Spain. *Chemosphere* 39(13):2271-2278.
- Meng G, Feng Y, Nie Z, et al. 2016. Internal exposure levels of typical POPs and their associations with childhood asthma in Shanghai, China. *Environ Res* 146:125-135. 10.1016/j.envres.2015.12.026.
- Mercado-Feliciano M, Bigsby RM. 2008a. The polybrominated diphenyl ether mixture DE-71 is mildly estrogenic. *Environ Health Perspect* 116(5):605-611.
- Mercado-Feliciano M, Bigsby RM. 2008b. Hydroxylated metabolites of the polybrominated diphenyl ether mixture DE-71 are weak estrogen receptor- α ligands. *Environ Health Perspect* 116(10):1315-1321.
- Messer A. 2010. Mini-review: Polybrominated diphenyl ether (PBDE) flame retardants as potential autism risk factors. *Physiol Behav* 100(3):245-249.
- Meylan WM, Howard PH. 1993. Computer estimation of the atmospheric gas-phase reaction rate of organic compounds with hydroxyl radicals and ozone. *Chemosphere* 26:2293-2299.

9. REFERENCES

- Microbiological Associates Inc. 1996. Pentabromodiphenyloxyde and octabromodiphenyl oxide. Maximization test in guinea pigs. Submitted to U.S. Environmental Protection Agency under TSCA Section FYI. OTS0001281.
- Miller VM, Sanchez-Morrissey S, Brosch KO, et al. 2012. Developmental coexposure to polychlorinated biphenyls and polybrominated diphenyl ethers has additive effects on circulating thyroxine levels in rats. *Toxicol Sci* 127(1):76-83.
- Miranda ML, Anthopoulos R, Wolkin A, et al. 2015. Associations of birth outcomes with maternal polybrominated diphenyl ethers and thyroid hormones during pregnancy. *Environ Int* 85:244-253. 10.1016/j.envint.2015.09.015.
- Morck A, Klasson Wehler E. 2001. Metabolism of decabromodiphenyl ether (BDE-209) in the rat. *Organohalogen Compounds* 52:9-12.
- Morck A, Hakk H, Orn U, et al. 2003. Decabromodiphenyl ether in the rat: Absorption, distribution, metabolism, and excretion. *Drug Metab Dispos* 31:900-907.
- Morreale de Escobar G, Obregon MJ, Escobar del Rey F. 2000. Is neuropsychological development related to maternal hypothyroidism or to maternal hypothyroxinemia? *J Clin Endocrinol Metab* 85(11):3975-3987.
- Morselli PL, Franco-Morselli R, Bossi L. 1980. Clinical pharmacokinetics in newborns and infants: Age-related differences and therapeutic implications. *Clin Pharmacokin* 5(6):485-527.
- Müller MH, Polder A, Brynildsrud OB, et al. 2016. Brominated flame retardants (BFRs) in breast milk and associated health risks to nursing infants in Northern Tanzania. *Environ Int* 89-90:38-47. 10.1016/j.envint.2015.12.032.
- Mumford SL, Kim S, Chen Z, et al. 2015. Persistent organic pollutants and semen quality: The LIFE Study. *Chemosphere* 135:427-435. 10.1016/j.chemosphere.2014.11.015.
- Nakari T, Pessala P. 2005. *In vitro* estrogenicity of polybrominated flame retardants. *Aquat Toxicol* 74(3):272-279.
- NAS/NRC. 1989. Report of the oversight committee. In: *Biologic markers in reproductive toxicology*. Washington, DC: National Academy of Sciences, National Research Council, National Academy Press, 15-35.
- Ni K, Lu Y, Wang T, et al. 2013. Polybrominated diphenyl ethers (PBDEs) in China: Policies and recommendations for sound management of plastics from electronic wastes. *J Environ Manage* 115:114-123.
- NIOSH. 2014. NIOSH pocket guide to chemical hazards. Atlanta, GA: National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention. <http://www.cdc.gov/niosh/npg/npgdcas.html>. December 19, 2014.
- Nomiyama K, Kanbara C, Ochiai M, et al. 2014. Halogenated phenolic contaminants in the blood of marine mammals from Japanese coastal waters. *Mar Environ Res* 93:15-22.

9. REFERENCES

- Norén K, Meironyté D. 1998. Contaminants in Swedish human milk. Decreasing levels of organochlorine and increasing levels of organochlorine compounds. *Organohalogen Compounds* 38:1-4.
- Norén K, Meironyté D. 2000. Certain organochlorine and organobromine contaminants in Swedish human milk in perspective of past 20-30 years. *Chemosphere* 40:1111-1123.
- Norris JM, Ehrmantraut JW, Gibbons CL, et al. 1973. Toxicological and environmental factors involved in the selection of decabromodiphenyl oxide as a fire retardant chemical. *Appl Polym Symp* 22:195-219.
- Norris JM, Kociba RJ, Humiston CG, et al. 1975a. The toxicity of decabromo diphenyl and octabromo biphenyl as determined by subacute and chronic dietary feeding studies in rats. *Toxicol Appl Pharmacol* 33:170.
- Norris JM, Kociba RJ, Schwetz BA, et al. 1975b. Toxicology of octabromobiphenyl and decabromodiphenyl oxide. *Environ Health Perspect* 11:153-161.
- Norstrom RJ, Simon M, Moisey J, et al. 2002. Geographical distribution (2000) and temporal trends (1981-2000) of brominated diphenyl ethers in Great Lakes herring gull eggs. *Environ Sci Technol* 36:4783-4789.
- NRC. 1993. Pesticides in the diets of infants and children. National Research Council. Washington, DC: National Academy Press.
- NRC. 2000. Decabromodiphenyl oxide. In: Toxicological risks of selected flame-retardant chemicals. National Research Council (US): Subcommittee on Flame-Retardant Chemicals. Washington, DC: National Academy Press, 72-98.
- NRC. 2002. Biosolids applied to land: Advancing standards and practices. Washington, DC: National Research Council, National Academy of Sciences, National Academy Press. <http://www.nap.edu/catalog/10426.html>. September 21, 2016.
- NTP. 1983. NTP technical report on the toxicology and carcinogenesis studies of a polybrominated biphenyl mixture (Firemaster FF-1) (CAS No. 67774-32-7) in F344/N rats and B6C3F1 mice (gavage studies). Research Triangle Park, NC: National Toxicology Program.
- NTP. 1986. NTP technical report on the toxicology and carcinogenesis studies of decabromodiphenyl oxide (CAS No. 1163-19-5) in F344/N rats and B6C3F1 mice (feed studies). Research Triangle Park, NC: National Toxicology Program.
- NTP. 2011. Report on carcinogens. 12th ed. Research Triangle Park, NC: U.S. Department of Health and Human Services, Public Health Service, National Toxicology Program. <http://ntp-server.niehs.nih.gov/ntp/roc/twelfth/roc12.pdf>. September 10, 2014.
- Nylund K, Asplund L, Jansson B, et al. 1992. Analysis of some polyhalogenated organic pollutants in sediment and sewage sludge. *Chemosphere* 24(12):1721-1730.
- Oberg T, Warman K, Bergstrom J. 1987. Brominated aromatics from combustion. *Chemosphere* 16(10-12):2451-2465.

9. REFERENCES

- Oberg M, Westerholm E, Fattore E, et al. 2010. Toxicity of Bromkal 70-5DE, a technical mixture of polybrominated diphenyl ethers, following 28 d of oral exposure in rats and impact of analysed impurities. *Chemosphere* 80(2):137-143.
- Ochiai M, Nomiyama K, Isobe T, et al. 2013. Accumulation of hydroxylated polychlorinated biphenyls (OH-PCBs) and implications for PCBs metabolic capacities in three porpoise species. *Chemosphere* 92:803-810.
- Ochiai S, Shimojo N, Yuka I, et al. 2014. A pilot study for foetal exposure to multiple persistent organic pollutants and the development of infant atopic dermatitis in modern Japanese society. *Chemosphere* 94:48-52.
- O'Connor JC, Frame SR, Davis LG, et al. 1999. Detection of thyroid toxicants in a tier 1 screening battery and alterations in thyroid endpoints over 28 days of exposure. *Toxicol Sci* 51:54-70.
- Oduanya DO, Okonkwo JO, Botha B. 2009. Polybrominated diphenyl ethers (PBDEs) in leachates from selected landfill sites in South Africa. *Waste Manag* 29(1):96-102.
- Ohta S, Ishizuka D, Nishimura H, et al. 2000. Real situation of contamination by polybrominated diphenyl ethers as flame retardants in market fish and mother milk of Japan. *Organohalogen Compounds* 47:218-221.
- Ohta S, Ishizuka D, Nishimura H, et al. 2002. Comparison of polybrominated diphenyl ethers in fish, vegetables, and meats and levels in human milk of nursing women in Japan. *Chemosphere* 46(5):689-696.
- Oliaei F, Weber R, Watson A. 2010. PBDE contamination in Minnesota landfills, waste water treatment plants and sediments as PBDE sources and reservoirs. *Organohalogen Compounds* 72:1346-1349.
- Olsson A, Vitnish M, Plikshs M, et al. 1999. Halogenated environmental contaminants in perch (*Perca fluviatilis*) from Latvian coastal areas. *Sci Total Environ* 239(1-3):19-30.
- Örn U, Klasson-Wehler E. 1998. Metabolism of 2,2',4,4'-tetrabromodiphenyl ether in rat and mouse. *Xenobiotica* 1998:199-211.
- Oros DR, Hoover D, Rodigari F, et al. 2005. Levels and distribution of polybrominated diphenyl ethers in water, surface sediments, and bivalves from the San Francisco Estuary. *Environ Sci Technol* 39(1):33-41.
- OSHA. 2013a. Air contaminants. Table Z-1-limits for air contaminants. Occupational safety and health standards. Code of Federal Regulations 29 CFR 1910.1000, Subpart Z. <http://www.gpo.gov/fdsys/pkg/CFR-2013-title29-vol6/pdf/CFR-2013-title29-vol6-sec1910-1000.pdf>. September 9, 2014.
- OSHA. 2013b. Process safety management of highly hazardous chemicals. Appendix A to 1910.119-List of highly hazardous chemicals, toxics and reactives (mandatory). Occupational safety and health standards. Code of Federal Regulations 29 CFR 1910.119, Subpart H. <http://www.gpo.gov/fdsys/pkg/CFR-2013-title29-vol5/pdf/CFR-2013-title29-vol5-sec1910-119.pdf>. September 9, 2014.

9. REFERENCES

- Oulhote Y, Chevrier J, Bouchard MF. 2016. Exposure to polybrominated diphenyl ethers (PBDEs) and hypothyroidism in Canadian women. *J Clin Endocrinol Metab* 101(2):590-598. 10.1210/jc.2015-2659.
- Owen GM, Brozek J. 1966. Influence of age, sex and nutrition on body composition during childhood and adolescence. In: Falkner F, ed. *Human development*. Philadelphia, PA: WB Saunders, 222-238.
- Pacyniak E, Hagenbuch B, Klaassen CD, et al. 2011. Organic anion transporting polypeptides in the hepatic uptake of PBDE congeners in mice. *Toxicol Appl Pharmacol* 257(1):23-31.
- Pacyniak E, Roth M, Hagenbuch B, et al. 2010. Mechanism of polybrominated diphenyl ether uptake into the liver: PBDE congeners are substrates of human hepatic OATP transporters. *Toxicol Sci* 115(2):344-353.
- Palha J, Fernandes R, de Escobar G, et al. 2000. Transthyretin regulates thyroid hormone levels in the choroid plexus, but not in the brain parenchyma: Study in a transthyretin-null mouse model. *Endocrinology* 141(9):3267-3272.
- Palha J, Hays M, Morreale de Escobar G, et al. 1997. Transthyretin is not essential for thyroxine to reach the brain and other tissues in transthyretin-null mice. *Am J Physiol* 272(3 Pt 1):E485-E493.
- Park HR, Loch-Caruso R. 2015. Protective effect of (+/-)-alpha-tocopherol on brominated diphenyl ether-47-stimulated prostaglandin pathways in human extravillous trophoblasts in vitro. *Toxicol in Vitro* 29(7):1309-1318. 10.1016/j.tiv.2015.05.015.
- Park JS, She J, Holden A, et al. 2011. High postnatal exposures to polybrominated diphenyl ethers (PBDEs) and polychlorinated biphenyls (PCBs) via breast milk in California: Does BDE-209 transfer to breast milk? *Environ Sci Technol* 45(10):4579-4585.
- Parolini M, Guazzoni N, Binelli A, et al. 2012. Polybrominated diphenyl ether contamination in soil, vegetation, and cow milk from a high-mountain pasture in the Italian Alps. *Arch Environ Contam Toxicol* 63(1):29-44.
- Patterson DG, Sjodin A, Bergman A. 2000. Brominated flame retardants in serum from US blood donors. *Organohalogen Compounds* 47:45-48.
- Pazin M, Pereira LC, Dorta DJ. 2015. Toxicity of brominated flame retardants, BDE-47 and BDE-99 stems from impaired mitochondrial bioenergetics. *Toxicol Mech Methods* 25(1):34-41. 10.3109/15376516.2014.974233.
- Pellacani C, Buschini A, Galati S, et al. 2012. Evaluation of DNA damage induced by 2 polybrominated diphenyl ether flame retardants (BDE-47 and BDE-209) in SK-N-MC cells. *Int J Toxicol* 31(4):372-379.
- Peltier MR, Koo HC, Getahun D, et al. 2015. Does exposure to flame retardants increase the risk for preterm birth? *J Reprod Immunol* 107:20-25. 10.1016/j.jri.2014.11.002.
- Pereira LC, Miranda LF, de Souza AO, et al. 2014. BDE-154 induces mitochondrial permeability transition and impairs mitochondrial bioenergetics. *J Toxicol Environ Health A* 77(1-3):24-36. 10.1080/15287394.2014.861337.
- Pessah IN, Cherednichenko G, Lein PJ. 2010. Minding the calcium store: Ryanodine receptor activation as a convergent mechanism of PCB toxicity. *Pharmacol Ther* 125(2):260-285.

9. REFERENCES

- Peters AJ, Coleman P, Jones KC. 1999. Organochlorine pesticides in UK air. *Organohalogen Compounds* 41:447-450.
- Petreas M, She J, Brown FR, et al. 2002. High PBDE concentrations in California human and wildlife populations. *Organohalogen Compounds* 58:177-180.
- Petreas M, She J, Brown FR, et al. 2003. High body burdens of 2,2',4,4'-tetrabromodiphenyl ether (BDE-47) in California women. *Environ Health Perspect* 111(9):1175-1179.
- Pettersson A, Karlsson M, van Bavel B, et al. 2002. Concentrations of polybrominated diphenylethers and thyroid hormones in human plasma from exposed workers. *Organohalogen Compounds* 58:269-272.
- Pettigrew A. 1993. Flame retardants (halogenated). In: Kroschwitz JI, Howe-Grant M, eds. *Kirk-Othmer encyclopedia of chemical technology. Explosives and propellants to flame retardants for textiles*. Vol. 10. 4th edition. New York, NY: John Wiley & Sons, 960-976.
- Pharmakon Research International, Inc. 1984. Initial submission: Acute oral toxicity in rats (14 days) of Saytex 115 (pentabromodiphenyloxide). Submitted to the U.S. Environmental Protection Agency under TSCA Section FYI. OTS0000972.
- Pi N, Chia SE, Ong CN, et al. 2016. Associations of serum organohalogen levels and prostate cancer risk: Results from a case-control study in Singapore. *Chemosphere* 144:1505-1512. 10.1016/j.chemosphere.2015.10.020.
- Polin D, Leavitt RA. 1984. Colestipol and energy restriction as an approach to hasten removal of PBBs from chickens. *J Toxicol Environ Health* 13:659-671.
- Polin D, Bursian SJ, Underwood MS, et al. 1991. Elimination of PBBs in rats, effect of mineral oil and/or feed restriction. *J Toxicol Environ Health* 33:197-212.
- Polin D, Lehning E, Pullen D, et al. 1985. Procedures to enhance withdrawal of xenobiotics from chickens. *J Toxicol Environ Health* 16:243-254.
- Poma G, Binelli A, Volta P, et al. 2014. Evaluation of spatial distribution and accumulation of novel brominated flame retardants, HBCD and PBDEs in an Italian subalpine lake using zebra mussel (*Dreissena polymorpha*). *Environ Sci Pollut Res Int* 21(16):9655-9664.
- Pomerantz I, Burke J, Firestone D, et al. 1978. Chemistry of PCBs and PBBs. *Environ Health Perspect* 24:133-146.
- Poon E, Powers BE, McAlonan RM, et al. 2011. Effects of developmental exposure to polychlorinated biphenyls and/or polybrominated diphenyl ethers on cochlear function. *Toxicol Sci* 124(1):161-168.
- Poon S, Wade MG, Aleksa K, et al. 2014. Hair as a biomarker of systemic exposure to polybrominated diphenyl ethers. *Environ Sci Technol* 48(24):14650-14658. 10.1021/es502789h.
- Qiu X, Bigsby RM, Hites RA. 2009. Hydroxylated metabolites of polybrominated diphenyl ethers in human blood samples from the United States. *Environ Health Perspect* 117(1):93-98.

9. REFERENCES

- Qiu X, Marvin CH, Hites RA. 2007. Dechlorane plus and other flame retardants in a sediment core from Lake Ontario. *Environ Sci Technol* 41(17):6014-6019.
- Quanrud D, Zerzghi H, Leung C, et al. 2011. Fate of endocrine disruptors following long-term land application of Class B biosolids and risks to public health. *Proc Water Environ Fed Residuals and biosolids* 2011(4):420-435. 10.2175/193864711802863201.
- Rahman F, Lanford KH, Scrimshaw MD, et al. 2001. Polybrominated diphenyl ether (PBDE) flame retardants. *Sci Total Environ* 275(1-3):1-17.
- Rao P, Kodavanti S, Zhang P. 2003. Effects of polybrominated diphenyl ethers and polychlorinated biphenyls on [3H] phorbol ester binding in rat neurons. *J Neurochem* 85(Suppl 1):13.
- Rasinger JD, Carroll TS, Lundebye AK, et al. 2014. Cross-omics gene and protein expression profiling in juvenile female mice highlights disruption of calcium and zinc signaling in the brain following dietary exposure to CB-153, BDE-47, HBCD or TCDD. *Toxicology* 321:1-12. 10.1016/j.tox.2014.03.006.
- Rawn DF, Ryan JJ, Sadler AR, et al. 2014. Brominated flame retardant concentrations in sera from the Canadian Health Measures Survey (CHMS) from 2007 to 2009. *Environ Int* 63:26-34.
- Rayne S, Ikonomidou MG, Antcliffe B. 2003a. Rapidly increasing polybrominated diphenyl ether concentrations in the Columbia River system from 1992 to 2000. *Environ Sci Technol* 37(13):2847-2854.
- Rayne S, Ikonomidou MG, Whale MD. 2003b. Anaerobic microbial and photochemical degradation of 4,4'-dibromodiphenyl ether. *Water Res* 37:551-560.
- Reistad T, Mariussen E, Fonnum F. 2002. The effect of brominated flame retardants on cell death and free radical formation in cerebellar granule cells. *Organohalogen Compounds* 57:391-394.
- Ren XM, Guo LH. 2012. Assessment of the binding of hydroxylated polybrominated diphenyl ethers to thyroid hormone transport proteins using a site-specific fluorescence probe. *Environ Sci Technol* 46(8):4633-4640.
- Ren A, Qiu X, Jin L, et al. 2011. Association of selected persistent organic pollutants in the placenta with the risk of neural tube defects. *Proc Natl Acad Sci USA* 108(31):12770-12775.
- Ren XM, Guo LH, Gao Y, et al. 2013. Hydroxylated polybrominated diphenyl ethers exhibit different activities on thyroid hormone receptors depending on their degree of bromination. *Toxicol Appl Pharmacol* 268(3):256-263.
- RePORTER. 2014. Polybrominated diphenyl ethers (PBDE). National Institutes of Health, Research Portfolio Online Reporting Tools. <http://projectreporter.nih.gov/reporter.cfm>. August 13, 2014.
- RePORTER. 2016. Polybrominated biphenyl ethers (PBDE). National Institutes of Health, Research Portfolio Online Reporting Tools. <https://projectreporter.nih.gov/reporter.cfm>. September 21, 2016.
- Reverte I, Domingo JL, Colomina MT. 2014. Neurodevelopmental effects of decabromodiphenyl ether (BDE-209) in APOE transgenic mice. *Neurotoxicol Teratol* 46:10-17. 10.1016/j.ntt.2014.08.003.

9. REFERENCES

- Reverte I, Klein AB, Domingo JL, et al. 2013. Long term effects of murine postnatal exposure to decabromodiphenyl ether (BDE-209) on learning and memory are dependent upon APOE polymorphism and age. *Neurotoxicol Teratol* 40:17-27.
- Rice CP, Chernyak SM, Begnoche L, et al. 2002. Comparisons of PBDE composition and concentration in fish collected from the Detroit River, MI and Des Plaines River, IL. *Chemosphere* 49(7):731-737.
- Rice DC, Reeve EA, Herlihy A, et al. 2007. Developmental delays and locomotor activity in the C57BL6/J mouse following neonatal exposure to the fully-brominated PBDE, decabromodiphenyl ether. *Neurotoxicol Teratol* 29(4):511-520.
- Rice DC, Thompson WD, Reeve EA, et al. 2009. Behavioral changes in aging but not young mice after neonatal exposure to the polybrominated flame retardant decaBDE. *Environ Health Perspect* 117(12):1903-1911.
- Richardson VM, Staskal DF, Ross DG, et al. 2008. Possible mechanisms of thyroid hormone disruption in mice by BDE 47, a major polybrominated diphenyl ether congener. *Toxicol Appl Pharmacol* 226(3):244-250.
- Rickenbacher U, McKinney JD, Oatley, SJ, et al. 1986. Structurally specific binding of halogenated biphenyls to thyroxine transport protein. *J Med Chem* 29:641-648.
- Riu A, Cravedi JP, Debrauwer L, et al. 2008. Disposition and metabolic profiling of [¹⁴C]-decabromodiphenyl ether in pregnant Wistar rats. *Environ Int* 34(3):318-329.
- Roberts SC, Bianco AC, Stapleton HM. 2015. Disruption of type 2 iodothyronine deiodinase activity in cultured human glial cells by polybrominated diphenyl ethers. *Chem Res Toxicol* 28(6):1265-1274. 10.1021/acs.chemrestox.5b00072.
- Robledo CA, Yeung E, Mendola P, et al. 2015a. Preconception maternal and paternal exposure to persistent organic pollutants and birth size: The LIFE study. *Environ Health Perspect* 123(1):88-94. 10.1289/ehp.1308016.
- Robledo CA, Yeung E, Mendola P, et al. 2015b. Supplemental material: Preconception maternal and paternal exposure to persistent organic pollutants and birth size: The LIFE study. *Environ Health Perspect*. 10.1289/ehp.1308016. <http://ehp.niehs.nih.gov/wp-content/uploads/123/1/ehp.1308016.s001.508.pdf>. September 19, 2016.
- Roper CS, Simpson AG, Madden S, et al. 2006. Absorption of [¹⁴C]-tetrabromodiphenyl ether (TeBDE) through human and rat skin *in vitro*. *Drug Chem Toxicol* 29(3):289-301.
- Rosenman KD, Anderson HA, Selikoff IJ, et al. 1979. Spermatogenesis in men exposed to polybrominated biphenyl (PBB). *Fertil Steril* 32:209-213.
- Roze E, Meijer L, Bakker A, et al. 2009. Prenatal exposure to organohalogens, including brominated flame retardants, influences motor, cognitive, and behavioral performance at school age. *Environ Health Perspect* 117(12):1953-1958.
- Rozman KK, Rozman TA, Williams J, et al. 1982. Effect of mineral oil and/or cholestyramine in the diet on biliary and intestinal elimination of 2,4,5,2',4',5'-hexabromodiphenyl in the Rhesus monkey. *J Toxicol Environ Health* 9:611-618.

9. REFERENCES

- Ryan JJ, Patry B. 2000. Determination of brominated diphenyl ethers (BDE's) and levels in Canadian human milks. *Organohalogen Compounds* 47:57-60.
- Ryan JJ, Rawn DF. 2014. The brominated flame retardants, PBDEs and HBCD, in Canadian human milk samples collected from 1992 to 2005; concentrations and trends. *Environ Int* 70:1-8. 10.1016/j.envint.2014.04.020.
- Rydén A, Nestor G, Jakobsson K, et al. 2012. Synthesis and tentative identification of novel polybrominated diphenyl ether metabolites in human blood. *Chemosphere* 88(10):1227-1234.
- Saegusa Y, Fujimoto H, Woo GH, et al. 2012. Transient aberration of neuronal development in the hippocampal dentate gyrus after developmental exposure to brominated flame retardants in rats. *Arch Toxicol* 86(9):1431-1442.
- Safe S. 1984. Polychlorinated biphenyls (PCBs) and polybrominated biphenyls (PBBs): Biochemistry, toxicology, and mechanism of action. *CRC Crit Rev Toxicol* 13:319-395.
- Safe SH, Zacharewski T. 1997. Organochlorine exposure and risk for breast cancer. In: Aldaz CM, Gould MN, McLachlan J, et al., eds. *Etiology of breast and gynecological cancers*. New York, NY: Wiley-Liss Inc., 133-145.
- Sagiv SK, Kogut K, Gaspar FW, et al. 2015. Prenatal and childhood polybrominated diphenyl ether (PBDE) exposure and attention and executive function at 9-12 years of age. *Neurotoxicol Teratol* 52(Pt B):151-161. 10.1016/j.ntt.2015.08.001.
- Sahlstrom LM, Sellstrom U, de Wit CA, et al. 2015. Feasibility study of feces for noninvasive biomonitoring of brominated flame retardants in toddlers. *Environ Sci Technol* 49(1):606-615. 10.1021/es504708c.
- Sakai SI, Honda Y, Takatsuki H, et al. 2001. Polybrominated substances in waste electrical and electronic plastics and their behavior in the incineration plants. *Organohalogen Compounds* 52:35-38.
- Sakamoto Y, Inoue K, Takahashi M, et al. 2013. Different pathways of constitutive androstane receptor-mediated liver hypertrophy and hepatocarcinogenesis in mice treated with piperonyl butoxide or decabromodiphenyl ether. *Toxicol Pathol* 41(8):1078-1092.
- Sand S, von Rosen D, Eriksson P, et al. 2004. Dose-response modeling and benchmark calculations from spontaneous behavior data on mice neonatally exposed to 2,2',4,4',5-pentabromodiphenyl ether. *Toxicol Sci* 81(2):491-501.
- Sanders JM, Chen LJ, Lebetkin EH, et al. 2006a. Metabolism and disposition of 2,2',4,4'-tetrabromodiphenyl ether following administration of single or multiple doses to rats and mice. *Xenobiotica* 36(1):103-117.
- Sanders JM, Lebetkin EH, Chen LJ, et al. 2006b. Disposition of 2,2',4,4',5,5'-hexabromodiphenyl ether (BDE153) and its interaction with other polybrominated diphenyl ethers (PBDEs) in rodents. *Xenobiotica* 36(9):824-837.
- Sandholm A, Emanuelsson BM, Klasson-Wehler E. 2003. Bioavailability and half-life of decabromodiphenyl ether (BDE-209) in rat. *Xenobiotica* 33(11):1149-1158.

9. REFERENCES

- Saquib Q, Siddiqui MA, Ahmed J, et al. 2016. Hazards of low dose flame-retardants (BDE-47 and BDE-32): Influence on transcriptome regulation and cell death in human liver cells. *J Hazard Mater* 308:37-49. 10.1016/j.jhazmat.2016.01.025.
- Sarkar D, Chowdhury JP, Singh SK. 2015. Effect of polybrominated diphenyl ether (BDE-209) on testicular steroidogenesis and spermatogenesis through altered thyroid status in adult mice. *Gen Comp Endocrinol*. 10.1016/j.ygcen.2015.11.009.
- Saunders NR, Ek CJ, Habgood MD, et al. 2008. Barriers in the brain: A renaissance? *Trends Neurosci* 31(6):279-286.
- Saunders NR, Liddelow SA, Dziegielewska KM. 2012. Barrier mechanisms in the developing brain. *Front Pharmacol* 3(10.3389/fphar.2012.00046):Article 46.
- Schang G, Robaire B, Hales BF. 2016. Organophosphate flame retardants act as endocrine-disrupting chemicals in MA-10 mouse tumor Leydig Cells. *Toxicol Sci* 150(2):499-509. 10.1093/toxsci/kfw012.
- Schecter A, Colacino J, Sjodin A, et al. 2010. Partitioning of polybrominated diphenyl ethers (PBDEs) in serum and milk from the same mothers. *Chemosphere* 78(10):1279-1284.
- Schecter A, Papke O, Harris TR, et al. 2006. Polybrominated diphenyl ether (PBDE) levels in an expanded market basket survey of U.S. food and estimated PBDE dietary intake by age and sex. *Environ Health Perspect* 114(10):1515-1520.
- Schecter A, Papke O, Tung KC, et al. 2005. Polybrominated diphenyl ether flame retardants in the U.S. population: Current levels, temporal trends, and comparison with dioxins, dibenzofurans, and polychlorinated biphenyls. *J Occup Environ Med* 47(3):199-211.
- Schecter A, Pavuk M, Papke O, et al. 2003. Polybrominated diphenyl ethers (PBDEs) in U.S. mother's milk. *Environ Health Perspect* 111(14):1723-1729.
- Scheuplein R, Charnley G, Dourson M. 2002. Differential sensitivity of children and adults to chemical toxicity. I. Biological basis. *Regul Toxicol Pharmacol* 35(3):429-447.
- Schnare DW, Ben M, Shields MG. 1984. Body burden reductions of PCBs, PBBs and chlorinated pesticides in human subjects. *Ambio* 13:378-380.
- Schreder ED, La Guardia MJ. 2014. Flame retardant transfers from U.S. households (dust and laundry wastewater) to the aquatic environment. *Environ Sci Technol* 48(19):11575-11583. 10.1021/es502227h.
- Schreiber T, Gassmann K, Gotz C, et al. 2010. Polybrominated diphenyl ethers induce developmental neurotoxicity in a human *in vitro* model: Evidence for endocrine disruption. *Environ Health Perspect* 118(4):572-578.
- Schröter-Kermani C, Helm D, Hermann T, et al. 2000. The German environmental specimen bank-application in trend monitoring of polybrominated diphenyl ethers in human blood. *Organohalogen Compounds* 47:49-52.
- Schussler GC. 2000. The thyroxine-binding proteins. *Thyroid* 10:141-149.

9. REFERENCES

- Sellström U, Jansson B. 1995. Analysis of tetrabromobisphenol A in a product and environmental samples. *Chemosphere* 31(4):3085-3092.
- Sellström U, Jansson B, Kierkegaard A, et al. 1993. Polybrominated diphenyl ethers (PBDE) in biological samples from the Swedish environment. *Chemosphere* 26(9):1703-1718.
- Sellström U, Kierkegaard M, Alsberg T, et al. 1999. Brominated flame retardants in sediments from European estuaries, the Baltic Sea and in sewage sludge. *Organohalogen Compounds* 40:383-386.
- Sellström U, Kierkegaard A, De Wit C, et al. 1998a. Polybrominated diphenyl ethers and hexabromocyclododecane in sediment and fish from a Swedish river. *Environ Toxicol Chem* 17(6):1065-1072.
- Sellström U, Soderstrom G, de Wit C, et al. 1998b. Photolytic debromination of decabromodiphenyl ether (DeBDE). *Organohalogen Compounds* 35:447-450.
- Serme-Gbedo YK, Abdelouahab N, Pasquier JC, et al. 2016. Maternal levels of endocrine disruptors, polybrominated diphenyl ethers, in early pregnancy are not associated with lower birth weight in the Canadian birth cohort GESTE. *Environ Health* 15(1):49. 10.1186/s12940-016-0134-z.
- Setchell BP, Waites GMH. 1975. The blood-testis barrier. In: Creep RO, Astwood EB, Geiger SR, eds. *Handbook of physiology: Endocrinology V*. Washington, DC: American Physiological Society, 143-172.
- Shah A, Coburn CG, Watson-Siriboe A, et al. 2011. Altered cardiovascular reactivity and osmoregulation during hyperosmotic stress in adult rats developmentally exposed to polybrominated diphenyl ethers (PBDEs). *Toxicol Appl Pharmacol* 256(2):103-113.
- Shaw SD, Berger ML, Brenner D, et al. 2009. Bioaccumulation of polybrominated diphenyl ethers and hexabromocyclododecane in the northwest Atlantic marine food web. *Sci Total Environ* 407(10):3323-3329.
- She J, Petreas M, Winkler J, et al. 2002. PBDEs in the San Francisco Bay area: Measurement in harbor seal blubber and human breast adipose tissue. *Chemosphere* 46(5):697-707.
- She J, Winkler J, Visita P, et al. 2000. Analysis of PBDEs in seal blubber and human breast adipose tissue samples. *Organohalogen Compounds* 47:53-56.
- Shi G, Yin H, Ye J, et al. 2013. Aerobic biotransformation of decabromodiphenyl ether (PBDE-209) by *Pseudomonas aeruginosa*. *Chemosphere* 93(8):1487-1493.
- Shin JY, Choi YY, Jeon HS, et al. 2010. Low-dose persistent organic pollutants increased telomere length in peripheral leukocytes of healthy Koreans. *Mutagenesis* 25(5):511-516.
- Shy CG, Huang HL, Chang-Chien GP, et al. 2011. Neurodevelopment of infants with prenatal exposure to polybrominated diphenyl ethers. *Bull Environ Contam Toxicol* 87(6):643-648.
- Shy CG, Huang HL, Chao HR, et al. 2012. Cord blood levels of thyroid hormones and IGF-1 weakly correlate with breast milk levels of PBDEs in Taiwan. *Int J Hyg Environ Health* 215(3):345-351.

9. REFERENCES

- Silberhorn EM, Glauert HP, Robertson LW. 1990. Carcinogenicity of polyhalogenated biphenyls: PCBs and PBBs. *Crit Rev Toxicol* 20(6):439-496.
- Simpson S, Gross MS, Olson JR, et al. 2015. Identification of polybrominated diphenyl ether metabolites based on calculated boiling points from COSMO-RS, experimental retention times, and mass spectral fragmentation patterns. *Anal Chem* 87(4):2299-2305.
- Sinjari T, Damerud P, Hallgren S. 1998. Competitive inhibition of 125I-thyroxin (T4) binding to choroid plexus by hydroxylated PCB metabolites. *Organohalogen Compounds* 37:241-244.
- Sjödín A, Carlsson H, Thuresson K, et al. 2001a. Flame retardants in indoor air at an electronics recycling plant and at other work environments. *Environ Sci Technol* 35(3):448-454.
- Sjödín A, Hagmar L, Klasson-Wehler E, et al. 1999a. Flame retardant exposure: Polybrominated diphenyl ethers in blood from Swedish workers. *Environ Health Perspect* 107(8):643-648.
- Sjödín A, Hagmar L, Klasson-Wehler E, et al. 2000. Influence of the consumption of fatty Baltic Sea fish on plasma levels of halogenated environmental contaminants in Latvian and Swedish men. *Environ Health Perspect* 108:1035-1041.
- Sjödín A, Jakobsson E, Kierkegaard A, et al. 1998. Gas chromatographic identification and quantification of polybrominated diphenyl ethers in a commercial product, Bromkal 70-5DE. *J Chromatogr A* 822(1):83-89.
- Sjödín A, Patterson DG Jr., Bergman A. 2001b. Brominated flame retardants in serum from U.S. blood donors. *Environ Sci Technol* 35(19):3830-3833.
- Sjödín A, Thuresson K, Hagmar L, et al. 1999b. Occupational exposure to polybrominated diphenyl ethers at dismantling of electronics. Ambient air and human serum analysis. *Organohalogen Compounds* 43:447-451.
- Sjodin A, Wong LY, Jones RS, et al. 2008. Serum concentrations of polybrominated diphenyl ethers (PBDEs) and polybrominated biphenyl (PBB) in the United States population: 2003-2004. *Environ Sci Technol* 42(4):1377-1384.
- Skarman E, Darnerud PO, Ohrvik H, et al. 2005. Reduced thyroxine levels in mice perinatally exposed to polybrominated diphenyl ethers. *Environ Toxicol Pharmacol* 19(2):273-281.
- Slotkin TA, Card J, Infante A, et al. 2013. BDE99 (2,2',4,4',5-pentabromodiphenyl ether) suppresses differentiation into neurotransmitter phenotypes in PC12 cells. *Neurotoxicol Teratol* 37:13-17.
- Smarr MM, Grantz KL, Zhang C, et al. 2016. Persistent organic pollutants and pregnancy complications. *Sci Total Environ* 1(551-552):285-291. 10.1016/j.scitotenv.2016.02.030.
- Smeds A, Saukko P. 2003. Brominated flame retardants and phenolic endocrine disrupters in Finnish human adipose tissue. *Chemosphere* 53:1123-1130.
- Smolnikar K, Dehnhardt M, Wiegand H. 2001. Perturbation by PBDE99 of calcium homeostasis after *in vitro* treatment. The second international workshop on brominated flame retardants. BFR 2001. Stockholm, Sweden, 273.

9. REFERENCES

- Soderström G, Sellström U, De Wit CA, et al. 2004. Photolytic debromination of decabromodiphenyl ether (BDE 209). *Environ Sci Technol* 38:127-132.
- Song W, Li A, Ford JC, et al. 2005. Polybrominated diphenyl ethers in the sediments of the Great Lakes. 2. Lakes Michigan and Huron. *Environ Sci Technol* 39(10):3474-3479.
- Stafford, CJ. 1983. Halogenated diphenyl ethers identified in avian tissues and eggs by GC/MS. *Chemosphere* 12(11/12):1487-1495.
- Stanley JS, Cramer PH, Thornburg KR, et al. 1991. Mass spectral confirmation of chlorinated and brominated diphenyl ethers in human adipose tissues. *Chemosphere* 23(8-10):1185-1186.
- Stapleton HM. 2006. Instrumental methods and challenges in quantifying polybrominated diphenyl ethers in environmental extracts: A review. *Anal Bioanal Chem* 386:807-817.
- Stapleton HM, Baker JE. 2003. Comparing polybrominated diphenyl ether and polychlorinated biphenyl bioaccumulation in a food web in Grand Bay, Lake Michigan. *Arch Environ Contam Toxicol* 45(2):227-234.
- Stapleton HM, Alaei M, Letcher RJ, et al. 2004. Debromination of the flame retardant decabromodiphenyl ether by juvenile carp (*Cyprinus carpio*) following dietary exposure. *Environ Sci Technol* 38(1):112-119.
- Stapleton HM, Eagle S, Anthopolos R, et al. 2011. Associations between polybrominated diphenyl ether (PBDE) flame retardants, phenolic metabolites, and thyroid hormones during pregnancy. *Environ Health Perspect* 119(10):1454-1459.
- Stapleton HM, Kelly SM, Pei R, et al. 2009. Metabolism of polybrominated diphenyl ethers (PBDEs) by human hepatocytes *in vitro*. *Environ Health Perspect* 117(2):197-202.
- Stapleton HM, Sjodin A, Jones RS, et al. 2008. Serum levels of polybrominated diphenyl ethers (PBDEs) in foam recyclers and carpet installers working in the United States. *Environ Sci Technol* 42(9):3453-3458.
- Staskal DF, Diliberto JJ, Birnbaum LS. 2006a. Impact of repeated exposure on the toxicokinetics of BDE 47 in mice. *Toxicol Sci* 89(2):380-385.
- Staskal DF, Diliberto JJ, DeVito MJ, et al. 2005. Toxicokinetics of BDE 47 in female mice: Effect of dose, route of exposure, and time. *Toxicol Sci* 83(2):215-223.
- Staskal DF, Hakk H, Bauer D, et al. 2006b. Toxicokinetics of polybrominated diphenyl ether congeners 47, 99, 100, and 153 in mice. *Toxicol Sci* 94(1):28-37.
- Stern GA, Ikononou MG. 2000. Temporal trends of polybrominated diphenyl ethers in SE Baffin beluga: Increasing evidence of long range atmospheric transport. *Organohalogen Compounds* 57:81-84.
- Stoker TE, Cooper RL, Lambright CS, et al. 2005. *In vivo* and *in vitro* anti-androgenic effects of DE-71, a commercial polybrominated diphenyl ether (PBDE) mixture. *Toxicol Appl Pharmacol* 207(1):78-88.
- Stoker TE, Ferrell J, Hedge JM, et al. 2003. Assessment of SE-71, a commercial polybrominated diphenyl ether (PBDE) mixture, in the EDSP male pubertal protocol. *Toxicologist* 72(S-1):135-136.

9. REFERENCES

- Stoker TE, Laws SC, Crofton KM, et al. 2004. Assessment of DE-71, a commercial polybrominated diphenyl ether (PBDE) mixture, in the EDSP male and female pubertal protocols. *Toxicol Sci* 78(1):144-155.
- Stoker TE, Cooper RL, Lambright CS, et al. 2005. *In vivo* and *in vitro* anti-androgenic effects of DE-71, a commercial polybrominated diphenyl ether (PBDE) mixture. *Toxicol Appl Pharmacol* 207(1):78-88.
- Strandberg B, Dodder NG, Basu I, et al. 2001. Concentrations and spatial variations of polybrominated diphenyl ethers and other organohalogen compounds in Great Lakes air. *Environ Sci Technol* 35(6):1078-1083.
- Strandman T, Koistinen J, Kiviranta H, et al. 1999. Levels of some polybrominated diphenyl ethers (PBDEs) in fish and human adipose tissue in Finland. *Organohalogen Compounds* 40:355-358.
- Strandman T, Koistinen J, Variainen T. 2000. Polybrominated diphenyl ethers (PBDEs) in placenta and human milk. *Chemosphere* 47:61-64.
- Streets SS, Henderson SA, Stoner AD, et al. 2006. Partitioning and bioaccumulation of PBDEs and PCBs in Lake Michigan. *Environ Sci Technol* 40(23):7263-7269.
- Stross JK, Smokler IA, Isbister J, et al. 1981. The human health effects of exposure to polybrominated biphenyls. *Toxicol Appl Pharmacol* 58:145-150.
- Stubbings WA, Harrad S. 2014. Extent and mechanisms of brominated flame retardant emissions from waste soft furnishings and fabrics: A critical review. *Environ Int* 71:164-175.
- Su G, Xia J, Liu H, et al. 2012. Dioxin-like potency of HO- and MeO- analogues of PBDEs' the potential risk through consumption of fish from eastern China. *Environ Sci Technol* 46(19):10781-10788.
- Sutton R, Sedlak MD, Yee D, et al. 2015. Declines in polybrominated diphenyl ether contamination of San Francisco Bay following production phase-outs and bans. *Environ Sci Technol* 49(2):777-784. 10.1021/es503727b.
- Swann RL, Laskowski DA, McCall PJ, et al. 1983. A rapid method for the estimation of the environmental parameters octanol/water partition coefficient, soil sorption constant, water to air, and water solubility. *Res Rev* 85:17-28.
- Szabo DT, Richardson VM, Ross DG, et al. 2009. Effects of perinatal PBDE exposure on hepatic phase I, phase II, phase III, and deiodinase 1 gene expression involved in thyroid hormone metabolism in male rat pups. *Toxicol Sci* 107(1):27-39. 10.1093/toxsci/kfn230.
- Ta TA, Koenig CM, Golub MS, et al. 2011. Bioaccumulation and behavioral effects of 2,2',4,4'-tetrabromodiphenyl ether (BDE-47) in perinatally exposed mice. *Neurotoxicol Teratol* 33(3):393-404.
- Takenaka S, Takahashi K. 1991. Enhancement of fecal excretion of polychlorinated biphenyls by the addition of rice bran fiber to the diet in rats. *Chemosphere* 22(3-4):375-381.

9. REFERENCES

- Talsness CE, Kuriyama SN, Sterner-Kock A, et al. 2008. *In utero* and lactational exposures to low doses of polybrominated diphenyl ether-47 alter the reproductive system and thyroid gland of female rat offspring. *Environ Health Perspect* 116(3):308-314.
- Talsness CE, Shakibaei M, Kuriyama S, et al. 2003. Ultrastructural changes in the ovaries of adult offspring following a single maternal exposure to low dose 2,2',4,4',5-pentabromodiphenyl ether. *Organohalogen Compounds* 61:88-91.
- Talsness CE, Shakibaei M, Kuriyama SN, et al. 2005. Ultrastructural changes observed in rat ovaries following in utero and lactational exposure to low doses of a polybrominated flame retardant. *Toxicol Lett* 157(3):189-202.
- Tan J, Loganath A, Chong YS, et al. 2009. Exposure to persistent organic pollutants in utero and related maternal characteristics on birth outcomes: A multivariate data analysis approach. *Chemosphere* 74(3):428-433.
- Tang Z, Huang Q, Cheng J, et al. 2014. Polybrominated diphenyl ethers in soils, sediments, and human hair in a plastic waste recycling area: A neglected heavily polluted area. *Environ Sci Technol* 48(3):1508-1516.
- Thomas K, Colborn T. 1992. Organochlorine endocrine disruptors in human tissue. In: Colborn T, Clement C, eds. *Chemically induced alterations in sexual and functional development: The wildlife/human connection*. Princeton, NJ: Princeton Scientific Publishing, 365-394.
- Thomsen C, Haug LS, Stigum H, et al. 2010. Changes in concentrations of perfluorinated compounds, polybrominated diphenyl ethers, and polychlorinated biphenyls in Norwegian breast-milk during twelve months of lactation. *Environ Sci Technol* 44(24):9550-9556.
- Thomsen C, Lundanes E, Becher G. 2001. Brominated flame retardants in plasma samples from three different occupational groups in Norway. *J Environ Monit* 3(4):366-370.
- Thureson K, Hoglund P, Hagmar L, et al. 2006. Apparent half-lives of hepta- to decabrominated diphenyl ethers in human serum as determined in occupationally exposed workers. *Environ Health Perspect* 114(2):176-181.
- Tittlemier SA, Halldorson T, Stern GA, et al. 2002. Vapor pressures, aqueous solubilities, and Henry's Law constants of some brominated flame retardants. *Environ Toxicol Chem* 21(9):1804-1810.
- Toft G, Lenters V, Vermeulen R, et al. 2014. Exposure to polybrominated diphenyl ethers and male reproductive function in Greenland, Poland and Ukraine. *Reprod Toxicol* 43:1-7. 10.1016/j.reprotox.2013.10.002.
- Tokarz JA, Ahn MY, Leng J, et al. 2008. Reductive debromination of polybrominated diphenyl ethers in anaerobic sediment and a biomimetic system. *Environ Sci Technol* 42(4):1157-1164.
- Toms LM, Sjodin A, Harden F, et al. 2009. Serum polybrominated diphenyl ether (PBDE) levels are higher in children (2-5 years of age) than in infants and adults. *Environ Health Perspect* 117(9):1461-1465.
- Trabert B, Chen Z, Kannan K, et al. 2015. Persistent organic pollutants (POPs) and fibroids: Results from the ENDO study. *J Expo Sci Environ Epidemiol* 25(3):278-285. 10.1038/jes.2014.31.

9. REFERENCES

- TRI14 2016. TRI explorer: Providing access to EPA's toxics release inventory data. U.S. Environmental Protection Agency, Office of Environmental Information. Toxics Release Inventory. <http://www.epa.gov/triexplorer/>. September 19, 2016.
- Trudel D, Scheringer M, von Goetz N, et al. 2011. Total consumer exposure to polybrominated diphenyl ethers in North America and Europe. *Environ Sci Technol* 45(6):2391-2397.
- Tseng LH, Hsu PC, Lee CW, et al. 2013. Developmental exposure to decabrominated diphenyl ether (BDE-209): Effects on sperm oxidative stress and chromatin DNA damage in mouse offspring. *Environ Toxicol* 28(7):380-389.
- Tseng LH, Li MH, Hsu PC, et al. 2006. Effects of in utero exposure to decabrominated diphenyl ether (PBDE 209) on developmental toxicity and liver enzyme activities in male mice. *Organohalogen Compounds* 68:1561-1564.
- Tseng LH, Li MH, Tsai SS, et al. 2008. Developmental exposure to decabromodiphenyl ether (PBDE 209): Effects on thyroid hormone and hepatic enzyme activity in male mouse offspring. *Chemosphere* 70(4):640-647.
- Tullo A. 2003. Great Lakes to phase out two flame retardants. *Chemical & Engineering News* 81(45):13.
- Turyk M, Fantuzzi G, Persky V, et al. 2015. Persistent organic pollutants and biomarkers of diabetes risk in a cohort of Great Lakes sport caught fish consumers. *Environ Res* 140:335-344. 10.1016/j.envres.2015.03.037.
- Turyk ME, Persky VW, Imm P, et al. 2008. Hormone disruption by PBDEs in adult male sport fish consumers. *Environ Health Perspect* 116(12):1635-1641.
- Ueno D, Kajiwara N, Tanaka H, et al. 2003. Global pollution monitoring of polybrominated diphenyl ethers (PBDEs) using skipjack tuna as a bioindicator. *Organohalogen Compounds* 61:37-40.
- USGS. 2012. Reconnaissance of contaminants in selected wastewater-treatment-plant effluent and stormwater runoff entering the Columbia River, Columbia River Basin, Washington and Oregon, 2008-10. Scientific Investigations Report 2012-5068. U.S. Geological Survey, U.S. Department of the Interior. <http://pubs.usgs.gov/sir/2012/5068/pdf/sir20125068.pdf>. September 22, 2016.
- Vagi SJ, Azziz-Baumgartner E, Sjodin A, et al. 2014. Exploring the potential association between brominated diphenyl ethers, polychlorinated biphenyls, organochlorine pesticides, perfluorinated compounds, phthalates, and bisphenol A in polycystic ovary syndrome: A case-control study. *BMC endocrine disorders* 14:86. 10.1186/1472-6823-14-86.
- van Bavel B, Dam M, Tysklind M, et al. 2001. Levels of polybrominated diphenyl ethers in marine mammals. *Organohalogen Compounds* 52:99-103.
- van Bavel B, Sundelin E, Lillback J, et al. 1999. Supercritical fluid extraction of polybrominated diphenyl ethers PBDEs from long-finned pilot whale (*Globicephala melas*) from the Atlantic. *Organohalogen Compounds* 40:359-362.

9. REFERENCES

- van den Dungen MW, Rijk JC, Kampman E, et al. 2015. Steroid hormone related effects of marine persistent organic pollutants in human H295R adrenocortical carcinoma cells. *Toxicol in Vitro* 29(4):769-778. 10.1016/j.tiv.2015.03.002.
- van den Hove MF, Beckers C, Devlieger H, et al. 1999. Hormone synthesis and storage in the thyroid of human preterm and term newborns: Effect of thyroxine treatment. *Biochimie* 81:563-570.
- Van der Ven LT, van de Kuil T, Leonards PE, et al. 2008a. A 28-day oral dose toxicity study in Wistar rats enhanced to detect endocrine effects of decabromodiphenyl ether (decaBDE). *Toxicol Lett* 179(1):6-14.
- van der Ven LT, van de Kuil T, Verhoef A, et al. 2008b. A 28-day oral dose toxicity study enhanced to detect endocrine effects of a purified technical pentabromodiphenyl ether (pentaBDE) mixture in Wistar rats. *Toxicology* 245(1-2):109-122.
- Van Vliet G. 1999. Merck AG thyroid symposium. Neonatal hypothyroidism: Treatment and outcome. *Thyroid* 9:79-84.
- Venkatesan AK, Halden RU. 2014. Brominated flame retardants in U.S. biosolids from the EPA national sewage sludge survey and chemical persistence in outdoor soil mesocosms. *Water Res* 55:133-142. 10.1016/j.watres.2014.02.021.
- Verma P, Gupta RK, Gandhi BS, et al. 2015. CDRI-08 Attenuates REST/NRSF-mediated expression of NMDAR1 Gene in PBDE-209-exposed mice brain. *eCAM* 2015:403840. 10.1155/2015/403840.
- Viberg H. 2009a. Exposure to polybrominated diphenyl ethers 203 and 206 during the neonatal brain growth spurt affects proteins important for normal neurodevelopment in mice. *Toxicol Sci* 109(2):306-311.
- Viberg H. 2009b. Neonatal ontogeny and neurotoxic effect of decabrominated diphenyl ether (PBDE 209) on levels of synaptophysin and tau. *Int J Dev Neurosci* 27(5):423-429.
- Viberg H, Eriksson P. 2011. Differences in neonatal neurotoxicity of brominated flame retardants, PBDE 99 and TBBPA, in mice. *Toxicology* 289(1):59-65.
- Viberg H, Fredriksson A, Eriksson P. 2002. Neonatal exposure to the brominated flame retardant 2,2',4,4',4-pentabromodiphenyl ether causes altered susceptibility of the cholinergic transmitter system in the adult mouse. *Toxicol Sci* 67:104-107.
- Viberg H, Fredrickson A, Eriksson P. 2003a. Neonatal PBDE 99 exposure causes dose-response related behavioural derangements that are not sex or strain specific in mice. *Toxicol Sci* 72(S-1):126.
- Viberg H, Fredriksson A, Eriksson P. 2003b. Neurotoxicity of different polybrominated diphenyl ethers, including PBDE 209. *Organohalogen Compounds* 65:9-11.
- Viberg H, Fredriksson A, Eriksson P. 2004a. Investigations of strain and/or gender differences in developmental neurotoxic effects of polybrominated diphenyl ethers in mice. *Toxicol Sci* 81(2):344-353.
- Viberg H, Fredriksson A, Eriksson P. 2004b. Neonatal exposure to the brominated flame-retardant, 2,2',4,4',5-pentabromodiphenyl ether, decreases cholinergic nicotinic receptors in hippocampus and affects spontaneous behaviour in the adult mouse. *Environ Toxicol Pharmacol* 17(2):61-65.

9. REFERENCES

- Viberg H, Fredriksson A, Eriksson P. 2005. Deranged spontaneous behaviour and decrease in cholinergic muscarinic receptors in hippocampus in the adult rat, after neonatal exposure to the brominated flame-retardant, 2,2',4,4',5-pentabromodiphenyl ether (PBDE 99). *Environ Toxicol Pharmacol* 20(2):283-288.
- Viberg H, Fredriksson A, Eriksson P. 2007. Changes in spontaneous behaviour and altered response to nicotine in the adult rat, after neonatal exposure to the brominated flame retardant, decabrominated diphenyl ether (PBDE 209). *Neurotoxicology* 28(1):136-142.
- Viberg H, Johansson N, Fredriksson A, et al. 2006. Neonatal exposure to higher brominated diphenyl ethers, hepta-, octa-, or nonabromodiphenyl ether, impairs spontaneous behavior and learning and memory functions of adult mice. *Toxicol Sci* 92(1):211-218.
- Viberg H, Mundy W, Eriksson P. 2008. Neonatal exposure to decabrominated diphenyl ether (PBDE 209) results in changes in BDNF, CaMKII and GAP-43, biochemical substrates of neuronal survival, growth, and synaptogenesis. *Neurotoxicology* 29(1):152-159.
- Vieira I, Sonnier M, Cresteil T. 1996. Developmental expression of CYP2E1 in the human liver: Hypermethylation control of gene expression during the neonatal period. *Eur J Biochem* 238(2):476-483.
- Villeneuve DL, Kannan K, Priest BT, et al. 2002. *In vitro* assessment of potential mechanism-specific effects of polybrominated diphenyl ethers. *Environ Toxicol Chem* 21(11):2431-2433.
- Vizcaino E, Grimalt JO, Lopez-Espinosa MJ, et al. 2011. Polybromodiphenyl ethers in mothers and their newborns from a non-occupationally exposed population (Valencia, Spain). *Environ Int* 37(1):152-157.
- von Meyerinck L, Hufnagel B, Schmoldt A, et al. 1990. Inductions of rat liver microsomal cytochrome P-450 by the pentabromo diphenyl ether bromkal 70 and half-lives of its components in the adipose tissue. *Toxicology* 61(2):259-274.
- Vulsma T, Gons MH, DeVijder JJM. 1989. Maternal-fetal transfer of thyroxine in congenital hypothyroidism due to a total organification defect or thyroid agenesis. *N Eng J Med* 321:13-16.
- Vuong AM, Yolton K, Webster GM, et al. 2016a. Prenatal polybrominated diphenyl ether and perfluoroalkyl substance exposures and executive function in school-age children. *Environ Res* 147:556-564. 10.1016/j.envres.2016.01.008.
- Vuong AM, Braun JM, Sjodin A, et al. 2016b. Prenatal polybrominated diphenyl ether exposure and body mass index in children up to 8 years of age. *Environ Health Perspect* 124(12):1891-1897. 10.1289/EHP139.
- Wainman BC, Kesner JS, Martin ID, et al. 2016. Menstrual cycle perturbation by organohalogen and elements in the Cree of James Bay, Canada. *Chemosphere* 149:190-201. 10.1016/j.chemosphere.2015.12.056.
- Wan Y, Choi K, Kim S, et al. 2010. Hydroxylated polybrominated diphenyl ethers and bisphenol A in pregnant women and their matching fetuses: Placental transfer and potential risks. *Environ Sci Technol* 44(13):5233-5239.

9. REFERENCES

- Wan Y, Wiseman S, Chang H, et al. 2009. Origin of hydroxylated brominated diphenyl ethers: Natural compounds or man-made flame retardants? *Environ Sci Technol* 43(19):7536-7542.
- Wang F, Liu W, Jin Y, et al. 2011a. Interaction of PFOS and BDE-47 co-exposure on thyroid hormone levels and TH-related gene and protein expression in developing rat brains. *Toxicol Sci* 121(2):279-291.
- Wang F, Wang J, Dai J, et al. 2010. Comparative tissue distribution, biotransformation and associated biological effects by decabromodiphenyl ethane and decabrominated diphenyl ether in male rats after a 90-day oral exposure study. *Environ Sci Technol* 44(14):5655-5660.
- Wang F, Wang J, Hu G, et al. 2011b. Tissue distribution and associated toxicological effects of decabrominated diphenyl ether in subchronically exposed male rats. *ISRN Toxicology* 2011:989251.
- Wang KL, Hsia SM, Mao IF, et al. 2011c. Effects of polybrominated diphenyl ethers on steroidogenesis in rat Leydig cells. *Hum Reprod* 26(8):2209-2217.
- Wang Y, Shi J, Li L, et al. 2013. Adverse effects of 2,2',4,4'-tetrabromodiphenyl ether on semen quality and spermatogenesis in male mice. *Bull Environ Contam Toxicol* 90(1):51-54.
- Wang Z, Na G, Ma X, et al. 2015. Characterizing the distribution of selected PBDEs in soil, moss and reindeer dung at Ny-Alesund of the Arctic. *Chemosphere* 137:9-13. 10.1016/j.chemosphere.2015.04.030.
- Warembourg C, Debost-Legrand A, Bonvallot N, et al. 2016. Exposure of pregnant women to persistent organic pollutants and cord sex hormone levels. *Hum Reprod* 31(1):190-198. 10.1093/humrep/dev260.
- Watanabe I. 1988. Behaviour of organobrominated compounds at the sediment phase in the environment. *Koshu Eisei Hen* 26:129-133.
- Watanabe I, Kashimoto T, Tatsukawa R. 1986. Confirmation of the presence of the flame retardant decabromodiphenyl ether in river sediment from Osaka, Japan. *Bull Environ Contam Toxicol* 36(6):839-842.
- Watanabe I, Kashimoto T, Tatsukawa R. 1987. Polybrominated biphenyl ethers in marine fish, shellfish and river and marine sediments in Japan. *Chemosphere* 16(10-12):2389-2396.
- Watanabe I, Kawano M, Tatsukawa R. 1995. Polybrominated and mixed polybromo/chlorinated dibenzo-p-dioxins and -dibenzofurans in the Japanese environment. *Organohalogen Compounds* 24:337-340.
- Watanabe W, Shimizu T, Hino A, et al. 2008. Effects of decabrominated diphenyl ether (DBDE) on developmental immunotoxicity in offspring mice. *Environ Toxicol Pharmacol* 26(3):315-319.
- Watanabe W, Shimizu T, Sawamura R, et al. 2010a. Effects of tetrabromobisphenol A, a brominated flame retardant, on the immune response to respiratory syncytial virus infection in mice. *Int Immunopharmacol* 10(4):393-397.
- Watanabe W, Shimizu T, Sawamura R, et al. 2010b. Functional disorder of primary immunity responding to respiratory syncytial virus infection in offspring mice exposed to a flame retardant, decabrominated diphenyl ether, perinatally. *J Med Virol* 82(6):1075-1082.

9. REFERENCES

- Watkins DJ, McClean MD, Fraser AJ, et al. 2011. Exposure to PBDEs in the office environment: Evaluating the relationships between dust, handwipes, and serum. *Environ Health Perspect* 119(9):1247-1252.
- Weiss J, Wallin E, Axmon A, et al. 2006. Hydroxy-PCBs, PBDEs, and HBCDDs in serum from an elderly population of Swedish fishermen's wives and associations with bone density. *Environ Sci Technol* 40(20):6282-6289.
- West JR, Smith HW, Chasis H. 1948. Glomerular filtration rate, effective renal blood flow, and maximal tubular excretory capacity in infancy. *J Pediatr* 32:10-18.
- Wester RC, Maibach HI, Bucks DA, et al. 1990. Percutaneous absorption and skin decontamination of PCBs: *In vitro* studies with human skin and *in vivo* studies in the Rhesus monkey. *J Toxicol Environ Health* 31:235-246.
- Whitehead TP, Crispo Smith S, Park JS, et al. 2015a. Concentrations of persistent organic pollutants in California women's serum and residential dust. *Environ Res* 136:57-66. 10.1016/j.envres.2014.10.009.
- Whitehead TP, Crispo Smith S, Park JS, et al. 2015b. Concentrations of persistent organic pollutants in California children's whole blood and residential dust. *Environ Sci Technol* 49(15):9331-9340. 10.1021/acs.est.5b02078.
- WHO. 1994a. Brominated diphenyl ethers. International programme on chemical safety. Environmental Health Criteria 162. World Health Organization. <http://www.inchem.org/documents/ehc/ehc/ehc162.htm>. December 17, 2014.
- WHO. 1994b. Brominated biphenyls. International programme on chemical safety. Environmental Health Criteria 152. World Health Organization. <http://www.inchem.org/documents/ehc/ehc/ehc152.htm>. December 17, 2014.
- WHO. 1998. Polybrominated dibenzo-p-dioxins and dibenzofurans. International Programme on Chemical Safety. Environmental Health Criteria. World Health Organization. <http://www.inchem.org/documents/ehc/ehc/ehc205.htm>. December 17, 2014
- WHO. 2006. Evaluation of certain food contaminants. Sixty-fourth report of the Joint/FAO/Who Expert Committee on Food Additives. Geneva, Switzerland: World Health Organization. http://whqlibdoc.who.int/trs/WHO_TRS_930_eng.pdf. December 17, 2014
- WHO. 2010. Guidelines for indoor air quality: Selected pollutants. Geneva, Switzerland: World Health Organization. http://www.euro.who.int/__data/assets/pdf_file/0009/128169/e94535.pdf. September 9, 2014.
- WHO. 2011. Guidelines for drinking-water quality. 4th ed. Geneva, Switzerland: World Health Organization. http://whqlibdoc.who.int/publications/2011/9789241548151_eng.pdf?ua=1. September 9, 2014.
- Widdowson EM, Dickerson JWT. 1964. Chemical composition of the body. In: Comar CL, Bronner F, eds. Mineral metabolism: An advanced treatise. Volume II: The elements Part A. New York, NY: Academic Press, 1-247.

9. REFERENCES

- Wiegand H, Desai D, Dehnhardt M, et al. 2001. Polyhalogenated hydrocarbon induced perturbation of intracellular calcium homeostasis: From astrocytes to human macrophages. *Organohalogen Compounds* 53:182-184.
- WIL Research Laboratories. 1984. 90-Day dietary study in rats with pentabromodiphenyl oxide (DE-71) (Final Report). Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. OTS0524336.
- WIL Research Laboratories. 1986. A range-finding teratology study in rats with DE-79. Submitted to U.S. Environmental Protection Agency under TSCA Section 8D. OTS0522298.
- Windham GC, Pinney SM, Voss RW, et al. 2015a. Brominated flame retardants and other persistent organohalogenated compounds in relation to timing of puberty in a longitudinal study of girls. *Environ Health Perspect* 123(10):1046-1052. 10.1289/ehp.1408778.
- Windham GC, Pinney SM, Voss RW, et al. 2015b. Supplemental material. Brominated flame retardants and other persistent organohalogenated compounds in relation to timing of puberty in a longitudinal study of girls. *Environ Health Perspect*. 10.1289/ehp.1408778. <http://ehp.niehs.nih.gov/wp-content/uploads/123/10/ehp.1408778.s001.acco.pdf>. September 19, 2016.
- Witt KL, Livanos E, Kissling GE, et al. 2008. Comparison of flow cytometry- and microscopy-based methods for measuring micronucleated reticulocyte frequencies in rodents treated with nongenotoxic and genotoxic chemicals. *Mutat Res* 649(1-2):101-113.
- Wolfe NL, Jeffers PM. 2000. Hydrolysis. In: Boethling SB, Mackay D, eds. *Handbook of property estimation methods for chemicals*. Boca Raton, FL: CRC Press LLC, 311-334.
- Wolff MS, Anderson HA, Camper F, et al. 1979a. Analysis of adipose tissue and serum from PBB (polybrominated biphenyl)-exposed workers. *J Environ Pathol Toxicol* 2:1397-1411.
- Wolff MS, Anderson HA, Rosenman KD, et al. 1979b. Equilibrium of polybrominated biphenyl (PBB) residues in serum and fat of Michigan residents. *Bull Environ Contam Toxicol* 21:775-781.
- Wolff MS, Anderson HA, Selikoff IJ. 1982. Human tissue burdens of halogenated aromatic chemicals in Michigan. *JAMA* 247(15):2112-2116.
- Wong F, Cousins IT, Macleod M. 2013. Bounding uncertainties in intrinsic human elimination half-lives and intake of polybrominated diphenyl ethers in the North American population. *Environ Int* 59:168-174.
- Woods R, Vallero RO, Golub MS, et al. 2012. Long-lived epigenetic interactions between perinatal PBDE exposure and MeCP2³⁰⁸ mutation. *Hum Mol Genet* 21(11):2399-2411.
- Wu CC, Bao LJ, Tao S, et al. 2016. Dermal uptake from airborne organics as an important route of human exposure to e-waste combustion fumes. *Environ Sci Technol* 50(13):6599-6605. 10.1021/acs.est.5b05952.
- Wu K, Xu X, Liu J, et al. 2010. Polybrominated diphenyl ethers in umbilical cord blood and relevant factors in neonates from Guiyu, China. *Environ Sci Technol* 44(2):813-819.

9. REFERENCES

- Xing T, Chen L, Tao Y, et al. 2009. Effects of decabrominated diphenyl ether (PBDE 209) exposure at different developmental periods on synaptic plasticity in the dentate gyrus of adult rats *in vivo*. *Toxicol Sci* 110(2):401-410.
- Xu L, Huo X, Zhang Y, et al. 2015b. Polybrominated diphenyl ethers in human placenta associated with neonatal physiological development at a typical e-waste recycling area in China. *Environ Pollut* 196:414-422. 10.1016/j.envpol.2014.11.002.
- Xu P, Lou X, Ding G, et al. 2014a. Association of PCB, PBDE and PCDD/F body burdens with hormone levels for children in an e-waste dismantling area of Zhejiang Province, China. *Sci Total Environ* 499:55-61. 10.1016/j.scitotenv.2014.08.057.
- Xu P, Lou X, Ding G, et al. 2015a. Effects of PCBs and PBDEs on thyroid hormone, lymphocyte proliferation, hematology and kidney injury markers in residents of an e-waste dismantling area in Zhejiang, China. *Sci Total Environ* 536:215-222. 10.1016/j.scitotenv.2015.07.025.
- Xu X, Liu J, Zeng X, et al. 2014b. Elevated serum polybrominated diphenyl ethers and alteration of thyroid hormones in children from Guiyu, China. *PLoS ONE* 9(11):e113699. 10.1371/journal.pone.0113699.
- Yamamoto H, Okumura T, Nishikawa Y, et al. 1997. Determination of decabromodiphenyl ether in water and sediment samples by gas chromatography with electron capture detection. *J AOAC Int* 80(1):102-106.
- Yan T, Xiang L, Xuejun J, et al. 2012. Spatial learning and memory deficit of low level polybrominated diphenyl ethers-47 in male adult rat is modulated by intracellular glutamate receptors. *J Toxicol Sci* 37(2):223-233.
- Yang C, Harrad S, M Abdallah, MAE et al. 2014. Polybrominated diphenyl ethers (PBDEs) in English freshwater lakes, 2008–2012. *Chemosphere* 110:41-47.
- Yang F, Alonso Aguirre A, Jin S, et al. 2008. Detection of polybrominated diphenyl ethers in tilapia (*Oreochromis mossambicus*) from O'ahu, Hawai'i. *J Environ Monit* 10(4):432-434.
- Yeh A, Kruse SE, Marcinek DJ, et al. 2015. Effect of omega-3 fatty acid oxidation products on the cellular and mitochondrial toxicity of BDE 47. *Toxicol in Vitro* 29(4):672-680. 10.1016/j.tiv.2015.01.015.
- Yu YX, Li JL, Zhang XY, et al. 2010b. Assessment of the bioaccessibility of polybrominated diphenyl ethers in foods and the correlations of the bioaccessibility with nutrient contents. *J Agric Food Chem* 58(1):301-308.
- Yu Z, Zheng K, Ren G, et al. 2010a. Identification of hydroxylated octa- and nona-bromodiphenyl ethers in human serum from electronic waste dismantling workers. *Environ Sci Technol* 44(10):3979-3985.
- Yuan J, Chen L, Chen D, et al. 2008. Elevated serum polybrominated diphenyl ethers and thyroid-stimulating hormone associated with lymphocytic micronuclei in Chinese workers from an E-waste dismantling site. *Environ Sci Technol* 42(6):2195-2200.
- Zegers BN, Lewis WE, Boon JP. 2000. Levels of some polybrominated diphenyl ether (PBDE) flame retardants in dated sediment cores. *Organohalogen Compounds* 47:229-232.

9. REFERENCES

- Zennegg M, Kohker M, Gerecke AC, et al. 2003. Polybrominated diphenyl ethers in whitefish from Swiss lakes and farmed rainbow trout. *Chemosphere* 51(7):545-553.
- Zennegg M, Munoz M, Schmid P, et al. 2013. Temporal trends of persistent organic pollutants in digested sewage sludge (1993-2012). *Environ Int* 60:202-208.
- Zhang S, Bursian SJ, Martin PA, et al. 2009. Reproductive and developmental toxicity of a pentabrominated diphenyl ether mixture, DE-71, to ranch mink (*Mustela vison*) and hazard assessment for wild mink in the Great Lakes region. *Toxicol Sci* 110(1):107-116.
- Zhang S, Kuang G, Zhao G, et al. 2013c. Involvement of the mitochondrial p53 pathway in PBDE-47-induced SH-SY5Y cells apoptosis and its underlying activation mechanism. *Food Chem Toxicol* 62:699-706.
- Zhang W, Cai Y, Sheng G, et al. 2011. Tissue distribution of decabrominated diphenyl ether (BDE-209) and its metabolites in sucking rat pups after prenatal and/or postnatal exposure. *Toxicology* 283(1):49-54.
- Zhang X, Feng M, Liu F, et al. 2014. Subacute oral toxicity of BDE-15, CDE-15, and HODE-15 in ICR male mice: Assessing effects on hepatic oxidative stress and metals status and ascertaining the protective role of vitamin E. *Environ Sci Pollut Res Int* 21(3):1924-1935.
- Zhang Z, Sun ZZ, Xiao X, et al. 2013a. Mechanism of BDE209-induced impaired glucose homeostasis based on gene microarray analysis of adult rat liver. *Arch Toxicol* 87(8):1557-1567.
- Zhang Z, Zhang X, Sun Z, et al. 2013b. Cytochrome P450 3A1 mediates 2,2',4,4'-tetrabromodiphenyl ether-induced reduction of spermatogenesis in adult rats. *PLoS ONE* 8(6):e66301. <http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0066301>. August 14, 2014.
- Zhang ZF, Shan Q, Zhuang J, et al. 2015a. Troxerutin inhibits 2,2',4,4'-tetrabromodiphenyl ether (BDE-47)-induced hepatocyte apoptosis by restoring proteasome function. *Toxicol Lett* 233(3):246-257. 10.1016/j.toxlet.2015.01.017.
- Zhang ZF, Zhang YQ, Fan SH, et al. 2015b. Troxerutin protects against 2,2',4,4'-tetrabromodiphenyl ether (BDE-47)-induced liver inflammation by attenuating oxidative stress-mediated NAD(+)-depletion. *J Hazard Mater* 283:98-109. 10.1016/j.jhazmat.2014.09.012.
- Zhao W, Cheng J, Gu J, et al. 2014. Assessment of neurotoxic effects and brain region distribution in rat offspring prenatally co-exposed to low doses of BDE-99 and methylmercury. *Chemosphere* 112:170-176. 10.1016/j.chemosphere.2014.04.011.
- Zhao Y, Ao H, Chen L, et al. 2011. Effect of brominated flame retardant BDE-47 on androgen production of adult rat Leydig cells. *Toxicol Lett* 205(2):209-214.
- Zheng J, Luo XJ, Yuan JG, et al. 2011. Levels and sources of brominated flame retardants in human hair from urban, e-waste, and rural areas in south China. *Environ Pollut* 159(12):3706-3713.
- Zhou T, Ross DG, De Vito MJ, et al. 2001. Effects of short-term *in vivo* exposure to polybrominated diphenyl ethers on thyroid hormones and hepatic enzyme activities in weaning rats. *Toxicol Sci* 61:76-82.

9. REFERENCES

- Zhou T, Taylor MM, DeVito MJ, et al. 2002. Developmental exposure to brominated diphenyl ethers results in thyroid hormone disruption. *Toxicol Sci* 66:105-116.
- Zhu LY, Hites RA. 2005. Brominated flame retardants in sediment cores from Lakes Michigan and Erie. *Environ Sci Technol* 39(10):3488-3494.
- Ziegler EE, Edwards BB, Jensen RL, et al. 1978. Absorption and retention of lead by infants. *Pediatr Res* 12(1):29-34.
- Zier B, Lenoir D, Lahaniatis ES, et al. 1991. Surface catalyzed halogenation-dehalogenation reactions of aromatic bromine compounds adsorbed on fly ash. *Chemosphere* 22(12):1121-1129.
- Zoeller RT, Crofton RM. 2000. Thyroid hormone action in fetal brain development and potential for distribution by environmental chemicals. *Neurotoxicology* 21(6):935-946.
- Zweidinger R, Cooper SD, Erickson MD, et al. 1979. Sampling and analysis for semivolatile brominated organics in ambient air. *Am Chem Soc Abstr Pap* 94:217-231.