9. REFERENCES

+*Abdel-Rahman MS, Saladin JJ, Bohman CE, et al. 1978. The effect of 2-hexanone and 2-hexanone metabolites on pupillomotor activity and growth. Am Ind Hyg Assoc J 39:94-99.

*Abdo KM, Graham DG, Timmons PR, et al. 1982. Neurotoxicity of continuous (90 days) inhalation of technical grade methyl butyl ketone in hens. J Toxicol Environ Health 9:199-215.

*Abou-Donia MB, Hu ZH, Lapadula DM, et al. 1991. Mechanisms of joint neurotoxicity of n-hexane, methyl isobutyl ketone and O-ethyl *O*-4-nitrophenyl phenylphosphonothioate in hens. J Pharmacol Exp Ther 257(1):282-289.

*Abou-Donia MB, Lapadula DM, Campbell G, et al. 1985a. The joint neurotoxic action of inhaled methyl butyl ketone vapor and dermally applied *0*-ethyl *0*-4nitrophenyl phenylphosphonothioate in hens: Potentiating effect. Toxicol Appl Pharmacol 79:69-82.

*Abou-Donia MB, Makkawy H-AM, Campbell GM. 1985b. Pattern of neurotoxicity of *n*-hexane, methyl *n*-butyl ketone, 2,5-hexanediol, and 2,5-hexanedione alone and in combination with *O*-ethyl *O*-4-nitrophenyl phenylphosphonothioate in hens. J Toxicol Environ Health 16:85-100.

*Abou-Donia MB, Lapadula DM, Campbell G, et al. 1985c. The synergism of n-hexane-induced neurotoxicity by methyl isobutyl ketone following subchronic (90 days) inhalation in hens: Induction of hepatic microsomal cytochrome P-450. Toxicol Appl Pharmacol 81:1-16.

*Abou-Donia MB, Makkawy H-AM, Graham DB. 1982. The relative neurotoxicities of *n*-hexane, methyl *n*-butyl ketone, 2,5-hexanediol, and 2,5-hexanedione following oral or intraperitoneal administration in hens. Toxicol Appl Pharmacol 62:369389.

*ACGIH. 1986. Documentation of the threshold limit values and biological exposure indices. 5th ed. Cincinnati, OH: American Conference of Governmental Industrial Hygienists.

*ACGIH. 2001. Methyl n-butyl ketone. In: Documentation of the threshold limit values and biological exposure indices, 7th edition. Cincinnati, OH: American Conference of Governmental Industrial Hygienists.

*ACGIH. 2003. Methyl n-butyl ketone. In: Documentation of the biological exposure indices, 7th edition. Cincinnati, OH: American Conference of Governmental Industrial Hygienists.

*ACGIH. 2015. Methyl n-butyl ketone. In: TLVs and BEIs based on the documentation of the threshold limit values for chemical substances and physical agents and biological exposure indices. Cincinnati, OH: American Conference of Governmental Industrial Hygienists.

*Adlercreutz H. 1995. Phytoestrogens: Epidemiology and a possible role in cancer protection. Environ Health Perspect 103(Suppl 7):103-112.

^{*} Cited in text

⁺ Cited in supplemental document

*AEGLs. 2016. Acute Exposure Guideline Levels (AEGLs) Values. U.S. Environmental Protection Agency. https://www.epa.gov/aegl/access-acute-exposure-guideline-levels-aegls-values#chemicals. March 22, 2016.

*AIHA. 2015. Current ERPG Values (2015). Fairfax, VA: American Industrial Hygiene Association. https://www.aiha.org/get-

involved/AIHAGuidelineFoundation/EmergencyResponsePlanningGuidelines/Documents/2015%20ERP G%20Levels.pdf. March 22, 2016.

+*Allen N, Mendell JR, Billmaier DJ, et al. 1975. Toxic polyneuropathy due to methyl n-butyl ketone: An industrial outbreak. Arch Neurol 32:209-218.

*Altman PL, Dittmer DS. 1974. Biological handbooks: Biology data book. Vol. III. 2nd ed. Bethesda, MD: Federation of American Societies of Experimental Biology, 1987-2008, 2041

*Ambrose D, Ellender JH, Lees EB, et al. 1975. Thermodynamic properties of organic oxygen compounds. XXXVIII. Vapour pressures of some aliphatic ketones. J Chem Thermodynamics 7:453-472.

*Andersen ME, Clewell HJ, Gargas ML, et al. 1987. Physiologically based pharmacokinetics and the risk assessment process for methylene chloride. Toxicol Appl Pharmacol 87(2):185-205.

*Andersen ME, Krishnan K. 1994. Relating *in vitro* to *in vivo* exposures with physiologically based tissue dosimetry and tissue response models. In: Salem H, ed. Animal test alternatives: Refinement, reduction, and replacement. New York, NY: Marcel Dekker, Inc., 9-25.

*Anderson RA, Harland WA. 1980. The analysis of volatiles in blood from fire fatalities. In: Proceedings of the Forensic Toxicology European Meeting. International Association of Toxicology. 279-292.

*Atkinson R. 1989. Kinetics and mechanisms of the gas-phase reactions of the hydroxyl radical with organic compounds. Journal of Physical and Chemical Reference Data. Monograph No. 1. American Chemical Society, American Institute of Physics. National Institute of Standards and Technology. http://www.nist.gov/data/PDFfiles/jpcrdM1.pdf. April 19, 2016.

*Atkinson R, Carter WP, Aschmann SM, et al. 1985. Atmospheric fates of organic chemicals: Prediction of ozone and hydroxyl radical reaction rates and mechanisms. Research Triangle Park, NC: U.S. Environmental Protection Agency, Office of Research and Development. EPA600385063. PB85241529.

*ATSDR. 1989. Decision guide for identifying substance-specific data needs related to toxicological profiles; Notice. Agency for Toxic Substances and Disease Registry. Fed Regist 54(174):37618-37634.

*ATSDR. 1999. Toxicological profile for n-hexane. Agency for Toxic Substances and Disease Registry, U.S. Department of Health and Human Services. http://www.atsdr.cdc.gov/toxprofiles/tp113.pdf. April 21, 2016. *ATSDR. 2008. Health consultation. Garfield county public health implications of ambient air exposure to volatile organic compounds as measured in rural, urban, and oil and gas development areas. Garfield County, Colorado. Agency for Toxic Substances & Disease Registry.

https://www.colorado.gov/pacific/sites/default/files/HHW_CSA_Ambient-Air-in-Garfield-County-HC-3.13.2008.pdf. August 4, 2017.

*ATSDR. 2015. 2-Hexanone. Full SPL data. Substance priority list (SPL) resource page. Agency for Toxic Substances and Disease Registry, Centers for Disease Control and Prevention. http://www.atsdr.cdc.gov/SPL/resources. November 1, 2016.

*Babeu L, Vaishnav DD. 1987. Prediction of biodegradability for selected organic chemicals. J Ind Microbial 2:107-115.

*Badings HT, De Jong C, Dooper RP. 1985. Automatic system for rapid analysis of volatile compounds by purge-and-cold trapping/capillary gas chromatography. J High Resolut Chromatogr Commun 8:755-763. 10.1002/jhrc.1240081111.

*Barnes DG, Dourson M. 1988. Reference dose (RfD): Description and use in health risk assessments. Regul Toxicol Pharmacol 8(4):471-486.

*Bavazzano P, Apostoli P, Balducci C, et al. 1998. Determination of urinary 2,5-hexanedione in the general Italian population. Int Arch Occup Environ Health 71(4):284-288.

*Berger GS, ed. 1994. Epidemiology of endometriosis. In: Endometriosis: Modern surgical management of endometriosis. New York, NY: Springer-Verlag, 3-7.

*Bierbaum PJ, Marceleno T. 1973. Survey of Brammer Manufacturing Company, Davenport, Iowa. Cincinnati, OH: National Institute for Occupational Safety and Health, Division of Field Studies and Clinical Investigations. PB81231318.

*Billmaier D, Yee HT, Allen N, et al. 1974. Peripheral neuropathy in a coated fabrics plant. J Occup Med 16:665-671.

+Branchflower RV, Pohl IX. 1981. Investigation of the mechanism of the potentiation of chloroforminduced hepatotoxicity and nephrotoxicity by methyl a-butyl ketone. Toxicol Appl Pharmacol 61:407-413.

+Branchflower RV, Schulick RD, George JW, et al. 1983. Comparison of the effects of methyl-n-butyl ketone and phenobarbital on rat liver cytochromes P-450 and the metabolism of chloroform to phosgene. Toxicol Appl Pharmacol 71:414-421.

+*Brown EM, Hewitt WR. 1984. Dose-response relationships in ketone-induced potentiation of chloroform hepato- and nephrotoxicity. Toxicol Appl Pharmacol 6:437-453.

*Brown KW, Donnelly KC. 1988. An estimation of the risk associated with the organic constituents of hazardous and municipal waste landfill leachates. Hazardous Wastes and Hazardous Materials 5:1-30.

*Bryant BH, Yamasaki H, Sandrof MA, et al. 2008. Spermatid head retention as a marker of 2,5-hexanedione-induced testicular toxicity in the rat. Toxicol Pathol 36:552-559.

*Buttery RG, Stern DJ, Ling LC. 1994. Studies on flavor volatiles of some sweet corn products. J Agric Food Chem 42:791-795. 10.1021/jf00039a038.

*Calvert JG, Pitts JN. 1966. Photochemistry of the polyatomic molecules. In: Photochemistry. New York, NY: John Wiley & Sons, Inc.

*Canter LW, Sabatini DA. 1994. Contamination of public ground water supplies by superfund sites. Int J Environ Stud 46:35-57. 10.1080/00207239408710909.

*Chou CHSJ, Holler J, De Rosa CT. 1998. Minimal risk levels (MRLs) for hazardous substances. J Clean Technol Environ Sci 7(1):1-24.

*Clewell HJ, Andersen ME. 1985. Risk assessment extrapolations and physiological modeling. Toxicol Ind Health 1(4):111-131.

*CLPSD. 1989, Contract Laboratory Program Statistical Database. Viar and Company, Management Services Division, Alexandria, VA. June 1986.

*Costa LG, Aschner M, Vitalone A, et al. 2004. Developmental neuropathology of environmental agents. Annu Rev Pharmacol Toxicol 44:87-110. 10.1146/annurev.pharmtox.44.101802.121424.

+*Couri D, Hetland LB, Abdel-Rahman MS, et al. 1977. The influence of inhaled ketone solvent vapors on hepatic microsomal biotransformation activities. Toxicol Appl Pharmacol 41:285-289.

+Cowlen MS, Hewitt WR, Schroeder F. 1984a. 2-Hexanone potentiation of [¹⁴C]chloroform hepatotoxicity: Covalent interaction of a reactive intermediate with rat liver phospholipid. Toxicol Appl Pharmacol 73:478-491.

+Cowlen MS, Hewitt WR, Schroeder F. 1984b. Mechanisms in 2-hexanone potentiation of chloroform hepatotoxicity. Toxicol Lett 22:293-299.

+*Davenport JG, Farrell DF, Sumi SM. 1976. Giant axonal neuropathy caused by industrial chemicals: Neurofilamentous axonal masses in man. Neurology 26:919923.

*Day EA, Anderson DF. 1965. Gas chromatographic and mass spectral identification of natural components of the aroma fraction of blue cheese. J Agric Food Chem 13:2-4.

*DeCaprio AP, Fowke JH. 1992. Limited and selective adduction of carboxyl-terminal lysines in the high molecular weight neurofilament proteins by 2,5-hexanedione *in vitro*. Brain Res 586:219-228.

*DeCaprio AP, Kinney EA, Fowke JH. 1997. Regioselective binding of 2,5-hexanedione to high-molecular-weight rat neurofilament proteins *in vitro*. Toxicol Appl Pharmacol 145:211-217.

*DHHS. 2016. CAS 591-78-6. 2-Hexanone. Household products database. U.S. Department of Health & Human Services, National Library of Medicine. https://householdproducts.nlm.nih.gov/. July 19, 2017.

*DiVincenzo GD, Hamilton ML, Kaplan CJ, et al. 1977. Metabolic fate and disposition of ¹⁴C-labeled methyl n-butyl ketone in the rat. Toxicol Appl Pharmacol 41:547-560.

*DiVincenzo GD, Hamilton ML, Kaplan CJ, et al. 1978. Studies on the respiratory uptake and excretion and the skin absorption of methyl n-butyl ketone in humans and dogs. Toxicol Appl Pharmacol 44:593-604.

*DiVincenzo GD, Kaplan CJ, Dedinas J. 1976. Characterization of the metabolites of methyl n-butyl ketone, methyl iso-butyl ketone, and methyl ethyl ketone in guinea pig serum and their clearance. Toxicol Appl Pharmacol 36:511-522.

*DOE. 2012. Protective action criteria (PAC): Chemicals with AEGLs, ERPGs, & TEELs. Definition of PACs (AEGLs, ERPGs or TEELs). Protective Action Criteria (PAC) with AEGLs, ERPGs, & TEELs: Rev. 27 for Chemicals of Concern - March 2012. Oak Ridge, TN: U.S. Department of Energy. http://energy.gov/ehss/protective-action-criteria-pac-aegls-erpgs-teels-rev-27-chemicals-concern-march-2012. March 24, 2015.

*DOE. 2016. Table 3: Protective Action Criteria (PAC) Rev. 28A based on applicable 60-minute AEGLs, ERPGs, or TEELs. The chemicals are listed by CASRN. February 2016. Oak Ridge, TN: U.S. Department of Energy. http://www.atlintl.com/DOE/teels/teel/Revision_28A_Table3.pdf. March 22, 2016.

+*Duckett S, Streletz IJ, Chambers RA, et al. 1979. 50 ppm MnBK subclinical neuropathy in rats. Experientia 35:1365-1367.

*Duguay AB, Plaa GL. 1995. Tissue concentrations of methyl isobutyl ketone, methyl n-butyl ketone and their metabolites after oral or inhalation exposure. Toxicol Lett 75(1-3):51-58.

*Dumont JP, Adda J. 1978. Occurrence of sesquiterpenes in mountain cheese volatiles. J Agric Food Chem 26:364-367.

+*Eben A, Flucke W, Mihail F, et al. 1979. Toxicological and metabolic studies of methyl nbutylketone, 2,5-hexanedione, and 2,5-hexanediol in male rats. Ecotoxicol Environ Safety 3:204-217.

+*Egan G, Spencer P, Schaumburg H, et al. 1980. n-Hexane-"free" hexane mixture fails to produce nervous system damage. Neurotoxicology 1:515-524.

*EIA. 2016. Energy in brief. Shale in the United States. U.S. Energy Information Administration. https://www.eia.gov/energy_in_brief/article/shale_in_the_united_states.cfm. November 02, 2016.

*Eitzer BD. 1995. Emissions of volatile organic chemicals from municipal solid waste composting facilities. Environ Sci Technol 29:896-902. 10.1021/es00004a009.

*Ek CJ, Dziegielewska KM, Habgood MD, et al. 2012. Barriers in the developing brain and neurotoxicology. Neurotoxicology 33(3):586-604. 10.1016/j.neuro.2011.12.009.

*EPA. 1981. Chemical hazard information profile Draft report methyl n-butyl ketone. Washington, DC: U.S. Environmental Protection Agency, Office of Pesticides and Toxic Substances.

*EPA. 1986. Gas chromatography/mass spectrometry for volatile organics method 8240. In: Test methods for evaluating solid waste. 3rd ed. SW-846. Washington, DC: U.S. Environmental Protection Agency, Office of Solid Waste and Emergency Response.

*EPA. 1987. U.S. Environmental Protection Agency. Fed Regist 52:11823-11826.

*EPA. 1988. Superfund record of decision. Forest Waste Disposal. Otisville, Michigan. Second remedial action. Washington, DC: U.S. Environmental Protection Agency. EPARODR0588062. http://nepis.epa.gov/Exe/ZyPURL.cgi?Dockey=91001K2I.txt. April 21, 2016.

*EPA. 1990. Interim methods for development of inhalation reference concentrations. Washington, DC: U.S. Environmental Protection Agency, Office of Health and Environmental Assessment, Office of Research and Development. EPA600890066A. PB90238890.

*EPA. 1997. Special report on environmental endocrine disruption: An effects assessment and analysis. Washington, DC: U.S. Environmental Protection Agency, Office of Pollution Prevention and Toxics. EPA630R96012.

*EPA. 2005. Toxic chemical release inventory reporting forms and instructions: Revised 2004 version. Section 313 of the Emergency Planning and Community Right-to-Know Act (Title III of the Superfund Amendments and Reauthorization Act of 1986). U.S. Environmental Protection Agency, Office of Environmental Information. EPA260B05001.

*EPA. 2009a. Toxicological review of 2-hexanone (CAS No. 591-78-6). U.S. Environmental Protection Agency. http://cfpub.epa.gov/ncea/iris/iris_documents/documents/toxreviews/1019tr.pdf. January 22, 2016.

*EPA. 2009b. National primary drinking water regulations. Washington, DC: U.S. Environmental Protection Agency, Office of Ground Water and Drinking Water. EPA816F090004. http://water.epa.gov/drink/contaminants/upload/mcl-2.pdf. March 4, 2015.

*EPA. 2012a. Benchmark dose technical guidance. Washington, DC: U.S. Environmental Protection Agency. EPA100R12001.

*EPA. 2012b. 2012 Edition of the drinking water standards and health advisories. Washington, DC: U.S. Environmental Protection Agency, Office of Water. EPA822S12001. https://www.epa.gov/sites/production/files/2015-09/documents/dwstandards2012.pdf. April 21, 2016.

*EPA. 2012c. 2-Hexanone. EPI SuiteTM-Estimation program interface. https://www.epa.gov/tsca-screening-tools/epi-suitetm-estimation-program-interface. June 14, 2016.

*EPA. 2014. Title 42 - The public health and welfare. Chapter 85 - Air pollution prevention and control. Subchapter I - programs and activities. Part A - Air quality and emission limitations. Hazardous air pollutants. U.S. Environmental Protection Agency. United States Code 42 USC 7412. https://www.gpo.gov/fdsys/pkg/USCODE-2014-title42/pdf/USCODE-2014-title42-chap85-subchapI-partA-sec7412.pdf. April 21, 2015.

*EPA. 2015a. Subchapter D-water programs. Designation of hazardous substances. U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 116.4. https://www.gpo.gov/fdsys/pkg/CFR-2015-title40-vol22/pdf/CFR-2015-title40-vol22-sec116-4.pdf. April 21, 2016.

*EPA. 2015b. Subpart A - General provisions. Determination of reportable quantities. U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 117.3. https://www.gpo.gov/fdsys/pkg/CFR-2015-title40-vol22/pdf/CFR-2015-title40-vol22-sec117-3.pdf. April 21, 2015. *EPA. 2015c. Subchapter J-Superfund, emergency planning, and community right-to-know programs. Designation of hazardous substances. U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 302.4. https://www.gpo.gov/fdsys/pkg/CFR-2015-title40-vol28/pdf/CFR-2015-title40-vol28/pdf/CFR-2015-title40-vol28-part302.pdf. April 21, 2016.

*EPA. 2015d. Subpart D - Specific toxic chemical listings. Chemicals and chemical categories to which this part applies. U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 372.65. https://www.gpo.gov/fdsys/pkg/CFR-2015-title40-vol28/pdf/CFR-2015-title40-vol28-sec372-65.pdf. April 21, 2016.

*EPA. 2015e. Subpart B - Manufacturers reporting - preliminary assessment information. Chemical lists and reporting periods. U.S. Environmental Protection Agency. Code of Federal Regulations 40 CFR 712.30. https://www.gpo.gov/fdsys/pkg/CFR-2015-title40-vol31/pdf/CFR-2015-title40-vol31-sec712-30.pdf. April 21, 2016.

*EPA. 2016a. National recommended water quality criteria - Aquatic life criteria table Washington, DC: U.S. Environmental Protection Agency, Office of Water, Office of Science and Technology. https://www.epa.gov/wqc/national-recommended-water-quality-criteria-aquatic-life-criteria-table. March 22, 2016.

*EPA. 2016b. National recommended water quality criteria - Human health criteria table. Washington, DC: U.S. Environmental Protection Agency, Office of Water, Office of Science and Technology. https://www.epa.gov/wqc/national-recommended-water-quality-criteria-human-health-criteria-table. March 22, 2016.

*FDA. 2013. Everything added to food in the United States (EAFUS). Washington, DC: U.S. Food and Drug Administration. http://www.accessdata.fda.gov/scripts/fcn/fcnnavigation.cfm?rpt=eafuslisting. January 8, 2014.

*Fedtke N, Bolt HM. 1986. Methodological investigations on the determination of n-hexane metabolites in urine. Int Arch Occup Environ Health 57:149-158.

*Fomon SJ. 1966. Body composition of the infant: Part 1: The male reference infant. In: Faulkner F, ed. Human development. Philadelphia, PA: WB Saunders, 239-246.

*Fomon SJ, Haschke F, Ziegler EE, et al. 1982. Body composition of reference children from birth to age 10 years. Am J Clin Nutr 35(Suppl 5):1169-1175.

*Girolami RL, Knight SG. 1955. Fatty acid oxidation by *Penicillium roqueforti*. Appl Microbiol 3(5):264-267.

*Giwercman A, Carlsen E, Keiding N, et al. 1993. Evidence for increasing incidence of abnormalities of the human testis: A review. Environ Health Perspect 101(Supp 2):65-71.

*Granvil CP, Sharkawi M, Plaa GL. 1994. Metabolic fate of methyl n-butyl ketone, methyl isobutyl ketone and their metabolites in mice. Toxicol Lett 70(3):263-267.

*Green DR, Le Pape D. 1987. Stability of hydrocarbon samples on solid-phase extraction columns. Anal Chem 59:699-703.

*Grey TC, Shrimpton DH. 1967. Volatile components of raw chicken breast muscle. Br Poult Sci 8:23-33.

*Grinsberg A. 2014. In the pits. Oil and gas wastewater disposal into open unlined pits and the threat to California's water and air. Clean Water Action, Clean Water Fund. http://www.cleanwateraction.org/sites/default/files/docs/publications/In%20the%20Pits.pdf. November 7, 2017.

*Gummin DD. 2015. Hydrocarbons. In: Hoffman RS, Lewin NA, Goldfrank LR, et al., eds. Goldfrank's toxicologic emergencies. 10th ed. New York, NY: McGraw-Hill Education, 1334-1345.

*Guzelian PS, Henry CJ, Olin SS. 1992. Similarities and differences between children and adults: Implications for risk assessment. Washington, DC: International Life Sciences and Press Institute Press.

*Hansch C, Leo A, Hoekman D. 1995. 2-Hexanone. In: Exploring QSAR. Hydrophobic, electronic, and steric constants. Washington, DC: American Chemical Society, 24.

+*Hansen LF, Nielsen GD. 1994. Sensory irritation and pulmonary irritation of n-methyl ketones: Receptor activation mechanisms and relationships with threshold limit values. Arch Toxicol 68(3):193-202.

*Hawthorne SB, Sievers RE. 1984. Emission of organic air pollutants from shale oil wastewaters. Environ Sci Technol 18(6):483-490. 10.1021/es00124a016.

*Hawthorne SB, Sievers RE, Barkley RM. 1985. Organic emissions from shale oil wastewaters and their implications for air quality. Environ Sci Technol 19:992-997.

*Hayes T. 2009. Sampling and analysis of water streams associated with the development of Marcellus shale gas. Gas Technology Institute. http://energyindepth.org/wp-content/uploads/marcellus/2012/11/MSCommission-Report.pdf. August 17, 2017.

*Helmig D, Arey J. 1992. Organic chemicals in the air at Whitaker's Forest/ Sierra Nevada Mountains, California. Sci Total Environ 112(2-3):233-250. 10.1016/0048-9697(92)90190-4

+Hewitt LA, Ayotte P, Plaa GL. 1986. Modifications in rat hepatobiliary function following treatment with acetone, 2-butanone, 2-hexanone, mirex, or chlordecone and subsequently exposed to chloroform. Toxicol Appl Pharmacol 83:465-473.

+Hewitt WR, Brown EM, Plaa GL. 1983. Relationship between the carbon skeleton length of ketonic solvents and potentiation of chloroform-induced hepatotoxicity in rats. Toxicol Lett 16:297-304.

+Hewitt WR, Miyajima H, Cote MG, et al. 1980. Acute alteration of chloroform- induced hepato- and nephrotoxicity by n-hexane, methyl n-butyl ketone and 2,5-hexanedione. Toxicol Appl Pharmacol 53:230-248.

*Hoel DG, Davis DL, Miller AB, et al. 1992. Trends in cancer mortality in 15 industrialized countries, 1969-1986. J Natl Cancer Inst 84(5):313-320.

*HSDB. 2009. 2-Hexanone. Hazardous Substances Data Bank. National Library of Medicine. http://toxnet.nlm.nih.gov. April 19, 2016.

*IARC. 2016. Agents classified by the IARC monographs. Volumes 1–115. Lyon, France: International Agency for Research on Cancer. http://monographs.iarc.fr/ENG/Classification/List_of_Classifications_Vol1-115.pdf. March 22, 2016.

*IRIS. 2009. 2-Hexanone; CASRN 591-78-6. Integrated Risk Information System. Washington, DC: U.S. Environmental Protection Agency. https://cfpub.epa.gov/ncea/iris/iris_documents/documents/subst/1019_summary.pdf. March 22, 2016.

+*Johnson BL, Setzer JV, Lewis TR, et al. 1977. Effects of methyl n-butyl ketone on behavior and the nervous system. Am Ind Hyg Assoc J 38:567-579.

+Kanada M, Miyagawa M, Sato M, et al. 1994. Neurochemical profile of effects of 28 neurotoxic chemicals on the central nervous system in rats: 1. Effects of oral administration on brain contents of biogenic amines and metabolites. Ind Health 32(3):145-164.

*Katz GV, O'Donoghue JL, DiVincenzo GD, et al. 1980. Comparative neurotoxicity and metabolism of ethyl n-butyl ketone and methyl n-butyl ketone in rats. Toxicol Appl Pharmacol 52:153-158.

*Kearns GL, Abdel-Rahman SM, Alander SW, et al. 2003. Developmental pharmacology-drug disposition, action, and therapy in infants and children. N Engl J Med 349(12):1157-1167. 10.1056/NEJMra035092.

*King JW. 1989. Fundamentals and applications of supercritical fluid extraction in chromatographic science. J Chromatogr Sci 27:355-364.

*Kinlin TE, Muralidhara R, Pittet AO, et al. 1972. Volatile components of roasted filberts. J Agric Food Chem 20:1021-1028.

*Komori M, Nishio K, Kitada M, et al. 1990. Fetus-specific expression of a form of cytochrome P-450 in human livers. Biochemistry 29(18):4430-4433.

+*Krasavage WJ, O'Donoghue JL. 1977. Chronic inhalation exposure of rats to methyl n-butyl ketone (MnBK). In: Initial Submission: Letter from Eastman Kodak Co to USEPA regarding toxicity studies of 2-hexanone and metabolites with cover letter dated 09/28/92. Eastman Kodak Co. Submitted under TSCA Section 8E to U.S. Environmental Protection Agency. OTS0555051. [Unpublished study to be peer reviewed]

+*Krasavage WJ, O'Donoghue JL, DiVincenzo GD, et al. 1980. The relative neurotoxicity of methyl-nbutyl ketone, n-hexane and their metabolites. Toxicol Appl Pharmacol 52:433-441.

*Krishnan K, Andersen ME. 1994. Physiologically based pharmacokinetic modeling in toxicology. In: Hayes AW, ed. Principles and methods of toxicology. 3rd ed. New York, NY: Raven Press, Ltd., 149-188.

*Krishnan K, Anderson ME, Clewell HJ, et al. 1994. Physiologically based pharmacokinetic modeling of chemical mixtures. In: Yang RSH, ed. Toxicology of chemical mixtures. Case studies, mechanisms, and novel approaches. San Diego, CA: Academic Press, 399-437.

*Kumar A, Alaimo CP, Horowitz R, et al. 2011. Volatile organic compound emissions from green waste composting: Characterization and ozone formation. Atmos Environ 45:1841-1848.

*Laity JL, Burstain IG, Appel BR. 1973. Photochemical smog and the atmospheric reactions of solvents. In: Tess RW, ed. Solvents theory and practice. Advances in chemistry series 124. Washington, DC: American Chemical Society, 95-112.

*Lande SS, Durkin PR, Christopher DE, et al. 1976. Investigation of selected potential environmental contaminants: Ketonic solvents. Washington, DC: U.S. Environmental Protection Agency, Office of Toxic Substances. EPA560276003. NTIS No. TR 76-500.

*Lapadula DM, Habig C, Gupta RP, et al. 1991. Induction of cytochrome P450 isozymes by simultaneous inhalation exposure of hens to n-hexane and methyl iso-butyl ketone (MiBK). Biochem Pharmacol 6(7):877-883.

*Lee EH, Kim J, Cho KS, et al. 2010. Degradation of hexane and other recalcitrant hydrocarbons by a novel isolate, Rhodococcus sp. EH831. Environ Sci Pollut Res Int 17(1):64-77. 10.1007/s11356-009-0238-x.

*Leeder JS, Kearns GL. 1997. Pharmacogenetics in pediatrics: Implications for practice. Pediatr Clin North Am 44(1):55-77.

*Leung H. 1993. Physiologically-based pharmacokinetic modelling. In: Ballantyne B, Marrs T, Turner P, eds. General and applied toxicology. Vol. 1. New York, NY: Stockton Press, 153-164.

*Livingston AL. 1978. Forage plant estrogens. J Toxicol Environ Health 4(2-3):301-324.

*LoPachin RM, DeCaprio AP. 2004. γ-Diketone neuropathy: Axon atrophy and the role of cytoskeletal protein adduction. Toxicol Appl Pharmacol 199(1):20-34. 10.1016/j.taap.2004.03.008.

*LoPachin RM, DeCaprio AP. 2005. Protein adduct formation as a molecular mechanism in neurotoxicity. Toxicol Sci 86(2):214-225.

*LoPachin RM, Gavin T. 2015. Toxic neuropathies: Mechanistic insights based on a chemical perspective. Neurosci Lett 596:78-83. 10.1016/j.neulet.2014.08.054.

*LoPachin RM, He D, Reid ML. 2005b. 2,5-Hexanedione-induced changes in the neurofilament subunit pools of rat peripheral nerve. Neurotoxicology 26(2):229-240. 10.1016/j.neuro.2004.09.007.

*LoPachin RM, Jortner BS, Reid ML, et al. 2003. γ -Diketone central neuropathy: Quantitative morphometric analysis of axons in rat spinal cord white matter regions and nerve roots. Toxicol Appl Pharmacol 193(1):29-46.

*Lopachin RM, Jortner BS, Reid ML, et al. 2005a. γ -Diketone central neuropathy: Quantitative analyses of cytoskeletal components in myelinated axons of the rat rubrospinal tract. Neurotoxicology 26(6):1021-1030. 10.1016/j.neuro.2005.04.008.

*LoPachin RM, Lehning EJ, Opanashuk LA, et al. 2000. Rate of neurotoxicant exposure determines morphologic manifestations of distal axonopathy. Toxicol Appl Pharmacol 167(2):75-86. 10.1006/taap.2000.8984.

*Lowery CE Jr., Foster JW, Jurtshuk P. 1968. The growth of various filamentous fungi and yeasts on nalkanes and ketones: I. Studies on substrate specificity. Arch Mikrobiol 60:246-254. *Lucas SV. 1984. GC/MS analysis of organics in drinking water concentrates and advanced waste treatment concentrates. Vol. 1. Analysis results for 17 drinking water, 16 advanced waste treatment and 3 process blank concentrates. Report to U.S. Environmental Protection Agency, Office of Research and Development, Research Triangle Park, NC, Battelle Columbus Laboratories, Columbus, OH. EPA-600184020a. PB85128221.

*Lukins HB, Foster JW. 1963. Methyl ketone metabolism in hydrocarbon utilizing mycobacteria. J Bacterial 85:1074-1087.

*Luster MI, Munson AE, Thomas PT, et al. 1988. Development of a testing battery to assess chemicalinduced immunotoxicity: National Toxicology Program's guidelines for immunotoxicity evaluation in mice. Fundam Appl Toxicol 10(2-19).

*MacLeod H, Jourdain JL, Poulet G, et al. 1984. Kinetic study of reactions of some organic sulfur compounds with OH radicals. Atmos Environ 18:2621-2626.

*Mallov JS. 1976. MBK neuropathy among spray painters. J Am Med Assoc 235:1455-1457.

*Marceleno T, Bierbaum PJ, Mallov JS. 1974. Survey of Premium Finishes, Incorporated, Cincinnati, OH. Cincinnati, OH: National Institute for Occupational Safety and Health, Division of Field Studies and Clinical Investigations. PB81242729.

*Mayr U, Butsch A, Schneider S. 1992. Validation of two *in vitro* test systems for estrogenic activities with zearalenone, phytoestrogens and cereal extracts. Toxicology 74(2-3):135-149.

*McCormick RL, Ratcliff MA, Christensen E, et al. 2015. Properties of oxygenates found in upgraded biomass pyrolysis oil as components of spark and compression ignition engine fuels. American Chemical Society. Energy Fuels 29(4):2453-2461. 10.1021/ef502893g.

+Mendell JR, Sahenk Z, Saida K, et al. 1977. Alterations of fast axoplasmic transport in experimental methyl n-butyl ketone neuropathy. Brain Res 133:107118.

+*Mendell JR, Saida K, Ganansia MF, et al. 1974. Toxic polyneuropathy produced by methyl n-butyl ketone. Science 185:787-789.

*Morselli PL, Franco-Morselli R, Bossi L. 1980. Clinical pharmacokinetics in newborns and infants: Age-related differences and therapeutic implications. Clin Pharmacokinet 5(6):485-527.

*Myers VB. 1983. Remedial activities at the Miami Drum site, Florida. In: National conference on management of uncontrolled hazardous waste sites. Silver Springs, MD: Hazardous Materials Control Research Institute, 354-357.

*Nakajima D, Ishii R, Kageyama S, et al. 2006. Genotoxicity of microbial volatile organic compounds. J Health Sci 52:148-153.

*NAS/NRC. 1989. Report of the oversight committee. Biologic markers in reproductive toxicology. Washington, DC: National Academy of Sciences, National Research Council, National Academy Press, 15-35.

*NIOSH. 1978. Criteria for a recommended standard: Occupational exposure in coal gasification plants. Cincinnati, OH: National Institute for Occupational Safety and Health. DHEW (NIOSH) Pub No. 78-191.

*NIOSH. 1984. Ketones I - method 1300. In: NIOSH manual of analytical methods. 3rd ed. Vol. 2. Cincinnati, OH: U.S. Department of Health and Human Services, National Institute for Occupational Safety and Health.

*NIOSH. 2003. Ketones I. Method 2555. NIOSH Manual of Analytical Methods (NMAM). National Occupational Safety and Health. http://www.cdc.gov/niosh/docs/2003-154/pdfs/2555.pdf. April 20, 2016.

*NIOSH. 2014. 2-Hexanone. Immediately Dangerous to Life or Health Concentrations (IDLH). Atlanta, GA: National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention. http://www.cdc.gov/niosh/idlh/591786.html. March 22, 2016.

*NIOSH. 2015. 2-Hexanone. NIOSH pocket guide to chemical hazards. Atlanta, GA: National Institute for Occupational Safety and Health, Centers for Disease Control and Prevention. http://www.cdc.gov/niosh/npg/npgd0325.html. March 22, 2016.

*Nomeir AA, Abou-Donia MB. 1985. Analysis of D-hexane, 2-hexanone, 2,5-hexanedione, and related chemicals by capillary gas chromatography and high performance liquid chromatography. Anal Biochem 151:381-388.

*NRC. 1993. Pesticides in the diets of infants and children. Washington, DC: National Research Council. National Academy Press. PB93216091.

*NTP. 2014. Report on carcinogens. Thirteenth edition. Research Triangle Park, NC: U.S. Department of Health and Human Services, Public Health Service, National Toxicology Program. http://ntp.niehs.nih.gov/pubhealth/roc/roc13/. March 22, 2016.

+*O'Donoghue JL, Krasavage WJ. 1979. Chronic inhalation exposure of cats to methyl n-butyl ketone (MnBK). In: Initial Submission: Letter from Eastman Kodak Co to USEPA regarding toxicity studies of 2-hexanone and metabolites with cover letter dated 09/28/92. Eastman Kodak Co. Submitted under TSCA Section 8E to U.S. Environmental Protection Agency. OTS0555051. [Unpublished study to be peer reviewed]

+*O'Donoghue JL, Krasavage WJ, Terhaar CJ. 1978. A comparative chronic toxicity study of methyl npropyl ketone, methyl n-butyl ketone, and hexane by ingestion. In: Initial Submission: Letter from Eastman Kodak Co to USEPA regarding toxicity studies of 2-hexanone and metabolites with cover letter dated 09/28/92. Eastman Kodak Co. Submitted under TSCA Section 8E to U.S. Environmental Protection Agency. OTS0555051. [Unpublished study to be peer reviewed]

*OSHA. 1995. 2-Hexanone. Method number OV2013. Occupational Safety & Health Administration, United States Department of Labor. https://www.osha.gov/dts/sltc/methods/partial/pv2031/2031.html. April 18, 2016.

*OSHA. 2015a. Gases, vapors, fumes, dusts, and mists. Appendix A to Part 1926.55-1970 American Conference of Governmental Industrial Hygienists' threshold limit values of airborne contaminants. Occupational Safety and Health Standards. Code of Federal Regulations 29 CFR 1926.55. https://www.gpo.gov/fdsys/pkg/CFR-2015-title29-vol8/pdf/CFR-2015-title29-vol8-sec1926-55.pdf. April 21, 2016.

*OSHA. 2015b. Subpart Z - Toxic and hazardous substances. Air contaminants. Occupational Safety and Health Standards. Code of Federal Regulations 29 CFR 1910.1000. https://www.gpo.gov/fdsys/pkg/CFR-2015-title29-vol6/pdf/CFR-2015-title29-vol6-sec1910-1000.pdf. April 21, 2016.

*OSHA. 2015c. Subpart Z - Toxic and hazardous substances. Air contaminants. Table Z - Shipyards. Occupational Safety and Health Standards. Code of Federal Regulations 29 CFR 1915.1000. https://www.gpo.gov/fdsys/pkg/CFR-2015-title29-vol7/pdf/CFR-2015-title29-vol7-sec1915-1000.pdf. April 21, 2016.

*Owen GM, Brozek J. 1966. Influence of age, sex and nutrition on body composition during childhood and adolescence. In: Falkner F, ed. Human development. Philadelphia, PA: WB Saunders, 222-238.

*PA DEP. 2011. Northeastern Pennsylvania Marcellus shale short-term ambient air sampling report. Pennsylvania Department of Environmental Protection. http://www.dep.state.pa.us/dep/deputate/airwaste/aq/aqm/docs/Marcellus_NE_01-12-11.pdf. August 17, 2017.

*Pankow JF, Rosen ME. 1988. Determination of volatile compounds in water by purging directly to a capillary column with whole column cryotrapping. Environ Sci Technol 22:398-405.

*Pellizzari ED, Castillo NP, Willis S, et al. 1979. Identification of organic components in aqueous effluents from energy-related processes. In: Van Hall CE, ed. Measurement of organic pollutants in water and wastewater. Philadelphia, PA: American Society for Testing and Materials, 256-274. ASTM Special Technical Publication No. 686.

*Perry JJ. 1968. Substrate specificity in hydrocarbon utilizing microorganisms. Antonie Van Leeuwenhoek 34:27-36.

*Perry DL, Chuang CC, Jungclaus GA, et al. 1979. Identification of organic compounds in industrial effluent discharges. Athens, GA: U.S. Environmental Protection Agency, Office of Research and Development, Environmental Research Laboratory. EPA600479016. PB294794.

+*Peters MA, Hudson PM, Dixon RL. 1981. The effect totigestational exposure to methyl n-butyl ketone has on postnatal development and behavior. Ecotoxicol Environ Safety 5:291-306.

*Plumb RHJ. 1992. The importance of volatile organic compounds as a disposal site monitoring parameter. In: Environmental science pollution control series 4, groundwater contamination and analysis at hazardous waste sites. New York, NY: Mercel Dekker, Inc, 173-197.

*Ramarathnam N, Rubin LJ, Diosady LL. 1991. Studies on meat flavor. 2. A quantitative investigation of the volatile carbonyls and hydrocarbons in uncured and cured beef and chicken. J Agric Food Chem 39:1839-1847.

*Raymond P, Plaa GL. 1995. Ketone potentiation of haloalkane-induced hepato- and nephrotoxicity. II. Implication of monooxygenases. J Toxicol Environ Health 46(3):317-328.

*RePORTER. 2016. 2-Hexanone. National Institutes of Health, Research Portfolio Online Reporting Tools. http://projectreporter.nih.gov/reporter.cfm. March 21, 2016.

*Richardson SD, Thruston ADJ, Caughran TV, et al. 1999. Identification of new ozone disinfection byproducts in drinking water. Environ Sci Technol 33:3368-3377. 10.1021/es981218c.

*RPSEA. 2012. Barnett and Appalachian shale water management and reuse technologies. Research Partnership to Secure Energy for America. Report No. 08122-05. Final.1. https://www.netl.doe.gov/File%20Library/Research/Oil-Gas/Natural%20Gas/shale%20gas/08122-05-final-report.pdf. August 11, 2017.

*RTECS. 2009. 2-Hexanone. Registry of Toxic Effects of Chemical Substances. National Institute for Occupational Safety and Health. http://www.cdc.gov/niosh-rtecs/MP155CC0.html. March 7, 2016.

+*Saida K, Mendell JR, Weiss HS. 1976. Peripheral nerve changes induced by methyl E-butyl ketone and potentiation by methyl ethyl ketone. Neuropathol Exp Neurol 35:207-225.

*Sato A, Nakajima T. 1979. Partition coefficients of some aromatic hydrocarbons and ketones in water, blood and oil. Br J Ind Med 36:231-234.

*Saunders NR, Ek CJ, Habgood MD, et al. 2008. Barriers in the brain: A renaissance? Trends Neurosci 31(6):279-286. 10.1016/j.tins.2008.03.003.

*Saunders NR, Liddelow SA, Dziegielewska KM. 2012. Barrier mechanisms in the developing brain. Front Pharmacol 3:Article 46. 10.3389/fphar.2012.00046.

*Scheuplein R, Charnley G, Dourson M. 2002. Differential sensitivity of children and adults to chemical toxicity. I. Biological basis. Regul Toxicol Pharmacol 35(3):429-447.

+*Schrenk HH, Yant WP, Patty FA. 1936. Acute response of guinea pigs to vapors of some new commercial organic compounds. Public Health Rep 51:624-631.

*Shackelford WM, Keith LH. 1976. Frequency of organic compounds identified in water. Athens, GA: U.S. Environmental Protection Agency, Office of Research and Development. EPA600476062. PB165470.

+*Smyth HF, Carpenter CP, Weil CS, et al. 1954. Range-finding toxicity data: List V. Arch Ind Hyg Occup Med 10:61-68.

+*Spencer PS, Schaumburg HH, Raleigh RL, et al. 1975. Nervous system degeneration produced by the industrial solvent methyl n-butyl ketone. Arch Neurol 32:219-222.

*Stutz DR, Janusz SJ. 1988. Protocol 56 ketones. In: Hazardous materials injuries: A handbook for pre-hospital care. 2nd ed. Beltsville, MD: Bradford Communications Corporation, 312-313.

*Takeoka GR, Flath RA, Guntert M, et al. 1988. Nectarine volatiles: Vacuum steam distillation versus headspace sampling. J Agric Food Chem 36:553-560.

*TCEQ. 2011. Superfund site discovery and assessment program. Removal action report. Dona Park residential removal site. Corpus Christi, Nueces County, TX. Houston, TX: Texas Commission on Environmental Quality.

https://www.tceq.texas.gov/assets/public/remediation/variousremediationsites/donapark/dona_park_rar_fi nal.pdf. August 10, 2016.

*TERA. 2014. Workplace Environmental Exposure Levels (WEEL). Cincinnati, OH: Toxicology Excellence for Risk Assessment. Occupational Alliance for Risk Science. http://www.tera.org/OARS/WEELs.pdf. March 22, 2016.

*Thomas K, Colborn T. 1992. Organochlorine endocrine disruptors in human tissue. In: Colborn T, Clement C, eds. Chemically induced alterations in sexual and functional development: The wildlife/human connection. Princeton, NJ: Princeton Scientific Publishing, 365-394.

*Thomas RG. 1990. Volatilization from water. In: Lyman WJ, Reehl WF, Rosenblatt DH, eds. Handbook of chemical property estimation methods. Washington, DC: American Chemical Society, 15-11-15-34.

*Topping DC, Morgott DA, O'Donoghue JL. 2001. Ketones of six to thirteen carbons. In: Bingham E, Cohrssen B, Powell CH, eds. Patty's toxicology. Vol. 6. 5th ed. New York, NY: John Willey & Sons, Inc., 195-363.

+*Union Carbide. 1977. Comparative toxicity to rats of methoxyacetone and five other aliphatic ketones in their drinking water with cover letter. Union Carbide Corporation. Submitted to the U.S. Environmental Protection Agency under TSCA Section 8D. Union Carbide Corporation. OTS206068.

*Vieira I, Sonnier M, Cresteil T. 1996. Developmental expression of CYP2E1 in the human liver: Hypermethylation control of gene expression during the neonatal period. Eur J Biochem 238(2):476-483.

*West JR, Smith HW, Chasis H. 1948. Glomerular filtration rate, effective renal blood flow, and maximal tubular excretory capacity in infancy. J Pediatr 32:10-18.

*White EL, Bus JS, Heck HD. 1979. Simultaneous determination of n-hexane, 2-hexanone and 2,5-hexanedione in biological tissues by gas chromatography mass spectrometry. Biomed Mass Spectrum 6(4):169-172.

*WHO. 2010. WHO guidelines for indoor air quality: Selected pollutants. Geneva, Switzerland: World Health Organization. http://www.euro.who.int/__data/assets/pdf_file/0009/128169/e94535.pdf. September 9, 2014.

*WHO. 2011. Guidelines for drinking-water quality. Geneva, Switzerland: World Health Organization. http://whqlibdoc.who.int/publications/2011/9789241548151_eng.pdf?ua=1. September 9, 2014.

*Widdowson EM, Dickerson JWT. 1964. Chemical composition of the body. In: Comar CL, Bronner F, eds. Mineral metabolism: An advance treatise. Volume II: The elements Part A. New York, NY: Academic Press, 1-247.

*Wieboldt RC, Adams GE, Later DW. 1988. Sensitivity improvement in infrared detection for supercritical fluid chromatography. Anal Chem 60:2422-2427.

*Yalkowsky SH, He Y. 2003. 2-Hexanone. In: Handbook of aqueous solubility data. Boca Raton, FL: CRC Press, 301.

*Yu RC, Hattis D, Landaw EM, et al. 2002. Toxicokinetic interaction of 2,5-hexanedione and methyl ethyl ketone. Arch Toxicol 75:643-652.

*Zhang L, Gavin T, DeCaprio AP, et al. 2010. γ -Diketone axonopathy: Analyses of cytoskeletal motors and highways in CNS myelinated axons. Toxicol Sci 117(1):180-189. 10.1093/toxsci/kfq176.

*Ziegler EE, Edwards BB, Jensen RL, et al. 1978. Absorption and retention of lead by infants. Pediatr Res 12(1):29-34.